

MALARIA CONTROL
BY ANTI-MOSQUITO MEASURES

Printed by Thacker's Press and Directories, Ltd.,
6, Bentinck Street, Calcutta
Published by Thacker Spink & Co (1933), Ltd.,
3, Esplanade East Calcutta

All rights reserved.

MALARIA CONTROL BY ANTI-MOSQUITO MEASURES

BY

GORDON COVELL

CIE MD (Lond) DPH DTM & H (Eng)

Lieut Colonel Indian Medical Service

Director Malaria Institute of India

SECOND EDITION

Revised enlarged and brought up to date by the Author

1941

CALCUTTA

THACKER SPINK & CO (1933) LTD

LONDON

W THACKER & CO 34 40, Ludgate Hill

PREFACE TO SECOND EDITION

In this edition a new chapter has been added dealing with the more important advances which have been made in malaria control methods during the past ten years. The various subjects have been treated in the same order as in the original edition.

The literature relating to anti mosquito measures has become very extensive and the bibliography, which is believed to be reasonably complete now includes 1,148 references as against 570 in the first edition. In cases where the title is not sufficiently explanatory, the subject is indicated by a footnote placed immediately below the reference.

January 1941

PREFACE TO FIRST EDITION

SINCE the immense importance of the rôle played by mosquitoes in the transmission of disease has come to be generally recognized, the literature dealing with the application of various anti mosquito measures has become so extensive that it is difficult for the average malaria worker to keep in touch with the different methods advocated. In the following pages an endeavour has been made to give some account of these methods in a convenient form.

It is impossible in a work of this kind to deal really adequately with the subject of drainage, which is largely a matter for the skilled engineer, but a brief description has been given of most of the points of practical importance to the malariologist. Other important measures, such as screening, oiling and the use of paris green, have been treated in considerable detail, whilst an attempt has been made to make some reference to all the measures which have been advocated in the literature.

An important feature of the book is the bibliography which contains 570 references to books and articles dealing with the subject. These have been arranged for convenience under the headings of the various measures dealt with. A list of the sources of various apparatus used in the application of anti mosquito measures is given as an appendix.

It is hoped that the description of the various measures which have proved of practical importance will make the book of value to the man in the field, and that the bibliography will be of service to malariologists generally

I wish to express my thanks to Dr G Macdonald, of the Malaria Survey of India*, for many helpful suggestions, and for kindly taking the photographs which are reproduced in Plates I and II

May 1931

* This organization is now designated The Malaria Institute of India

CONTENTS

Part I

PROTECTION AGAINST BITES OF MOSQUITOES

| | PAGE |
|------------------------------|------|
| 1 NETS | 1 |
| 2 MOSQUITO BOOTS | 3 |
| 3 VEILS GLOVES AND HAND FANS | 3 |
| 4 REPELLENTS OR CULICIFUGES | 3 |
| 5 SCREENING | 5 |

Part II

MEASURES DIRECTED AGAINST ADULT MOSQUITOES

| | |
|--|----|
| 1 DESTRUCTION BY HAND | |
| Hand nets Swatters Chloroform Tubes etc | 11 |
| 2 TRAPS | |
| Lefroy's Fletcher's Richmond's Sergeant's Proudlock's Clemesha's Strickland and Chowdhury's James Chatti Traps Traps on Screens Trap holes Illuminated Traps | 12 |
| 3 FUMIGATION | |
| Pyrethrum Powder Sulphur Dioxide Cresol Lysol Creolin Camphor and Carbolic Acid Formalin Chlorine Mercuric Chloride Waste Tobacco Smoke Datura etc | 18 |
| 4 SPRAYS | |
| Gomha's Brug and Van Slooten's Mansell's Lefroy's Army Fly Spray Flit Mosquit Flytox Whiz Flyosan Shelltox Necrosene Fly dead etc | 23 |
| 5 JUNGLE CLEARING | 27 |
| 6 CULTIVATION OF ALLEGED DETREMENT TREES AND PLANTS | |
| Eucalyptus Nim Ch naberry Castor Oil P paja Lavender Clover Pines Casuarina Sun flower etc | 27 |
| 7 USE OF BATS | 28 |
| 8 ZOOTROPIC LAXIS (Animal Prophylax) | 29 |

Part III

MEASURES DIRECTED AGAINST LARVÆ OF MOSQUITOES

1 Description of Methods Used

| | |
|--|----|
| 1 JUNGLE-CLEARING | |
| Where useful where dangerous | 35 |
| 2 DRAINAGE | |
| Open Drainage Subsoil Irrigation Vertical Drainage | 36 |

| | PAGE |
|--|------|
| 3 OILING | |
| How Oil Is Spread ng of Oils Choice of Oil Kerosene Oil Crude or Fuel Oil Ant malar al Oil Pester ne Waste Motor | 44 |
| 4 CHEMICAL LARVICIDES | |
| Paris green—Method of Act on Requirements Choice of Diluent Screen ng Mix ng Methods of Appl cat on Hand bellows Knapsack Sprayers Rotary Blowers Elect ric Blowers Dust ng from Aeroplanes Methods of Determin ng Toxic ty Su tabi ty of Diluent etc Strength to be Used Quant ty to be Used Frequency of Dust ng Cost of Appl cat on Effects on Other Forms of Life Advantages and Disadvantages | 57 |
| Paraformaldehyde (Stoxal) Cresol Cyllin Sanitas Okol Panama Canal Larvic ds Crude Carbolic Acid Creosote Borax Lime Soap Paristan Isotex Empran n Emulsified Necrosene Hycol | 71 |
| 5 SULPHATE OF COPPER | 77 |
| 6 VEGETABLE LARVICIDES | |
| Detritus Cactus Paste etc | 78 |
| 7 LARVICIDAL FISH | 79 |
| 8 DETERRENT AQUATIC PLANTS | |
| <i>Azolla Lemna Wolffia</i> Water hyacinth <i>Utricularia Chara</i> <i>Cladophora</i> | 81 |
| 9 BIOLOGICAL CONTROL OF MOSQUITO BREEDING | 83 |
| <i>B Methods Advocated in Special Circumstances</i> | |
| 1 SWAMPS AND MARSHES | 85 |
| 2 SALT WATER SWAMPS | 86 |
| 3 LAKES AND RESERVOIRS | 87 |
| 4 PONDS AND TANKS | 87 |
| 5 LARGE RIVER BEDS | 88 |
| 6 SMALL STREAMS | 88 |
| 7 HILL-FOOT SEEPAGES | 89 |
| 8 RICE FIELDS | 89 |
| 9 IRRIGATION CHANNELS AND DITCHES | 89 |
| 10 IRRIGATION AND WATER LOGGING | 90 |
| 11 DEPRESSIONS IN THE GROUND | 90 |
| 12 BORROW PITS | 91 |
| 13 SMALL POOLS | 91 |
| 14 HOLES IN ROCKS AND TREES | 91 |

CONTENTS

| | | |
|----|--------------------------|------|
| 15 | WELLS | PAGE |
| 16 | CISTERNS | 91 |
| 17 | FIRE BUCKETS | 92 |
| 18 | RAIN WATER GUTTERS | 93 |
| 9 | WATER USED IN BUILDING | 93 |
| 7 | STAND PIPES AND HYDRANTS | 93 |
| | SLUICE VALVES | 93 |

Part IV

RECENT ADVANCES IN MALARIA CONTROL

| | | | | |
|---|------------|---|---|----|
| 1 | REPELLENTS | 1 | <i>Protection Against Bites of Mosquitoes</i> | 91 |
| 2 | SCREENING | | | 91 |

| | | | | |
|---|----------------|--|---|----|
| 1 | TRAPS | | <i>Measures Directed Against Adult Mosquitoes</i> | 91 |
| 2 | SPEARS | | | 91 |
| 3 | ZOOPOPHYLLAXIS | | | 91 |
| 4 | HOUSING | | | 96 |

| | | |
|----------|---|-----|
| | <i>C Measures Directed Against Larvae of Mosquitoes</i> | 100 |
| DRAINAGE | | 101 |
| OILING | | |

| | | |
|----------------------|--|-----|
| CHEMICAL LARVICIDES | | 102 |
| Paris green | | 104 |
| VEGETABLE LARVICIDES | | |
| Pyrethrum Emulsion | | 107 |
| Cashew nut Shell Oil | | 109 |

| | | |
|--------------------------------------|--|-----|
| NATURALISTIC CONTROL MEASURES | | 111 |
| Chemical Measures | | |
| Pollution of Water | | 113 |
| Changing the Salt | | |
| Physical Measures—Filling by Silting | | |
| Sluicing | | |
| Flooding | | |
| Fluctuating Water Level | | |
| Intermittent Drying | | |
| Agitation of the | | |
| Water Surface | | |
| Stagnating or Ponding | | |
| Muddying | | |
| Shading | | |
| Biological Measures—Larvivorous Fish | | |
| Changing the Fauna and | | |
| Flora | | |

| | |
|---|-----|
| 1) General Conduct of Anti malarial Campaigns | 118 |
| | 119 |

BIBLIOGRAPHY

ADDITIONS TO BIBLIOGRAPHY

| | |
|---|-----|
| APPENDIX I—List of SOURCES OF ANTI MOSQUITO EQUIPMENT | 121 |
| " II—TABLE OF EQUIVALENTS | 166 |
| INDEX | 217 |



LIST OF ILLUSTRATIONS

| | PAGE |
|---|------|
| Fig 1 Method of counting the mesh of cotton netting | 2 |
| " 2 Two types of wire gauze | 7 |
| " 3 Crinoline trap | 14 |
| " 4 Sergeant's trap | 16 |
| " 5 Two systems of laying out drains | 37 |
| " 6 Cross section of key wall | 39 |
| " 7 Cross section of three types of subsoil drains | 40 |
| " 8 Subsoil tile drain longitudinal section | 42 |
| " 9 Vertical drainage | 44 |
| " 10 Oil ball and float | 51 |
| " 11 Screener for dust | 61 |
| " 12 Mixer for use with paris green | 62 |
| " 13 Contour drain for hill side seepage | 85 |
| PLATE I A Oil Sprayer with Double Action Pump (Lalwood No 28) | 52 |
| B High Pressure Oil Sprayer (Bur Oaks Kent No 4) | 52 |
| PLATE II - A Haversack for the application of paris green by hand | 63 |
| B Rotary Blower (Peerless Dust Gun) for paris green | 63 |

PART I. PROTECTION AGAINST BITES OF MOSQUITOES

1. Nets—The use of nets as a protection from the bites of mosquitoes has been practised from very early times. Ross³⁷ notes that they were employed by the Romans and were alluded to by Herodotus, Horace, Juvenal and other early writers. They still remain the most important of all measures of personal protection against the bites of mosquitoes. A mosquito net should always be used in any malarious locality, whatever other precautions may be adopted.

The size of the openings in the netting is important and this is determined not only by the number of holes to the inch but also by the thickness of the cotton of which the netting is made. The mesh is stated in terms of the sum of the number of holes counted along a line of the warp and a line of the bobbin falling within an area of one square inch, the hole at the corner of the square where the two lines meet being counted twice (Fig. 1). Cotton is graded according to weight and in terms of the ratio of the accepted factor of 840 yards to one lb. The mosquito netting supplied to the British Army in India is of .5 26 mesh, woven of 30's cotton. This means that the sum of the holes counted along a line of the warp and a line of the bobbin in one square inch is 25 or 26 and that the cotton of which it is woven goes thirty times 840 yards to one lb. It does *not* mean that there are .5 or .6 holes to the square inch or linear inch of netting. In fact the number of holes to the square inch would be about 150. Netting of this type is however not infrequently misleadingly referred to as being of '25 26 holes to the square inch'.

The material should be white, to allow of easy detection of mosquitoes. The top as well as the sides of the net should be of netting and not of calico, as the latter excludes air. The best pattern is the rectangular net, and the next best a bell net with a circular hoop a yard or two above the bed, to allow of the necessary stretching of the net. Not a single rent or hole in the net should be allowed, and there should be no openings in it for the purpose of entering the net.

If possible a large bed and net should be used, so that the hands, knees and elbows may not be pushed against the net during




FIG. 1

Method of counting the mesh of cotton netting. The mesh is the sum of the number of holes counted along the lines AB and AC, the hole at A being counted twice (after MacArthur²⁰).

sleep, and thus be bitten through it by mosquitoes outside. If these are not available, a loose valance of gauze or a strip of calico should be sewn round the lower part of the net for a height of about 9 inches above the upper surface of the bed. The net should be made especially full at the corners, and it is an advantage to have webbing sewn diagonally across the top of the net to take the strain.

The net should be hung inside the poles and tucked continuously all round under the mattress, and not be allowed to hang down to the floor.

It should be let down before dark in

the evening, and should be stretched as tight as possible in every direction to allow the air to pass freely through it. When going to bed a thorough search should be made for any mosquitoes which may be inside the net, preferably with an electric torch. The net should be put up in the morning by collecting the hanging portion together, twisting it into a coil and throwing it over the top bar of the frame.

When travelling by train in malarious localities, difficulty is often experienced in suspending a net in a railway compartment. Small hooks attached to rubber discs, which will adhere by

suction to any smooth surface, can now be obtained, and will be found most useful for this purpose

2. Mosquito Boots are very useful to protect the ankles in the evening. These may be made either of soft leather, or of canvas. Wellington boots may also be used for this purpose *

3. Veils and Gloves are sometimes used to protect men when on guard at night. Ross³⁷ recommended the use of a hand-fan, an excellent substitute for which is a palm leaf

4. Repellents or Culicifuges.—The common drawback of all these is that after a certain length of time their effect wears off, but they have their value where men have to be out at night for limited periods in malarious situations. They are also useful for application when people are sitting out of doors during those sultry evenings which are so common in the tropics in the malaria season. Dover's preparation (formula d) is particularly suitable. Its smell is not unpleasant, and it may be applied as a pomade to the hair to deter mosquitoes from biting the neck and face⁴⁶⁰. The requirements to be aimed at are as follows —

- (i) It must spread easily
- (ii) It must be of such consistency as to make it adhere strongly to the skin
- (iii) Its base must hinder the too rapid volatilization of the substances which it contains

Bacot and Talbot⁴⁵⁸ concluded that the protection afforded by culicifuges does not result from the dislike of the insect for the preparation but from the latter obscuring the attractiveness of the human odour. They found that culicifuges prepared with grease were more easily applied but not so lasting as those put up with wax. The least possible proportion of the inert base should be used in making up the preparation

* A pillow case drawn up over the feet and legs is a useful precaution for ladies when dining. Joss sticks (e.g., 'Katol') may also be burnt under the table to keep mosquitoes away

A vast number of culicifuges have been recommended the composition of a few of them being given below —

- (a) *Bamber Oil* —Oil of citronella $1\frac{1}{2}$ parts liquid paraffin
1 part coconut oil 2 parts carbolic acid 1 per cent
- (b) Vaseline 100 parts naphthalene 10 parts camphor
1 part
- (c) Castor oil 1 part alcohol 1 part oil of lavender
1 part
- (d) Oil of citronella $\frac{1}{2}$ oz spirits of camphor $\frac{1}{4}$ oz cedar
wood oil $\frac{1}{4}$ oz white petroleum jelly 2 ozs (Dover's
formula⁴⁶⁰)
- (e) Fowler¹⁶ in Palestine found vermujell¹ the best repellent
for field use
- (f) Anti mosquito sprays such as Flit sprayed over
the feet and ankles are useful in repelling mosquitoes
for a considerable period (see under Sprays)

Among other substances recommended may be mentioned sulphur petroleum bergamot oil thuja oil cinamic aldehyde anisic oil a mixture of tar and oil oil of eucalyptus ammonia powdered sandal wood oil of peppermint lemon juice and vinegar oil of turpentine etc

Rudolfs⁴⁶⁶ found that the best effects were gained by mixing the repellent with vaseline talcum powder corn starch etc He found that pyrethrum extract (with vaseline) and clove oil gave protection for from 100 to 120 minutes whilst oil of lime pennyroyal and rose geranium were effective for 90 minutes

Mention should also be made of the carefully conducted experiments of Bunker and Hirschfelder⁴⁵⁷ into the respective merits of a very large number of repellents

Whatever preparation may be used should be applied liberally, especially about the neck and wrists as the sweating induced by tropical climates tends to make the effects transient

Coogle⁴⁵⁸ in the U S A noted that anophelines were less numerous under railway bridges the timbers of which had been

treated with creosote oil than under the neighbouring road bridges not so treated and he recommended the application of creosote oil to the walls and ceilings of houses as a mosquito repellent Fermi²⁸, however who tried this method in Italy reported that it was ineffectual

(See *Bibliography* Nos 456-467)

5 Screening of houses barracks etc

The use of screening especially where electricity is also available for light and fans has made a great difference to the health and comfort of those who are able to occupy well built houses in the tropics It is however essential to screen not only sleeping quarters but also other living rooms and more especially verandahs and to provide if possible overhead fans in these also Otherwise the inevitable result will be that people will sit about outside the building in the hot weather till a late hour at night, probably with a large part of their persons exposed to the bites of mosquitoes

In the case of troops it should be insisted on that the men go inside the screened barracks at sunset and remain there If this is impracticable strict orders should be issued that shorts must be replaced by slacks at sunset and that either jackets must be worn or the sleeves of the shirt pulled down

The value of screening has been generally recognized for many years in America Africa Italy and Malaya in which countries it is extensively employed During the last few years it has been introduced in the barracks of British troops in certain cantonments in India with excellent results The following are the principal points to be observed —

The building to be screened must be well built and in good repair

Door frames should be made of seasoned wood with iron brackets at the corners and should not sag on their hinges A double wire attached diagonally across the lower half of the door and tightened up with a turnbuckle will tend to prevent this

No attempt at edge fitting should be made. The door should be made to fit against a $\frac{3}{4}$ inch batten all round the inside, and sufficient space allowed for the door to swell in wet weathers without scraping at the bottom.

A light strip of wood should be nailed across the door frame at about the height of a man's shoulders. This is to push against when opening the door.

There should be two fastenings for each door, one half-way up the top section and one near the bottom. The lower panel of the door may be strengthened by wire netting,* as a protection against kicks and against dogs attempting to enter or leave the building. Doors should open outwards, so that any mosquitoes resting on them may be driven out when they are opened. They should be placed if possible on the windward side of the house, for mosquitoes tend to congregate on the leeward side. It is of great advantage to have double doors, with a porch or vestibule at least 6 feet in length between them. Strong springs should be fitted to the doors to ensure that they shall remain tightly closed when not in use.

Every aperture in the building should be screened. In the case of chimneys Coogler⁴⁷¹ recommends the suspension of a wire cage containing naphthalene balls inside the chimney about two feet from the top. It may be mentioned, however, that Bunker and Hirschfelder⁴⁵⁷ state that mosquitoes appear to be indifferent to the odour of naphthalene.

No removable screens or shutters should be used.

No screening should be used within one foot of the floor, because it is liable to be damaged when the floor is scrubbed.

No furniture should be placed against the screening.

* If the gauze employed is of copper, the protecting netting should also be of copper and not of plain or galvanized steel in order to avoid galvanic action. The aperture of the netting should be sufficiently small to prevent the ends of canes, umbrellas, etc., from passing through and damaging the gauze.

Orenstein⁴⁵⁴ states that if copper gauze is used it should be fastened with copper tacks, and the rows of tacks covered by thin strips of wood, to avoid electrolyzation

As regards the type of gauze to be used for screening, there are several points to be considered

(1) *Size of Aperture* —As in the case of mosquito netting, the size of the openings in the screen is of the first importance. It is insufficient merely to state the number of openings to the linear inch because the size of aperture will vary with the diameter of the wire used. It may be increased or reduced by 50 per cent by the adoption of a thinner or thicker wire (Fig 2). The diameter of the wire may be expressed by the number of the gauge, but




FIG 2

Two types of wire gauze. Both are of the same mesh, but the aperture of B is less than that of A by one third, owing to the greater diameter of the wire.

unfortunately there are several different gauges in use, each of which denotes a different diameter of wire for the same number. Thus Imperial Standard Wire Gauge (I S W G) No 30, denotes a diameter of 0.0124 inch whilst Steel Wire Gauge (S W G), No 30 denotes a diameter of 0.014 inch. To avoid confusion it is therefore best to specify sizes of wire in decimals of an inch. Another source of confusion is that some authors in referring to size of aperture speak of the area in square inches, whilst others speak of the breadth of the opening in linear

inches The latter notation is the more usual and will be adopted here

The gauze recommended by MacArthur⁴³¹ as being sufficient to exclude *Stegomyia fasciata* from buildings was of 14 mesh and 30 I S W G which gives an aperture of 0.0590 inch. Le Prince and Orenstein³⁴ found that an aperture of 0.0485 inch would permit the passage of this species only under stress of circumstances whilst an aperture of 0.046 inch would exclude them altogether. Farle⁴⁷³ states that in Porto Rico a mesh of 16 with a diameter of wire of 0.009 or 0.010 inch (aperture 0.0595-0.0530 inch) will keep out practically all mosquitoes whilst a mesh of 14 with a diameter of wire of 0.015 inch (aperture 0.056 inch) is sufficient for most places. Although other observers have carried out experiments with various types of gauze they almost invariably neglect to state the one essential point viz the size of aperture. Further research on this subject is urgently required.

The gauze used in the barracks of British troops in India is of 14 mesh and 28.30 S W G giving an aperture of 0.055 to 0.057 inch. The standard adopted by the Punjab Government Public Works Department is of 12 mesh and 25 S W G giving an aperture of 0.063 inch. The size of aperture recommended by Messrs Christie Ltd of Glasgow for the screening of houses is 0.0408 inch (18 mesh and 28 I S W G). This gives a ventilation area of 53.93 per cent as compared with 68.23 per cent given by a gauze of 14 mesh and 30 I S W G.

(ii) *Strength of Wire*—It is obvious that other things being equal the greater the diameter of the wire used the stronger will be the gauze. It is equally obvious however that the use of a thicker wire will mean the exclusion of more air and light. Our aim therefore should be to obtain the thinnest wire which will give the required strength. *The gauge usually recommended is 28-30 I S W G denoting a diameter of 0.0148 to 0.0124 inch or 28-30 S W G denoting a diameter of 0.016 to 0.014 inch.* In order to obtain the greatest amount of air and daylight a

heavier gauze may be used in places subject to the greatest amount of wear, such as doors and the lower half of verandah screens, etc., a lighter type being employed for less exposed situations

(iii) *Durability of Screen*—The material of which the gauze is made is of great importance. Painted* steel or galvanized iron wire are the cheapest, but are not suitable except in a very dry climate. Darling³⁷⁴ found that in a warm moist atmosphere galvanized iron and brass screening corroded rapidly whilst copper and phosphorized bronze both resisted deterioration admirably. When tested in a salt laden atmosphere the copper screening also became badly corroded, the bronze resisting to a much greater degree. He therefore concluded that phosphorized bronze was best for places near the sea, but that for other places in the tropics the less expensive copper screening was just as effective, provided the copper content was high (about 90 per cent), and the amount of iron as low as possible (not more than $\frac{1}{2}$ per cent).

MacArthur⁴⁸¹ recommends the use of 'Monel metal' as being the best for use in a moist climate. This consists of nickel 57 per cent, copper 28 per cent, other metal 5 per cent. It is a natural alloy of metals, and contains no tin, zinc or antimony. A disadvantage of this is its high cost. Earle⁴⁷³ states that in Porto Rico the only two kinds of wire which have been found durable are of Monel metal, diameter 0.009–0.010 inch, and of copper or bronze, diameter 0.015 inch.

Wurtz⁴⁹² recommended the use of ordinary cotton netting, treated with a solution of commercial silicate of potassium in its own volume of water. This is painted on several times with a brush, without drying between the coats, and is said to make the cotton much stronger and more resistant to water or fire.

The gauze now being tried in the British troops' barracks in India is composed of copper 90 per cent, and tin 10 per cent.

* It should be noted that painting reduces the size of the aperture.

The essential requirement for screening is that the work should be thoroughly carried out. Frequent inspection is necessary for the detection of rents in the wire and defects in the wooden framework the latter being especially likely to develop where there is extreme variation in humidity between the dry and wet seasons.

(See *Bibliography* Nos 468-493)

PART II.

MEASURES DIRECTED AGAINST ADULT MOSQUITOES.

1. Destruction by Hand.—This method has received increasing attention in recent years, and in view of the fact that recently fed and therefore possibly infected mosquitoes may be killed in this way it has an important bearing in the prevention of malaria. Children may be taught to use the methods enumerated below and Ross³⁷ has suggested that the offer of one anna for 50 anophelines would probably result in a very extensive catch in any village.

(a) *Hand nets* may be made cheaply by tying a piece of flexible cane or wire in the form of a loop to which a bag of white netting about 18 inches deep is attached. In lofty rooms a long handled net may be used.

(b) *Swatters* consisting of a piece of wire gauze about 6 inches square fixed to a two foot wooden handle may be used and are especially effective on wire screens.

(c) A *chloroform tube* 7 inches long by 1 inch in diameter may be used. A layer of rubber bands or pieces of old tyres one inch thick is placed in the bottom of the tube held in place by a plug of absorbent cotton covered with a disc of blotting paper or cork. A few c.c. of chloroform are poured in and are absorbed by the rubber. This will last for two or three weeks. A test tube with a plug of cotton wool soaked in benzene or petrol may also be used or a test tube wetted inside with kerosene. To catch the mosquito remove the cork and place the mouth of the tube quickly over the insect as it rests on some object. An electric torch is a great assistance.

(d) A tin with some petrol or kerosene in the bottom, fixed to the end of a long stick, is useful in catching mosquitoes on ceilings

(e) Curtains etc., may be shaken and the mosquitoes flying out caught by hand if the latter is first dipped in soapy water

The above methods may be used to supplement screening

2. Traps.—The use of traps for catching mosquitoes was suggested by Nuttall and Shipley⁵¹⁷ in 1902, as the result of their experiments to ascertain the colour most attractive to mosquitoes*. Various different forms of trap have been devised, several of which are described below

Traps are used for three chief purposes, (i) for reducing the number of mosquitoes (ii) to find out the relative prevalence of the different species of mosquitoes and (iii) in order to gauge the effect of control measures. In the last two cases the traps are placed in various quarters of the area under observation and counts are made of the mosquitoes captured in each trap daily. This is a valuable method for estimating the efficiency of anti-larval measures and an increase in the catch in any particular locality will often lead to the detection of some breeding place which would otherwise be overlooked.

Traps may be used either inside or outside buildings. James' trap²⁷³ was designed for use in the shaded corner of a garden, and Zetek⁵²¹ recorded that in camps where traps were placed on the leeward side of buildings a large number of anophelines were caught, resulting in a reduction in the incidence of malaria.

When used inside houses, barracks etc., the success of traps largely depends on the type of building in which they are employed. In well built barracks with whitewashed walls the proportion of mosquitoes caught will be considerable because the traps form a more attractive resting place than the bare white

* These observers found that dark blue was the most attractive colour, and it is suggested that this would probably be the most effective colour to use for the inside of the traps here described.

walls. But in ill lighted, 'kuchcha' built buildings where there are many dark corners, crannies in the walls, etc. providing suitable resting places for mosquitoes only a small proportion of these insects will be found in the traps. The placing of baits in traps (excepting human baits see p 18), has not been attended with success* and in those described below the object aimed at has been merely to provide an attractive resting place for mosquitoes.

(a) *Lefroy's Trap*⁵¹³ — This is a wooden box 12 inches long by 12 inches wide and 9 inches deep lined with dark green baize and having a hinged door. The floor of the box is covered with tin. The trap is set in a dark corner at night and in the morning the door is closed and the insects killed by dropping a teaspoonful of benzene or chloroform through a small hole in the top of the box fitted with a movable plate.

(b) *Fletcher's Trap*⁵¹¹ — This consists of a wooden skeleton of a box with a hinged lid covered on all sides with black mosquito netting. This is contained inside an open topped wooden box, painted black inside. The whole is placed in a suitable position at night with the hinged top of the inner box open. In the morning the lid is closed with a metal hook fastener and the inner box lifted out by handles provided for the purpose. The mosquitoes may be killed by placing this on the bare ground in full sunlight.

(c) *Richmond and Mendis Traps*²³⁸ — These observers have made a number of experiments with various traps of their own device in barracks. The most successful of these were —

- (i) *The Crinoline Trap* — A 'skirt' of navy blue or black cloth with an upper opening 4 inches in diameter (which can be closed with a purse string) is suspended from the roof about 6 feet above the floor level. A bamboo hoop 3 feet in diameter is attached to the inside of the skirt 2½ feet from the top, and below

* Meta* has suggested the employment of small p g stives as traps for outside use (see 'Zooprophylaxis').

the hoop the skirt is continued in the form of a fringe hanging vertically downward for a distance of 3 feet. The lower opening can be closed by a purse string, which runs through loops round the *outside* of the skirt (Fig 3). At night the upper opening is closed, whilst the lower is left open. In the morning the lower opening is closed, and an inverted glass jam jar is inserted into the upper one. The trap itself is then well shaken with the result that the imprisoned insects fly upwards towards the light into the jar, and can be removed alive.




FIG. 3

Crinoline trap (after Richmond and Mendis²²⁹)

If living mosquitoes are not required, the insects may be killed by simply spraying them with 'Flit' or other spray, without previously closing the lower opening.

- (ii) *Box Traps*—These are made of wood, 3 feet by 2 feet by 2 feet, lined inside with black cloth, open at the

end with no door but fitted with a detachable black muslin lining with a purse string at the open end. This lining is provided with loops which slip over hooks inside the box. The purse string is closed in the morning when the sleeve may be detached and the insects removed alive if desired.

- (iii) *Wire Traps*—These are of the same dimensions as the box traps but consist of 4 strong wire frames which can be joined together with string to form the framework of the trap. A black cloth covering is slipped over this and a detachable sleeve (as described above) attached inside. The advantage of this type over the box trap is its easy portability which makes it particularly suitable for field service conditions.

Of the three forms of trap described above Richmond and Mendis found the box and wire traps the most effective. One trap to each 5 000 cubic feet of space in a well built British barrack room with a capacity of 20 000 cubic feet captured 60 to 65 per cent of the anophelines present. The traps were placed 2 or 3 feet above the level of the floor.

- (d) *Sergeant's Trap*—This is a portable wooden trap described by Legroux²² and stated by him to have been designed by Dr Etienne Sergeant. One end of the trap is smaller than the other and at this end there is a gauze door overhung by a continuation of the roof of the trap (Fig 4). The door is left open at night and is closed in the morning. At the larger end of the trap there is a gauze-covered window with a wooden shutter. By opening the shutter the catch can be viewed and counted.

- (e) *Proudlock's Traps*. These are ingeniously devised patent traps made in two principal types.

- (i) *The Household Type* for use in hungilows etc consists of a small wooden box painted black inside with a vertical entrance in front fitted with a

shutter, and a sliding panel at the back. When the latter is opened a glass window is disclosed through which the catch may be viewed. When this is exposed to sunlight the insects are quickly killed and may be emptied out through a small hole at the side of the trap fitted with a shutter.




FIG 4

Sergeant's trap (after Legroux²¹⁰) The gauze door in front is opened to allow mosquitoes to enter. The shutter over the circular gauze covered window at the back is closed.

- (11) *The Research Type* is a more elaborate trap consisting of an elongated box with a funnel at one end through which the mosquitoes enter. At the rear end at the side of the box there is a small opening with a shutter over which can be fitted a slide rather like the plate holder of a camera but with a glass face. In the morning a plunger is inserted at the front end of the box and the shutter at the rear end opened together with a corresponding shutter in the slide. The plunger is then pushed home and the mosquitoes are driven into the slide box which can then be removed.

(f) *Clemesha's Trap*—This is an ordinary tea plucking basket, plastered inside and out with cowdung and covered with a lid, also plastered in which there is a central opening 6 inches in diameter. The trap is hung up a little above the floor in a

cowshed or other building with some grass or hay in the bottom. The catch is collected by first closing the opening in the lid and then placing the trap under a mosquito net. The operator enters the net, opens the lid and shakes the trap, the mosquitoes being caught in test tubes on the inside of the net.

(g) *Strickland and Choudhury's Trap*⁵¹²—This is made from a tea box, 18 inches by 18 inches by 18 inches, blackened inside, and with a circular hole in the side which can be closed by means of a swivelled door. Strickland and Chowdhury recorded better results with this trap than with Clemesha's trap.

(h) *James' Trap*⁵¹³—A rectangular wooden framework, 5 feet by 3 feet by 3 feet, is covered with mosquito netting, one end of it being a door on hinges. The trap is placed on the ground in a shaded corner of a garden and covered with sacking and a thick tarpaulin, so that the interior is dark and cool. Two or three plants are placed inside each trap and several near the door, which is left partly open at night. At about 8 or 9 a.m., the vegetation round is disturbed as much as possible, and straw and paper torches are burnt in all the surrounding outhouses and buildings to drive out the mosquitoes, most of which take shelter in the trap. About half an hour later the door of the trap is closed, the sacking and tarpaulin removed and the trap placed in strong sunlight for one or two hours when all the mosquitoes are killed.

(i) *Chatt's Traps*—These are the ordinary earthenware water vessels commonly used in tropical countries. They are half filled with water, and placed in corners of buildings. Mosquitoes may be collected from them by means of a cylindrical frame covered with gauze, which is continued upwards in the form of a sleeve. They are useful for catching culicines but are ineffective for anophelines.

(j) *Traps on Screens*—These are essentially labyrinths of wire gauze on wooden frameworks, which are fixed on to the inside or outside of screens of buildings. Most mosquitoes are caught while trying to get in. The traps act automatically, and need only be emptied at long intervals. They have been used with great success in Panama.⁵¹⁴

McHardy⁵¹⁵ in Dar es Salaam has devised a 'trap house' with these traps fixed on the wire gauze which screens it. The trap house is made in sections, so that it can be easily taken down and erected wherever desired. African natives sleep inside it acting as a 'bait'.

(L) *Trap holes*—Blin⁵¹⁰ described experiments in which he dug oblique holes in the ground near houses infested with mosquitoes. The insects which came to rest in them to seek darkness and protection from the heat were destroyed in the daytime with a torch made with a wisp of straw or a stick dipped in kerosene.

(I) *Illuminated Traps* have been used in the U S A with considerable success (Owen⁵¹⁸, Headlee⁵¹²). Certain species of mosquitoes are attracted by a moderate light, though if the light be too strong it repels them. The trap used by Headlee is fashioned from a sugar tin, 12 inches high and 10 to 12 inches in diameter, with the bottom removed and replaced by a cone of wire gauze, which projects into its interior. There is an opening not more than $\frac{1}{4}$ inch in diameter at the summit of the cone. Light is supplied by a small electric bulb connected with a battery.

(See *Bibliography*, Nos 510-521.)

3. Fumigation.—Whichever method is used all chinks in the doors and windows should first be closed by hanging blankets over them. It is best to leave one window uncovered and to darken all the others. A white sheet is placed on the floor in front of the uncovered window. The reason for this is that in most cases mosquitoes are not killed by fumigation, but are only stupefied. They will for the most part fall on the sheet, and can then easily be seen, swept up and burnt.

(a) *Pyrethrum Powder* (*Pyrethri flores*)*—This is made from the dried flower heads of the chrysanthemum (*Pyrethrum cinerariaefolium*). It forms the main ingredient of

* This must not be confused with *Pyrethra radix* (pellitory root), which has also been recommended as a fumigant.

many insect powders, e.g., 'Keating's Powder,' and owes its insecticidal activity to a mixture of acids and esters. It also contains oleoresins to which it probably owes its value as a fumigant and since these are volatile it is essential that the powder shall be fresh. It is also important to ensure that it is not adulterated with the powdered stems of the plant, which is a common practice.

The powder may be used dry and puffed or blown into the air of a room. For fumigation it is heaped up in an iron pot into a little pyramid, which is lighted at the top and burns slowly, giving off a dense and pungent smoke. A layer of sawdust 2 to 3 inches deep placed in the bottom of the pot in the form of a crater permits complete combustion thus saving 10 to 15 per cent of the fumigant which would otherwise be wasted. The powder may also be moistened and made up into small cones which burn readily when dry. One pound at least is necessary for 1,000 cubic feet of space (Boyce³ recommends 3 lbs). The room should be kept closed for a period of 3 hours.

Other methods are to heat the powder on a metal plate over a kerosene lamp or to puff it from an insufflator into a burning gas jet. The fumes are not noxious to human beings.

(b) *Sulphur Dioxide*—This may be applied in various ways.

- (1) For operations on a large scale the use of a Clayton machine has proved most effective for destroying mosquitoes in yellow fever epidemics. Various types of Clayton machine are on the market. At Accra a machine mounted on a bogie carriage is always kept ready for action. For treating ships, the machines are installed in special boats. There is also a light type of Clayton machine which can be carried by two men, it is air-cooled, and is capable of saturating 6 000 cubic feet of space per hour with a 15 per cent strength of gas (Parsons and Brook²⁶).

McHardy⁵¹⁵ in Dar es Salaam has devised a 'trap house' with these traps fixed on the wire gauze which screens it. The trap house is made in sections so that it can be easily taken down and erected wherever desired. African natives sleep inside it, acting as a 'bait'.

(A) *Trap holes*—Blin⁵¹⁰ described experiments in which he dug oblique holes in the ground near houses infested with mosquitoes. The insects which came to rest in them to seek darkness and protection from the heat were destroyed in the daytime with a torch made with a wisp of straw or a stick dipped in kerosene.

(I) *Illuminated Traps* have been used in the U S A with considerable success (Owen⁵¹⁸, Headlee⁵¹²). Certain species of mosquitoes are attracted by a moderate light, though if the light be too strong it repels them. The trap used by Headlee is fashioned from a sugar tin 12 inches high and 10 to 12 inches in diameter, with the bottom removed and replaced by a cone of wire gauze, which projects into its interior. There is an opening not more than $\frac{1}{4}$ inch in diameter at the summit of the cone. Light is supplied by a small electric bulb connected with a battery.

(See *Bibliography*, Nos 510-521.)

3. Fumigation—Whichever method is used all chinks in the doors and windows should first be closed by hanging blankets over them. It is best to leave one window uncovered and to darken all the others. A white sheet is placed on the floor in front of the uncovered window. The reason for this is that in most cases mosquitoes are not killed by fumigation, but are only stupefied. They will for the most part fall on the sheet, and can then easily be seen, swept up and burnt.

(a) *Pyrethrum Powder* (*Pyrethri flores*)*—This is made from the dried flower heads of the chrysanthemum (*Pyrethrum cinerariaefolium*). It forms the main ingredient of

* This must not be confused with *Pyrethri radix* (pellitory root) which has also been recommended as a fumigant.

many insect powders e.g. 'Heatings Powder' and owes its insecticidal activity to a mixture of acids and esters. It also contains oleoresins to which it probably owes its value as a fumigant and since these are volatile it is essential that the powder shall be fresh. It is also important to ensure that it is not adulterated with the powdered stems of the plant which is a common practice.

The powder may be used dry and puffed or blown into the air of a room. For fumigation it is heaped up in an iron pot into a little pyramid which is lighted at the top and burns slowly giving off a dense and pungent smoke. A layer of sawdust 2 to 3 inches deep placed in the bottom of the pot in the form of a crater permits complete combustion thus saving 10 to 15 per cent of the fumigant which would otherwise be wasted. The powder may also be moistened and made up into small cones which burn readily when dry. One pound at least is necessary for 1 000 cubic feet of space (Boyce³ recommends 3 lbs). The room should be kept closed for a period of 3 hours.

Other methods are to heat the powder on a metal plate over a kerosene lamp or to puff it from an insufflator into a burning gas jet. The fumes are not noxious to human beings.

(b) *Sulphur Dioxide*—This may be applied in various ways.

- (1) For operations on a large scale the use of a Clayton machine has proved most effective for destroying mosquitoes in yellow fever epidemics. Various types of Clayton machine are on the market. At Accra a machine mounted on a bogie carriage is always kept ready for action. For treating ships the machines are installed in special boats. There is also a light type of Clayton machine which can be carried by two men; it is air cooled and is capable of saturating 6 000 cubic feet of space per hour with a 15 per cent strength of gas (Parsons and Brool²³⁰).

McHardy⁵¹⁵ in Dar es Salaam has devised a 'trap house' with these traps fixed on the wire gauze which screens it. The trap house is made in sections so that it can be easily taken down and erected wherever desired. African natives sleep inside it, acting as a bait.

(k) *Trap holes*—Blin⁵¹⁶ described experiments in which he dug oblique holes in the ground near houses infested with mosquitoes. The insects which came to rest in them to seek darkness and protection from the heat were destroyed in the daytime with a torch made with a wisp of straw or a stick dipped in kerosene.

(l) *Illuminated Traps* have been used in the U. S. A. with considerable success (Owen⁵¹⁸, Headlee⁵¹⁷). Certain species of mosquitoes are attracted by a moderate light, though if the light be too strong it repels them. The trap used by Headlee is fashioned from a sugar tin, 12 inches high and 10 to 12 inches in diameter, with the bottom removed and replaced by a cone of wire gauze, which projects into its interior. There is an opening not more than $\frac{1}{4}$ inch in diameter at the summit of the cone. Light is supplied by a small electric bulb connected with a battery.

(See *Bibliography*, Nos. 510-521.)

3. Fumigation—Whichever method is used all chinks in the doors and windows should first be closed by hanging blankets over them. It is best to leave one window uncovered and to darken all the others. A white sheet is placed on the floor in front of the uncovered window. The reason for this is that in most cases mosquitoes are not killed by fumigation, but are only stupefied. They will for the most part fall on the sheet, and can then easily be seen, swept up and burnt.

(a) *Pyrethrum Powder* (*Pyrethri flores*)*—This is made from the dried flower heads of the chrysanthemum (*Pyrethrum cinerariaefolium*). It forms the main ingredient of

* This must not be confused with *Pyrethri radix* (pellitory root) which has also been recommended as a fumigant.

(c) *Cresylic Acid Derivatives*—

- (i) *Cresol Vapour*—This is probably the cheapest fumigant. Four to five ounces of Liq. cresol sap are required per 1 000 cubic feet. It may be slowly vaporized over a kerosene stove of the 'Beatrice' type (in which the flame is guarded by a metal chimney with a mica window) the cresol being placed in some enamelled receptacle which should completely cover the grating on which it rests. The receptacle should be as large and deep as possible so long as its position is secure. Constant observation is necessary and a bucket of water should be at hand in case the cresol should catch fire. The stove can be put out usually in about half an hour's time (the time varies according to the size of the room and the amount of cresol used) after which the room should be closed for another $1\frac{1}{2}$ to 2 hours. It may also be evaporated in kerosene tins over charcoal braziers. The braziers should be glowing when put into a room otherwise the fumes given off are liable to blacken the walls and furniture. Richmond and Mendis²³⁸ found that the cost of fumigating with cresol vapour including the price of the charcoal used was 1 65 annas per 1 000 cubic feet. Another method of vaporizing cresol is to ignite dung cakes and pour the cresol over them when half burnt.

Although the cost of cresol fumigation is so low the amount of preparation and precautions required for its use are much greater than in the case of sulphur. Opinions differ as to the relative efficiency of the two methods. Richmond and Mendis consider sulphur fumigation to be the more satisfactory though the cost (about $3\frac{1}{2}$ annas per 1 000 cubic feet of space) is more than twice as great as with cresol. Hanafin * on the other hand considers cresol to be the more effective and has obtained excellent results

* Private communication

- (11) It may be applied in the form of 'Sulphume,' a proprietary preparation of liquid sulphur dioxide put up in metal cylinders holding about 20 ozs. This was however not found very effective in a single experiment conducted by Parsons and Brook²³⁶
- (111) Sulphur may be placed in an iron pot, and this placed on bricks in a tub or other receptacle containing an inch of water in the bottom. It is readily ignited by sprinkling methylated spirit or kerosene over it and lighting it. It is usually stated that 2 lbs of sulphur should be burnt for 1 000 cubic feet of space with exposure to the fumes for 2 to 3 hours (but see below)
- (iv) *Giles Sulphur Cones*—These consist of one part each of nitre and charcoal to eight of sulphur made up into pastilles weighing 4 ozs each by means of a little gum water and dried in the sun. Giles⁹ recommended that one pastille should be burnt for every 1 000 cubic feet of space but this would only be equivalent to 3.2 ozs of sulphur. Richmond and Mendis²³⁸ found that 4 pastilles (12.8 ozs of sulphur) were effective in barracks for every 1 000 cubic feet of space whilst for fumigating tents 6 pastilles (19.2 ozs of sulphur) were required per 1 000 cubic feet. Even this amount, however, is much less than the quantity of sulphur specified by other authors as necessary.

Richmond and Mendis²³⁸ found the use of Giles' cones a very convenient method for sulphur fumigation. They are placed on tin plates on the floor, and when lighted burn quickly, producing a maximum concentration of sulphur fumes in a very short time.

A great disadvantage of sulphur is that it tarnishes metal work and injures pianos, sewing machines, watches, telephones, etc. It is also harmful to food stuffs.

destructor of mosquitoes but its use is attended with too much risk to make it suitable for general use²³⁶

(g) *Mercuric Chloride* has been used by Guiteras²³³ in America sublimated in a porcelain dish over an alcohol lamp. He used 25 gm. to 1 000 cubic feet of space allowing it to act for 2 to 3 hours. He claimed that its poisonous qualities do not constitute a real danger that it is very simple and effective and requires very little apparatus.

(h) *Waste Tobacco* has been used with effect in Russia²³⁷ mixed with chopped hay or straw and burnt on iron sheets. It is left to smoulder for 16 hours and 30–40 gm. are required per cubic metre of space.

(i) *Smoke Production* —The principle of this measure is to fill the building with smoke quickly to drive mosquitoes out of holes and corners and to attract them to one spot where they can be destroyed. The doorway is covered with a white sheet made as taut as possible. A pile of dry straw is lighted and then covered with damp straw or manure. When the smoke fills the building the insects fly to the sheet where they can be killed.²³⁹

(j) *Datura stramonium* in powder may be used (8 ozs. to 1 000 cubic feet) made up with nitre or saltpetre in the proportion of 1 to 3 of datura. The mixture may be burnt on a tin pan or shovel. Smith²⁴¹ states that the fumes are not poisonous to human beings, are not injurious to fabrics or metals and can be used with entire safety.

Many other substances have been tried as fumigants such as chloral vapour, dried orange peel, *Artemisia (santonica)* pastilles of pellitory root (*Pyrethri radix*, the dried root of *Anacyclus pyrethrum*), dinitrocresol, etc.

The relative merits of fumigation and spraying are discussed on p. 25.

(See *Bibliography* Nos. 218–241.)

4 Sprays —Experiments with various sprays as destructors of mosquitoes were carried out by Giemsa and Muhlens²⁴⁰ in 1911 following the appearance of an article by Stenel²³⁸ who called

with it in Lahore No precaution was taken to seal up the rooms treated beyond closing the doors and windows

(ii) *Cresol and Lysol* —A mixture of equal parts of these was used by Parsons and Brook²³⁵ with satisfactory results

(iii) *Creolin and I-zal* —Alexander²¹⁸ in Accra made various experiments with these preparations Creolin was first used then a mixture of creolin and izal and finally izal alone the latter proving very satisfactory It was found necessary to use enamelled receptacles otherwise soot was deposited The technique is similar to that for cresol

(d) *Camphor and Carbolic Acid* (*Mimm's Culicide* ') — Equal parts of camphor and crystallized carbolic acid are fused together into a liquid by gentle heat Three to four ounces of the mixture must be vaporized for each 1 000 cubic feet of space This is done by placing the liquid in a wide shallow pan over a spirit or petroleum lamp when white fumes are given off To prevent the mixture from burning the fumes should not be allowed to come into close contact with the flame of the lamp The room must be kept closed for two hours The vapour is not dangerous to human life except when very dense but it produces a headache if too freely breathed²⁴

(e) *Formic Acid* —If commercial formalin (40 per cent formic aldehyde) and potassium permanganate are brought together, formic acid fumes are liberated which are toxic to mosquitoes For 1 000 cubic feet of space 10 ozs of potassium permanganate crystals are placed in a bucket containing 1 to 2 pints of water and 1 pint of formalin is poured into it An advantage of this method is that no heating apparatus is required so that there is no danger of fire, also there is no ill effect on furniture²³⁶

(f) *Chlorine Gas* —This may be applied from cylinders containing liquid chlorine or may be generated by pouring crude sulphuric acid on to chloride of lime It is a powerful rapid and economical

and petroleum ether. They did not confirm the results of Brug and Van Slooten, as they found that in the case of all their experimental preparations a large proportion of the insects recovered after some hours in the fresh air, whereas with the proprietary preparations this was not the case.

The cost of these preparations however, is so high that it would appear best to use a locally prepared mixture on the lines of that recommended by Brug and Van Slooten: the mosquitoes stupefied by the spray being swept up and burnt immediately afterwards. One such preparation has been used with success by Mansell⁵⁰⁷ in Peshawar (formula No. 6). Sprays of this nature should not be used in the vicinity of naked lights. For small rooms they may be applied by means of the hand sprayers or atomisers which are now obtainable almost everywhere at a trifling cost. For larger buildings such as barracks high pressure sprayers, such as the 'Four Oaks' Sprayer, have been used with success, though in the very lofty barracks usually built in the tropics it is sometimes found difficult to secure effective results.

As regards the relative merits of fumigation and spraying, the latter would appear to offer many advantages. James,* working with artificially infected *A. maculipennis* arrived at the conclusion that infected anophelines usually pass their whole life in the house where the infection occurred. If this were of universal application, fumigation or spraying even at long intervals would appear to be an effective means of controlling malaria. Subsequent investigations by many workers in different parts of the world have however shown that there is normally a very considerable to and fro movement of anophelines in and out of buildings, and that even shortly after a blood meal anophelines may change their abode (Klugler and Lieberman⁵⁴³, Richmond and Mendis³⁸, and others). Either method therefore to be successful must be repeated at frequent intervals (every 3 or 4 days) and in the case of fumigation the necessity for evacuating rooms or barracks

* *Trans. Roy. Soc. Trop. Med. and Hyg.*, XX, p. 143 (1926).

attention to the neglect of this method of malaria prevention. Mixtures containing cresol and phenol were found too irritating to the human respiratory tract, and the same objection applied to solutions of quassia. Finally Giemsa evolved a mixture containing pyrethrum tincture, soft soap and glycerin, which constituted an effective spray free from the above objections (see formula No 1 below). The pyrethrum tincture was however expensive, and Giemsa⁵⁰³ conducted further experiments to find a cheaper spray. He found that soap solutions alone gave good results and that the toxic properties of a number of other substances besides pyrethrum were enhanced by the addition of comparatively small quantities of soap. He recommended the use of soap solutions containing formaldehyde as being both cheap and effective.

In recent years a large number of proprietary sprays have been put on the market, most of them having a basis of kerosene oil. The following may be mentioned — Flit, Flytox, Mosquil, Shell tox, Whiz, Flyosan, Fly ded, Necrosene Rids, Aho Spree, and Abis. Richmond and Mendis³³, experimenting with Flit, Mosquil, kerosene and Army Fly Spray, found Flit the most effective but considered that the somewhat greater success attained with this preparation, compared with Mosquil and kerosene, was scarcely worth the extra cost. Mosquil was more effective than kerosene, but four times as costly, whilst Army Fly Spray (formula No 3) was not satisfactory.

Brug and Van Slooten⁴⁰⁷ found that a mixture of kerosene with 2 per cent of carbon tetrachloride was effective, and was superior to Flit in that it did not produce stains on white fabrics. This gave as good results as a mixture which they termed 'Rids locally prepared' (formula No 5).

Swellengrebel and his co workers³⁰³ carried out experiments with various proprietary sprays, and with a large number of experimental preparations consisting of kerosene oil with the addition of small proportions of carbon tetrachloride, methyl salicylate, turpentine, creosote, oil of cajaput, aniline oil, petrol

(5) *Rods locally prepared* (Brug and Van Slooten)—

| | |
|----------------------|------|
| Kerosene oil | 89 6 |
| Carbon tetrachloride | 7 |
| Methyl alcohol | 3 4 |

(6) *Mosses Spray*—

| | |
|---|---------------------------|
| Carbon tetrachloride (medicinally pure) | 1 |
| Synthetic oil of wintergreen | 2 |
| Kerosene oil and quality | 97 |
| Naphthalene | $\frac{1}{2}$ lb per gall |

(See *Bibliography* Nos 494-509)

5 Cutting down of Jungle and Other Vegetation—

Many species of anophelines spend the daytime sheltering amongst long grass or in bushes and other coarse vegetation and the clearing away of these in the vicinity of dwellings is therefore of value in the campaign against adult mosquitoes (see also p 35)

6 Cultivation of Alleged Deterrent Trees and Plants—

Eucalyptus trees mimosa trees chinaberry trees castor oil plants papaya lavender clover and various aquatic plants have been said to have a deterrent effect on mosquitoes but experiments have proved that none of them have any direct effect against adults

de Herelle¹³⁸ seeking to explain the relative immunity from malaria of certain tracts of country (Tuscany Argentina Holland) in spite of the presence of anopheline mosquitoes noted that clover was grown in such countries and suggested that coumarin ingested by the female insect from the clover might have an effect on the malaria parasite harboured by it comparable with that which quinine has on the parasite in man

Willcocks¹⁵⁴ quoted de Herelle's remarks and suggested that the relative immunity of cultivated Egypt from malaria might likewise be due to the presence of clover He advocated the introduction of clover and other leguminous plants and trees into Palestine Cyprus Irak Northern India the Sudan and Uganda His paper was widely commented on by the lay press

for periods of several hours makes the prosecution of an effective campaign difficult or impossible. Moreover with sulphur dioxide the tarnishing of metals and injury to watches etc. constitute a great drawback which renders the procedure most unpopular among troops. With sprays the whole operation can be performed in a very short time without the necessity of removing any of the contents of the building and the rooms can be re occupied within half an hour of spraying. Another advantage of spraying is that it is also effective against sandflies on which fumigation appears to have but little effect²³⁸. Spraying may be used with advantage to supplement screening.

The formulæ of various sprays are given below —

| | <i>Parts</i> |
|---|--------------|
| (1) <i>G emsa s Spray—</i> | |
| Pyrethrum tincture (90 parts pyrethrum extracted in | |
| 100 parts methylated spirit) | 580 |
| Potash soap | 180 |
| Glycerin | 240 |

The above mixture to be diluted with 90 volumes of water before use. 100 grammes of the fluid suffice for 50 cubic metres of space.

| | |
|----------------------------|------------------------|
| (2) <i>Lefroy s Spray—</i> | lbs |
| Pyrethrum | o |
| | gall |
| Alcohol | 1 |
| Safrol | 1 |
| Soap | q s to |
| | form an emulsion |
| | (usually about 10 ozs) |

In hot climates add $\frac{1}{2}$ to 2 per cent castor oil. Dilute one in 30 before use.

| | <i>Parts</i> |
|------------------------------|--------------|
| (3) <i>Ar n s Fly Spray—</i> | |
| Pulv s pyrethrum | 117 |
| Spirit rectificatus | 91° (or q s) |
| Camphor oil | 780 |
| Sapo durus pulv s | o 37 |

| | |
|--|-----|
| (4) <i>B ug and Van Slooten s Spray—</i> | |
| Kerosene oil | 100 |
| Carbon tetrachloride | o |

could not be induced to live in the roosts. Nelson⁵¹ has pointed out that bats are of little value against mosquitoes, their chief food consisting of other insects. It is generally recognized that the employment of these creatures is not likely to prove of practical utility in mosquito control.

(See *Bibliography*, Nos 48-54.)

8. Zooprophyllaxis.—Animal prophylaxis against malaria was suggested by Roubaud⁵⁵⁴ in 1919, although Rizzi⁵⁵³ claims that this method of protection was first recognized and practised in Italy. Roubaud pointed out that *A. maculipennis* only feeds on man in the absence of other sources of mammalian blood and suggested that man could have complete protection from the attacks of this species by keeping cattle in the proximity of human dwellings. He stated that in regions where cattle were plentiful a race of these mosquitoes has evolved which have adapted themselves to feed exclusively on animals. Other authors have pointed out that whilst the presence of cattle may attract mosquitoes to themselves, the total population of anophelines in a locality is largely increased where cattle are plentiful. There are also other factors to be considered, one being that the presence of large numbers of cattle goes hand in hand with increased agricultural prosperity, better drainage, clearing of jungle and a betterment of the social and economic condition of the population all of which may tend to diminish the incidence of malaria. It may further be pointed out that the presence of anophelines in a cowshed does not necessarily mean that these have fed on the cattle occupying it. Numerous experiments by means of the precipitin test have shown how frequent it is to find human blood in mosquitoes caught in cattle byres, stables, etc. and, conversely to find the blood of cattle and other domestic animals in those captured in human habitations.

Symes⁵⁵⁵ records that in Taveta, Kenya Colony, the women and children have for years lived in huts together with their cattle. The parasite rate is 50-60 per cent. and the spleen rate 85-100 per cent, whilst it is usual to find the majority of the malaria-carrying

Mayne¹⁴⁸ carried out laboratory tests with coumarin and various species of Indian anophelines, and concluded (1) that coumarin did not exert any deleterious effect on the lives of the mosquitoes feeding on it, and (2) that coumarin, taken with the liquid extracted from a clover blossom, would have no deleterious action on oocysts and sporozoites undergoing development in the infected mosquito

There is no definite proof that clover has any effect on the incidence of malaria, and the only recorded experiments which have been carried out as to the effect of coumarin on mosquitoes or malaria parasites indicate that it has no effect on either under natural conditions

The planting of certain trees and plants, however, has a good effect by drying up water and reducing the level of the subsoil water. Amongst others which have been recommended in India for this purpose are —Pines, Casuarina, Nim, Eucalyptus, Sun flower, Canna. Castor oil plant, Chrysanthemum

The planting of belts of high trees* as a screen against the diffusion of mosquitoes is a measure of practical importance, and it is an advantage to have a screen of forest between breeding places and human habitations. If such a natural defence exists it should on no account be removed, as has been done on occasions with disastrous results †

(See *Bibliography*, Nos 124-151)

7. The Use of Bats as Mosquito-destroyers.—That bats destroyed mosquitoes has been known for centuries, and the fact was mentioned by Pliny in his work on natural history. Campbell¹⁴⁹ in 1909 built a bat roost at San Antonio, Texas, the object being to breed bats which would destroy the local mosquitoes and would also prove a paying proposition on account of the production of guano. Subsequently the experiment was repeated in Italy, but without success the chief difficulty being that the bats

* As distinct from low jungle or scrub

† The question of planting shade giving trees and plants as an anti larval measure is discussed later (p. 36)

PART III.

MEASURES DIRECTED AGAINST THE LARVÆ OF MOSQUITOES.

A.—DESCRIPTION OF METHODS USED

Before describing the methods employed in dealing with larvæ, it is necessary to point out the extreme importance of a careful preliminary survey of the area in question so that it may be ascertained which species of anophelines are present, and where the breeding places of each are situated. Attention may thus be concentrated on measures to deal with the breeding places of the principal malaria carriers. For example, it is worse than useless to embark on expensive measures to drain a collection of water that may be harbouring either no larvæ at all or only those of harmless mosquitoes whilst the less obvious breeding places of the malaria carriers are overlooked.

Attention is also drawn to the fact that, when the favourite breeding places of any particular species have been eliminated, it will tend to breed in other situations so that careful periodical inspections of the area concerned are essential.

A point of great practical importance is the extent of the area over which it is necessary to practice anti larval measures. No strict rules can be laid down on this point, for a great number of factors must be taken into consideration, and the final decision will depend on a close study of the local conditions prevailing, such as the species of malaria carrying anophelines concerned, the intensity of breeding, the direction of the prevailing wind, etc.

As a general rule, in India, anti larval measures should be undertaken for a distance of at least half a mile from the periphery

anophelines close to the sleeping place of the human beings, in spite of the proximity of the fire, and the much greater accessibility of the cattle. He considers that the presence of the cattle in this region gives very little protection.

In India especially in the Punjab, a season of heavy rainfall following a year of drought is frequently the precursor of an outbreak of malaria in epidemic form. Cragg⁵⁷⁸ has suggested that since the number of cattle which normally form a protection to the population is decreased in proportion to the severity of the preceding drought whilst the number of anophelines is increased by the heavy rains, this may be an important factor in the genesis of these epidemics. He further suggested that the great extension of irrigation works primarily intended as a protection against famine, may prove to be in itself a great anti malarial measure for with an assured supply of water there will be sufficient fodder to keep the cattle alive in times of drought.

Close association with cattle does not always prevent an epidemic of malaria. In the Punjab epidemic of 1908 the cattle zone of Amritsar city was one of the worst epidemic areas. Fry⁵⁴⁰ however points out that though those living in this zone suffered owing to the presence of cattle the rest of the town was probably protected thereby from the abnormal mosquito population. He suggests that if cattle sheds were arranged on the outskirts of a village instead of indiscriminately as is usually the case, the inhabitants would be more free from the attacks of mosquitoes.

Fermi⁵³⁸ states that if animal prophylaxis is to be effective the cattle sheds should be situated at a distance of 50 to 100 yards from human dwellings.

A very large amount of literature dealing with the subject of zoophilism has appeared during the last 10 years. It is impossible to include a complete bibliography, but references to a number of the more important papers are given at the end of the book.

(See *Bibliography*, Nos 522-567.)

this connection C J Wilson^{5 6} as the result of experiments with a specially constructed trap made with mosquito netting in the Federated Malay States came to the conclusion that the result of trap catches was more valuable as an indication of the species of anophelines frequenting dwellings than the results of catches by collectors. Out of 1526 mosquitoes caught in the trap in some coolie lines in 12 nights 167 were anophelines of which 22 were *A. maculatus*. Two trained collectors searched in the lines each morning during the same period and caught 564 mosquitoes of which only 25 were anophelines and none were *A. maculatus* *.

A record is kept of the number of each species taken in each catching station. This is a most valuable means of testing the efficiency of control measures and will frequently lead to the detection of some breeding place which would otherwise be overlooked.

After the usual preliminary malaria survey has been carried out the following procedure is suggested —

(1) First select catching stations one at least being situated on each side of the area to be protected. Regular catching should preferably be started at least six weeks before the beginning of the malaria season. Indeed it would be better still if catching and other observations could be carried out for one complete season before instituting control measures but this is seldom possible in practice.

(2) Institute rigid anti larval measures over all actual and potential breeding places of malaria carrying anophelines for a distance of one quarter of a mile from the periphery of the area to be protected. Control measures should be initiated *before* the annual increase in the numbers of adult malaria carrying anophelines occurs, i.e. usually about one month before the commencement of the malaria season.

* The experiments here referred to were actually carried out by Mr B A R Guter the collector sleeping inside the trap.

of the area to be controlled. This may have to be extended in certain directions if breeding of malaria carrying anophelines outside this radius is very intense or if there is a strong prevailing wind from a particular quarter. On the other hand it may not always be necessary to undertake rigid control measures for half a mile in every direction. Thus Senior White⁸³ obtained excellent results by instituting anti larval measures for a quarter mile radius round certain camps in a very malarious area in the Vizagapatam Agency (Madras Presidency India) where a railway was being constructed the area being extended in one direction nearly three quarters of a mile from the centre of the main camp on account of the prevailing wind.

On the other hand Hackett⁴¹⁸ in Italy has found it necessary to control mosquito breeding for a distance of 3 kilometres (nearly 2 miles) from the periphery of the area requiring protection. Evidently *A. maculipennis* is a stronger flier than most of the malaria carrying anophelines of India with the possible exception of *A. sinica*.

Many experiments have been carried out with a view to ascertaining the distance of flight of anophelines and it has been shown that on occasions certain species will fly for distances of several miles. But in considering the size of the area over which it is necessary to practice anti larval measures it must be remembered as Ross³⁷ has pointed out that it is not a question of the power of flight of anophelines or how far they can fly but how far they actually do fly on the average.

In instituting control measures much useful information may be obtained by the use of catching stations or observation stations. Certain shelters are selected in various quarters of the camp village or other locality which is to be protected. These are visited regularly (daily if possible during the first season of control) and the same collector catches as many adult anophelines as possible in a given time say 15 minutes in each spot. Or traps may be placed in selected places and the catches collected daily. In

particular part of the area and attention can then be mainly directed to these special places

Temporary measures such as the application of paris green oil etc should be discontinued at the end of the usual period of the malaria season in the particular locality concerned To continue them after this date will merely in the author's opinion, result in a useless expenditure of labour and money

Other things being equal permanent measures i.e the complete abolition of breeding places should be adopted wherever possible Temporary measures such as oiling etc should only be employed (a) when the cause of breeding is temporary (e.g. during the construction of docks railways etc) or (b) when the cost of permanent measures is prohibitive It may be remarked here that permanent measures although they may involve a greater initial outlay than those of a temporary nature frequently prove more economical in the long run owing to the constantly recurring expenses of the latter and the necessity for maintaining a large inspecting staff

1 Clearing of Jungle* or Scrub—This is of value in the following respects

- (a) It removes the sheltering places of adult mosquitoes
- (b) It promotes evaporation and therefore the drying up of collections of water
- (c) It discloses breeding places which otherwise may be overlooked

A word of warning on this subject is however necessary Certain dangerous malaria carrying species of anophelines e.g. *A. maculatus*, *A. minimus*† and *A. ludlowi* prefer breeding places exposed to bright sunshine The indiscriminate clearing

* Considerable confusion has been caused in India in the use of the word jungle some authors alluding to forests of high trees by this term others to low bush or scrub It is used in the latter sense here

† This is the case in India but in the Philippine Islands *A. minimus* is said to favour streams heavily shaded by bamboo (*Neliazis**)

(iii) If any particularly dangerous breeding places such as a ravine stream or seepage outcrop exist between the quarter mile and half mile radius control these also

(iv) Scrutinize the daily records from the catching stations. The capture of a considerable number of malaria carrying anophelines in any particular catching station may indicate that control measures must be extended in that direction beyond the half mile radius (probably in the direction from which the prevailing wind is blowing) or that some particular breeding place has been overlooked

(v) Wherever possible catching stations should also be instituted in some village in the neighbourhood where the conditions are as nearly as possible the same as in the area under control but where no anti-malarial measures are being carried out. A record should be kept of anopheline catches, spleen rates, morbidity and mortality rates etc. in this village which may be compared with those observed in the controlled area. It must be remembered that the incidence of malaria like that of other diseases may vary very considerably in different years and if no records are kept of an uncontrolled area as well as of the one where anti-malarial measures have been instituted entirely erroneous conclusions may be reached as to the efficacy of the campaign. The danger of over-estimating the good effects of anti-malaria measures is especially great because as a rule such measures are undertaken during the period immediately following an unusually severe malaria year when the incidence of the disease would in any case tend to become less.

Our aim is of course to control the smallest possible area that will give good results. Every hundred yards added to the radius of the controlled area means a large increase in expenditure. It is usually found that expenses incurred in anti-malaria measures during the first year can be very considerably reduced during subsequent malaria seasons. It may for instance be found that all the malaria carrying anophelines are breeding in some

particular part of the area, and attention can then be mainly directed to these special places

Temporary measures, such as the application of paris green, oil, etc., should be discontinued at the end of the usual period of the malaria season in the particular locality concerned. To continue them after this date will merely, in the author's opinion, result in a useless expenditure of labour and money

Other things being equal permanent measures, i.e., the complete abolition of breeding places should be adopted wherever possible. Temporary measures such as oiling etc., should only be employed (a) when the cause of breeding is temporary (e.g., during the construction of docks railways etc.), or (b) when the cost of permanent measures is prohibitive. It may be remarked here that permanent measures although they may involve a greater initial outlay than those of a temporary nature frequently prove more economical in the long run, owing to the constantly recurring expenses of the latter and the necessity for maintaining a large inspecting staff

1. Clearing of Jungle* or Scrub—This is of value in the following respects —

- (a) It removes the sheltering places of adult mosquitoes
- (b) It promotes evaporation and therefore the drying up of collections of water
- (c) It discloses breeding places which otherwise may be overlooked

A word of warning on this subject is however necessary. Certain dangerous malaria carrying species of anophelines, e.g., *A. maculatus*, *A. minimus*† and *A. ludlowi*, prefer breeding places exposed to bright sunshine. The indiscriminate clearing

* Considerable confusion has been caused in India in the use of the word 'jungle,' some authors alluding to forests of high trees by this term others to low bush or scrub. It is used in the latter sense here

† This is the case in India, but in the Philippine Islands *A. minimus* is said to favour streams heavily shaded by bamboo (Mielzinsky)

of jungle in places where such species exist may therefore be followed by disastrous results. This occurred in a district in the Federated Malay States where the wholesale clearing of jungle though proving inimical to the shade loving *A. umbrosus* provided excellent breeding places for the more dangerous *A. maculatus*.

In Assam where *A. minimus* is the principal malaria carrier an immense amount of harm has been caused by the clearing of jungle from the edges of streams and experiments are now being carried out with a view to finding the most suitable shade giving plants for use along the banks of streams in tea gardens. For this purpose it is important that the plants and trees shall give dense shade and that they shall be evergreen. It has been found that *duranta* (the thorny privet hedge) and *eupatorium* are suitable for narrow channels whilst for broader streams the jack fruit tree (*Artocarpus integrifolia*) appears to be efficacious (Ramsay⁸⁰).

The clearing of forests has also resulted in serious malaria epidemics in the high plateau region of Argentina where *A. albivittatus*, *A. tarsimaculatus* and *A. argyritarsis* all prefer sunlit water for their breeding places.

2 Drainage—It is impossible to treat this large and important subject fully here. Some of the chief points will be dealt with and for further details the reader is referred to the list of references given in the *Bibliography* under this heading.

(a) *Open drains*—These should be narrow and deep rather than broad and shallow. Their banks should be kept clear of vegetation and sloped to an angle of about 45 degrees. Tributaries should enter at an acute angle or curve where possible and not at right angles in order to lessen the deposition of silt and debris at the point of junction. The bottom of the drain should be rounded and not V shaped.

In dealing with swampy areas drains may be made either parallel with one another or arranged in 'herring bone' fashion.

The former method is the more economical. For example in draining an area of 14 acres with lines of drains laid 100 feet apart, the herring bone system would require 300 feet more of drain than the parallel¹⁸¹ (Fig 5). In British Guiana there is employed a method of making straight parallel drains and cultivating the intervening areas the plants grown taking up a large amount of water. In the case of hill foot seepages the best method is to construct a system of contour drains to catch seepage at the point at which it arises (see p. 89).




FIG. 5

Two systems of laying out drains in areas of equal size. In both the drains are the same distance apart, but the length of drain in A is considerably greater than in B (after King¹⁸¹).

Drains should be as few and as short as possible and the fall of the ground should be carefully considered before commencing work. The excavation for drainage should start at the outfall end. Sharp bends should be avoided wherever possible. The main drain should be constructed first and the tributaries afterwards. In some cases the use of dynamite has been found helpful in the construction of drains¹⁸⁵.

Open drains may be either *pukka* i.e. lined with concrete, brick stone etc. or *kuchcha* i.e. open earth drains. A *pukka*

drain should never be constructed without first making a kuchcha one to determine the requisite depth of the drain and to see whether the flow is satisfactory

Pukka drains should be made with a central deeper channel or 'cunette' to take the water along quickly when its level is low and to minimize the extent of the breeding area. At the point of junction with a side channel the opposite side of the drain should be strengthened and raised to prevent overflow. 'Weep holes' should be made in the side of the drain so that the subsoil water may get into it. These should slope downwards towards the bottom of the drain. It is also an advantage to construct key walls at right angles to the drain at intervals especially where there is a curve in its course to prevent water from outside tearing away the earth supporting the side walls (Fig 6). Key walls




FIG 6

Cross section of key wall (after Le Prince)

should extend 6 inches to a foot or more into the ground below the bottom lining of the drain and in all cases weep holes should be provided just above the key wall. In wide drains the side walls may be almost vertical.

It is not always necessary to line the entire drain the lining of the bottom and sides up to 3 inches above the normal water line for small ditches being sufficient as a rule. A drain may be roughly lined with flat stone, the interspaces being filled in with small stone

and sealed roughly with cement mortar. Where flat stone is not available concrete made with gravel or small stone, in a layer of about 2 inches in thickness and reinforced with 2 inch mesh wire netting, may be used.

In places where a drain has to pass beneath a road by means of a culvert, the grade of the drain should be increased, to prevent the accumulation of silt or débris. The bottom of the culvert should be lined with stone or concrete. A screen of vertical rods may be provided at the entrance of the pipe or culvert, and in the case of an earth drain the bottom should be lined for a distance of about 6 feet with flat stone or timber at the point of discharge, so as to prevent erosion and the formation of a pocket.

In the case of kuchcha drains, if the flow of water is very swift undermining of the banks may occur, or if there is some temporary obstruction excessive local scouring may ensue removing soil from below the grade line of the bottom of the drain and causing a 'pot hole'. Subsequently, if the rest of the drain becomes dry (as in storm water drains), pools will remain in such situations, which may become anopheline breeding places. A small temporary channel should then be made to connect these pools and drain off the water. It is sometimes possible to effect this rapidly in the case of a drain with a muddy bottom by dragging a small log downstream in its bed.

The necessity for repeated re grading, cleaning and oiling of open earth drains makes their upkeep expensive. They are of no use in localities subject to very heavy rainfall as when this occurs the system will be wrecked. Lined drains last much longer, are more easily cleaned, require less inspection and are frequently ultimately less costly than open earth drains¹⁸⁴.

(b) *Subsoil drains*—The drain in this case is formed by a series of unjointed tile pipes laid end to end close together in trenches beneath the ground, the water entering from below at the points of junction of the pipes. If soft spots are found in the bottom of the trench (which must of course be properly graded),

drain should never be constructed without first making a kuchcha one to determine the requisite depth of the drain and to see whether the flow is satisfactory.

Pukka drains should be made with a central deeper channel or 'cunette' to take the water along quickly when its level is low and to minimize the extent of the breeding area. At the point of junction with a side channel the opposite side of the drain should be strengthened and raised to prevent overflow. 'Weep holes' should be made in the side of the drain so that the subsoil water may get into it. These should slope downwards towards the bottom of the drain. It is also an advantage to construct 'key walls' at right angles to the drain at intervals especially where there is a curve in its course to prevent water from outside tearing away the earth supporting the side walls (Fig 6). Key walls




FIG 6

Cross section of key wall (after Le Prince)

should extend 6 inches to a foot or more into the ground below the bottom lining of the drain and in all cases weep holes should be provided just above the key wall. In wide drains the side walls may be almost vertical.

It is not always necessary to line the entire drain the lining of the bottom and sides up to 3 inches above the normal water line for small ditches being sufficient as a rule. A drain may be roughly lined with flat stone, the interspaces being filled in with small stone

and sealed roughly with cement mortar. Where flat stone is not available concrete made with gravel or small stone, in a layer of about 2 inches in thickness and reinforced with 2 inch mesh wire netting may be used.

In places where a drain has to pass beneath a road by means of a culvert the grade of the drain should be increased, to prevent the accumulation of silt or debris. The bottom of the culvert should be lined with stone or concrete. A screen of vertical rods may be provided at the entrance of the pipe or culvert and in the case of an earth drain the bottom should be lined for a distance of about 6 feet with flat stone or timber at the point of discharge, so as to prevent erosion and the formation of a pocket.

In the case of kuchcha drains if the flow of water is very swift undermining of the banks may occur or if there is some temporary obstruction excessive local scouring may ensue removing soil from below the grade line of the bottom of the drain and causing a 'pot hole'. Subsequently, if the rest of the drain becomes dry (as in storm water drains) pools will remain in such situations, which may become anopheline breeding places. A small temporary channel should then be made to connect these pools and drain off the water. It is sometimes possible to effect this rapidly in the case of a drain with a muddy bottom by dragging a small log downstream in its bed.

The necessity for repeated re grading, cleaning and oiling of open earth drains makes their upkeep expensive. They are of no use in localities subject to very heavy rainfall as when this occurs the system will be wrecked. Lined drains last much longer, are more easily cleaned, require less inspection and are frequently ultimately less costly than open earth drains¹⁸⁴.

(b) *Subsoil drains*—The drain in this case is formed by a series of unjointed tile pipes laid end to end close together in trenches beneath the ground, the water entering from below at the points of junction of the pipes. If soft spots are found in the bottom of the trench (which must of course be properly graded),

drain should never be constructed without first making a kutchra one to determine the requisite depth of the drain and to see whether the flow is satisfactory.

Pukka drains should be made with a central deeper channel or 'cunette' to take the water along quickly when its level is low and to minimize the extent of the breeding area. At the point of junction with a side channel the opposite side of the drain should be strengthened and raised to prevent overflow. 'Weep holes' should be made in the side of the drain so that the subsoil water may get into it. These should slope downwards towards the bottom of the drain. It is also an advantage to construct 'key walls' at right angles to the drain at intervals especially where there is a curve in its course to prevent water from outside tearing away the earth supporting the side walls (Fig. 6). Key walls




FIG. 6

Cross section of key wall (after Le Prince.)

should extend 6 inches to a foot or more into the ground below the bottom lining of the drain and in all cases weep holes should be provided just above the key wall. In wide drains the side walls may be almost vertical.

It is not always necessary to line the entire drain the lining of the bottom and sides up to 3 inches above the normal water line for small ditches being sufficient as a rule. A drain may be roughly lined with flat stone, the interspaces being filled in with small stone

and sealed roughly with cement mortar. Where flat stone is not available concrete made with gravel or small stone in a layer of about 2 inches in thickness and reinforced with 2 inch mesh wire netting may be used.

In places where a drain has to pass beneath a road by means of a culvert the grade of the drain should be increased to prevent the accumulation of silt or debris. The bottom of the culvert should be lined with stone or concrete. A screen of vertical rods may be provided at the entrance of the pipe or culvert and in the case of an earth drain the bottom should be lined for a distance of about 6 feet with flat stone or timber at the point of discharge so as to prevent erosion and the formation of a pocket.

In the case of kuchcha drains if the flow of water is very swift undermining of the banks may occur or if there is some temporary obstruction excessive local scouring may ensue removing soil from below the grade line of the bottom of the drain and causing a 'pot hole'. Subsequently if the rest of the drain becomes dry (as in storm water drains) pools will remain in such situations, which may become anopheline breeding places. A small temporary channel should then be made to connect these pools and drain off the water. It is sometimes possible to effect this rapidly in the case of a drain with a muddy bottom by dragging a small log downstream in its bed.

The necessity for repeated re grading, cleaning and oiling of open earth drains makes their upkeep expensive. They are of no use in localities subject to very heavy rainfall as when this occurs the system will be wrecked. Lined drains last much longer, are more easily cleaned, require less inspection and are frequently ultimately less costly than open earth drains¹⁸⁴.

(b) *Subsoil drains*—The drain in this case is formed by a series of unjointed tile pipes laid end to end close together in trenches beneath the ground, the water entering from below at the points of junction of the pipes. If soft spots are found in the bottom of the trench (which must of course be properly graded),

drain should never be constructed without first making a kuchcha one to determine the requisite depth of the drain and to see whether the flow is satisfactory

Pukka drains should be made with a central deeper channel or 'cunette' to take the water along quickly when its level is low and to minimize the extent of the breeding area At the point of junction with a side channel the opposite side of the drain should be strengthened and raised to prevent overflow 'Weep holes' should be made in the side of the drain so that the subsoil water may get into it These should slope downwards towards the bottom of the drain It is also an advantage to construct key walls at right angles to the drain at intervals especially where there is a curve in its course to prevent water from outside tearing away the earth supporting the side walls (Fig 6) Key walls




FIG 6

Cross section of key wall (after Le Prince)

should extend 6 inches to a foot or more into the ground below the bottom lining of the drain and in all cases weep holes should be provided just above the key wall In wide drains the side walls may be almost vertical

It is not always necessary to line the entire drain the lining of the bottom and sides up to 3 inches above the normal water line for small ditches being sufficient as a rule A drain may be roughly lined with flat stone the interspaces being filled in with small stone

and sealed roughly with cement mortar. Where flat stone is not available concrete made with gravel or small stone, in a layer of about 2 inches in thickness and reinforced with 2 inch mesh wire netting may be used.

In places where a drain has to pass beneath a road by means of a culvert the grade of the drain should be increased, to prevent the accumulation of silt or debris. The bottom of the culvert should be lined with stone or concrete. A screen of vertical rods may be provided at the entrance of the pipe or culvert, and in the case of an earth drain the bottom should be lined for a distance of about 6 feet with flat stone or timber at the point of discharge, so as to prevent erosion and the formation of a pocket.

In the case of kuchcha drains, if the flow of water is very swift undermining of the banks may occur or if there is some temporary obstruction excessive local scouring may ensue removing soil from below the grade line of the bottom of the drain and causing a 'pot hole'. Subsequently if the rest of the drain becomes dry (as in storm water drains) pools will remain in such situations, which may become anopheline breeding places. A small temporary channel should then be made to connect these pools and drain off the water. It is sometimes possible to effect this rapidly in the case of a drain with a muddy bottom by dragging a small log downstream in its bed.

The necessity for repeated re grading, cleaning and oiling of open earth drains makes their upkeep expensive. They are of no use in localities subject to very heavy rainfall as when this occurs the system will be wrecked. Lined drains last much longer, are more easily cleaned, require less inspection and are frequently ultimately less costly than open earth drains¹⁸⁴.

(b) *Subsoil drains*—The drain in this case is formed by a series of unjointed tile pipes laid end to end close together in trenches beneath the ground, the water entering from below at the points of junction of the pipes. If soft spots are found in the bottom of the trench (which must of course be properly graded),

drain should never be constructed without first making a kuchcha one to determine the requisite depth of the drain and to see whether the flow is satisfactory

Pukka drains should be made with a central deeper channel or 'cunette' to take the water along quickly when its level is low and to minimize the extent of the breeding area. At the point of junction with a side channel the opposite side of the drain should be strengthened and raised to prevent overflow. 'Weep holes' should be made in the side of the drain so that the subsoil water may get into it. These should slope downwards towards the bottom of the drain. It is also an advantage to construct key walls at right angles to the drain at intervals especially where there is a curve in its course to prevent water from outside tearing away the earth supporting the side walls (Fig 6). Key walls




FIG 6

Cross section of key wall (after Le Prince)

should extend 6 inches to a foot or more into the ground below the bottom lining of the drain and in all cases weep holes should be provided just above the key wall. In wide drains the side walls may be almost vertical.

It is not always necessary to line the entire drain the lining of the bottom and sides up to 3 inches above the normal water level for small ditches being sufficient as a rule. A drain may be roughly lined with flat stone, the interspaces being filled in with small stone

and sealed roughly with cement mortar. Where flat stone is not available concrete made with gravel or small stone, in a layer of about 2 inches in thickness and reinforced with 2 inch mesh wire netting may be used.

In places where a drain has to pass beneath a road by means of a culvert, the grade of the drain should be increased to prevent the accumulation of silt or débris. The bottom of the culvert should be lined with stone or concrete. A screen of vertical rods may be provided at the entrance of the pipe or culvert and in the case of an earth drain the bottom should be lined for a distance of about 6 feet with flat stone or timber at the point of discharge so as to prevent erosion and the formation of a pocket.

In the case of kuchcha drains if the flow of water is very swift undermining of the banks may occur or if there is some temporary obstruction excessive local scouring may ensue removing soil from below the grade line of the bottom of the drain and causing a 'pot hole'. Subsequently if the rest of the drain becomes dry (as in storm water drains) pools will remain in such situations, which may become anopheline breeding places. A small temporary channel should then be made to connect these pools and drain off the water. It is sometimes possible to effect this rapidly in the case of a drain with a muddy bottom by dragging a small log downstream in its bed.

The necessity for repeated re grading, cleaning and oiling of open earth drains makes their upkeep expensive. They are of no use in localities subject to very heavy rainfall as when this occurs the system will be wrecked. Lined drains last much longer, are more easily cleaned, require less inspection and are frequently ultimately less costly than open earth drains.¹⁸⁴

(b) *Subsoil drains*—The drain in this case is formed by a series of unjointed tile pipes laid end to end close together in trenches beneath the ground, the water entering from below at the points of junction of the pipes. If soft spots are found in the bottom of the trench (which must of course be properly graded),

drain should never be constructed without first making a kutchcha one to determine the requisite depth of the drain and to see whether the flow is satisfactory

Pukka drains should be made with a central deeper channel or 'cunette' to take the water along quickly when its level is low and to minimize the extent of the breeding area. At the point of junction with a side channel the opposite side of the drain should be strengthened and raised to prevent overflow. 'Weep holes' should be made in the side of the drain so that the subsoil water may get into it. These should slope downwards towards the bottom of the drain. It is also an advantage to construct 'key walls' at right angles to the drain at intervals especially where there is a curve in its course to prevent water from outside tearing away the earth supporting the side walls (Fig. 6). Key walls




FIG. 6

Cross section of key wall (after Le Prince)

should extend 6 inches to a foot or more into the ground below the bottom lining of the drain and in all cases weep holes should be provided just above the key wall. In wide drains the side walls may be almost vertical.

It is not always necessary to line the entire drain the lining of the bottom and sides up to 3 inches above the normal water line for small ditches being sufficient as a rule. A drain may be roughly lined with flat stone, the interspaces being filled in with small stone

(one fifth for main drains over 6 inches in diameter) during ordinary dry weather (at least 2 or 3 days after moderate rainfall) they will be sufficient to do their work during floods (Evans¹⁶⁸) The pipes should be laid in an absolutely straight line with as few changes of gradient as possible

Greasy water and house waste must not be allowed to discharge into any part of the system Where pipes come near the surface proper bridge crossings are necessary to protect them from being crushed by carts etc The outlet of the subsoil drain must be well above the high water level of the stream lake or ditch into which it discharges Inspections should be made to see that the outlets do not become clogged with silt or other deposit Metal rods placed vertically at the outlet will prevent the entry of small animals which might die in the pipes and cause blockage

Subsoil drainage in connection with mosquito control is used for the following purposes —

- (i) To lower the ground water so that pools of surface water will be more readily absorbed
- (ii) To intercept seepage
- (iii) To deal with hill streams in ravines

(i) Where this method is used for lowering the subsoil water the lines of pipes are usually spaced from 50 to 150 feet apart They are generally laid from 2 to 4 feet deep If laid nearer to the surface the water is drained away more rapidly but the area drained is less extensive The pipes are laid in the bottom of a perfectly graded narrow trench with their ends abutting one another closely and are then covered with clay or earth A grade of 1 foot or more per 400 feet is desirable¹⁶⁴

(ii) To intercept seepage the drain is constructed above the line of outcrop at the time of maximum flow approximately at right angles to the flow deep enough to collect the seepage and with a grade of not less than 1 foot in 200 feet The pipes are laid with open joints or about one-eighth to one fourth inch apart and the trench filled with stones The uppermost layer should be

stones are rammed into place until a solid foundation is obtained. Higler¹⁰⁸ has found it useful where gravel is available to lay the tile pipe on about 10 cm. depth of gravel and to cover it with 10-20 cm. of the same material. He has found that in pipes so laid clogging has been negligible whilst in others it is of periodic occurrence. The laying of the pipes should be commenced at the outlet of the drain and continued upwards as the trench is made. In fine easily disintegrated soils the pipe joints should be covered with a wrapping of canvas, palm leaves or other material to prevent clogging. The pipes should be of circular bore from 3 to 12 inches in diameter and 1 to 2 feet in length. As a rule the internal diameter




FIG. 7

Cross section of three types of subsoil drains. A Tile drain (after Le Prince⁹) B Stone drain C Pole drain

of the pipes should not be less than 4 inches even for hill foot drains whilst main drains are usually from 6 to 8 inches in diameter. Where more capacity is required two or more drains may be laid side by side. The thickness of the drain should be three eighths of an inch for pipes of 3 or 4 inches in diameter, half an inch for those of 6 inches in diameter and five-eighths of an inch for those of 8 inches in diameter. It is impossible to lay down rigid rules for the capacity of the system but in general it may be said that if the pipes are calculated to run not more than one quarter full

disadvantage is its cost, which is a grave objection, especially when dealing with ravine streams. Senior White⁸ remarks that in the case of streams which run only for two months or so after the slackening of the monsoon, oiling is probably more economical.

The cost of subsoil hill land drainage in and around the town of Kuala Lumpur, Federated Malay States, was about £5 16s 0d per acre. The cost of draining agricultural hill land in the Malay States ranges from £1 15s 0d to £4 10s 0d per acre. The cost of maintenance has not exceeded 10 per cent of the first cost in the year following construction and 5 per cent in the succeeding years. It must be realized that maintenance must commence the day after construction is completed (L'vans¹⁶⁸).

In place of tile pipes subsoil drains may be constructed with sides, roof and floor made of flat stones (Fig. 7 B). Drains of this type varying in size from 6 inches to 12 inches wide and from 9 inches to 15 inches deep were installed at Meenglas, Bengal in 1917-1919 at an average cost of 11½ annas per foot run, and after 8 years working with very little cost in maintenance they were found to be in as good condition as ever¹⁹.

Where stones are not available larch or other poles may be laid lengthwise in the bottom of the drain, one on each side and one on the top forming a conduit (Fig. 7 C).

Another method is simply to make deep trenches and fill them with stones, but in this case it is necessary to dig them up and clean out the silt at intervals.

(c) *Vertical drainage*. This may prove a useful and economical measure especially in the case of swamps and marshes. The bed of the marsh is probably of silt or clay which retains the water, but underneath this there may be a permeable or fissured rock which will give drainage. In order to drain the marsh a pit is sunk down to the permeable rock, the exposed surface of which is blasted. The pit should be filled with rock filling, a vertical line of pipe being carried to the level of the marsh bottom. The pipe should be surrounded with stone or gravel. Spigot and socket sewer pipes or iron pipes coated can be used, or a circular pit built

of small sized stones, and should extend 2 or 3 inches above the surface of the ground¹⁸⁴

(iii) In dealing with hill streams in ravines, it is necessary that the pipes should be buried *at least* 3 feet deep*. The pipes should be covered by a layer of heavy stone, the smaller stones being placed above as before, but in order to prevent silt from reaching the pipes and blocking them, it is necessary for the topmost layer itself (i.e., the floor of the ravine) to be of silt. In wet climates a layer of grass will grow over this, and will prove efficacious against scouring. It is most important that trees




FIG. 8

Subsoil tile drain longitudinal section
(after Le Prince¹⁸⁴)

growing within a distance of some 40 feet from the pipes shall be rooted out, otherwise their roots will grow down into the pipe junctions and block them.

The advantages claimed for subsoil drainage as compared with open drains are that it is self cleaning maintains itself permits of rapid inspection needs very little attention, requires no oblong and permits of no exposure of water accessible to mosquitoes. It should be pointed out, however that the laying of pipes should as a rule, only be undertaken by a skilled engineer. Its principal

* A Circular ** issued by the Public Works Department F. M. S. in 1911 laid down that there should be not less than 4 feet of cover over the pipes.

How Oil Kills Mosquito Larvæ

It was originally supposed that the lethal effect of oil on larvæ was due solely to the fact that it suffocated them by cutting off their air supply. A great deal of investigation has since been carried out as to the manner in which oil kills larvæ, the various theories put forward being as follows —

- (i) Suffocation of the larvæ by producing a surface film which cuts off their supply of air
- (ii) Blocking of the respiratory tubes by particles of oil
- (iii) Toxic action of oil vapours
- (iv) Toxic action of the oil directly by contact
- (v) Toxic action of the oil indirectly by solubility in the water
- (vi) Reduction of the surface tension making it difficult for the larvæ to remain at the surface and thus causing them to drown

Freeborn and Atsatt³⁴⁰ found that the toxicity of oils increases with their volatility and that the volatile constituents contain the principles which produce the primary lethal effect but that in the case of those oils with negligible volatility (boiling point over 250°C) the lethal action may be due either to blocking of the respiratory tubes, poisoning by contact or even by suffocation. They did not consider that the reduction of surface tension or toxicity from solubility in water had any effect.

More recent researches have in the main confirmed these observations. Green³⁴⁵ showed that the rapidity with which larvæ die depends on the volatility and toxicity of the oil, that the larvæ obtain their fatal dose of oil in a very short time (less than 1 second with petrol and within 1 minute with an absolutely non-toxic oil, Nujol) and that *Culex* larvæ take a very much longer time to die than *Aopheles* (6 to 8 times as long).

Hacker³⁵⁰ pointed out that oil frequently kills larvæ without the production of a complete film. He laid stress on the importance of differential wetting in the susceptibility of larvæ to the effects of oil. Oil on the surface of the water acts like a

up in dry stone masonry. A circular strainer should be put in round the mouth of the sinkage (Home¹⁷⁷). Probably several of these pits will be necessary if the marsh is a large one. A certain amount of grading will be required in the marsh leading to the sink holes (Fig. 9).

(See Bibliography, Nos. 155-195.)




FIG. 9

Vertical drainage (after Home¹⁷⁷)

3. Oiling.—The use of oil to diminish the number of mosquitoes was recommended in a note in *The American Advertiser*, Philadelphia, in its issue of August 29th 1793 (McFarland³⁶³). It was also suggested in a work entitled 'Omniapa Otiosores,' said to have been written by R. Southey, which was published in London in 1812. The first published account of scientifically conducted experiments with oil as a mosquito larvicide appears to have been that of Howard³⁵³, which appeared in 1892.

The addition of small quantities of a vegetable oil also greatly increases the spreading of mineral oils. According to Hacker³⁵⁰ vegetable oils probably spread by virtue of the free fatty acids which they contain. Leak³⁶² recorded that the addition of 1 per cent of castor oil to kerosene increased the spread 25 times. He found that 2 ozs. of kerosene covered an area 5 yards in diameter, whilst on the addition of 1 per cent of castor oil an unbroken film 30 yards in diameter was formed.

If the oil contains too much active material (OH groups, COOH groups etc.) the excess leaves the oil, satisfies the affinity of the water for active groups and therefore prevents the oil from spreading. This is a point of practical importance, since the addition of too much cresol, vegetable oil etc., will actually decrease the spreading power instead of increasing it. A method of testing whether too much active material has been added to, or is contained in an oil is by shaking up a sample with water, if this water then prevents the spread of oil it has been inactivated by the excess of active material.

The spreading of oils is also increased by oxidation. Hacker³⁵⁰ showed that kerosene and crude oil increased in spreading power when exposed to air for 8 days. It was found by Johnson³⁵⁸ that oil soaked sawdust caused a better film when left for 24 hours before use, and also that certain waste oils from garages did not spread well until they were soaked with sawdust. Hacker considers that the improvement in spreading power was in this case probably due to oxidation and not to the filtering out of carbon particles by the sawdust, as Johnson suggested. Sollmann³⁷⁵ found that the spreading of kerosene was increased by adding asphalt and Hacker suggests that this may be because the tar causes oxidation of paraffins by catalytic action. Hacker also noted that the toxicity of kerosene was increased by oxidation as well as the spreading power.

The presence of dust or grease on the water, or of soap or soapy material in the water, prevents oil from spreading, and this has to be taken into account when considering the amount of

dry object, and attracts the spiracles (which are also dry) When contact takes place the oil is able to wet the spiracles and enter the breathing tubes Contact with oil is sufficient to kill larvæ

Oil also tends to deter the adults from depositing their eggs, whilst it kills the algae forming with them a dense floating mass which probably has a deleterious effect, and in any case reduces the food supply of the larvæ In its place there appears a closely felted alga attached to the stones or sand forming the bed of the stream or pool, the presence of which forms a useful indication as to whether oiling has been carried out effectually⁴⁵

The spreading power of oils

The popular opinion is that the lighter kerosenes spread better than the crude oils, but some authors state the reverse Hacker³⁵⁰, in his laboratory experiments, found great difficulty in producing uniform films from the same amount of oil dropped on the same amount of water in the same size Petri dish He explained this phenomenon by the formation in some cases of an 'invisible film' of greasy substances derived from the fingers during the preparation of the dish, inhibiting the spread of oil This may be the explanation of the conflicting statements which have been made regarding the spreading power of various oils

The spreading of oils, however, does not depend on viscosity, and a thin oil does not necessarily show better spreading powers than a thick Ginsburg³⁴⁹ has shown that the tar acids having hydroxyl (OH) groups (e.g., phenols, cresols, xlenols), increase the spreading power of mineral oils, as also do the monohydric alcohols pine oil and turpentine Cresols and xlenols proved more efficient than the other compounds, the addition of one gallon of crude cresol containing 95 per cent cresylic acids to 100 gallons of fuel oil greatly increasing the spreading and penetration of the oil on salt and fresh waters covered with dead organic matter and vegetation In laboratory experiments it has been found that the addition of 1 per cent cresol to fuel oil increases the spreading power by 50 per cent The duration of the oil film is also appreciably increased by this treatment

(a) *Kerosene oil*—This may be applied alone but there are certain objections to its use—

- (i) The film is so thin that a very slight disturbance of the water surface by floating debris projecting vegetation ripples caused by the wind or current etc., break the continuity of the film
- (ii) It is expensive and likely to be stolen by unscrupulous employees
- (iii) It is transparent and therefore likely to be wasted by the persons applying it because it is difficult to see whether the film is satisfactory
- (iv) There is risk of fire e.g. where sparks from a passing railway train may drop into it

The quantity recommended for use by Howard³⁵³ was one ounce to 15 square feet of surface the film produced lasting about 10 days

(b) *Crude oil and other unrefined oils*—The liquid fuel of the petroleum companies is not properly speaking crude oil but a refuse after some of the more volatile oils have been distilled off. Its composition varies considerably in different consignments. Watson⁴⁵ notes that some consignments can be used alone whilst others need the addition of one part of kerosene to 8 parts of liquid fuel before use with a sprayer. The dark tarry solution of extractions which is found at the bottom of the oil drums is toxic to larvae and is very useful for application in springs etc. The addition of from 1 to 2½ per cent of a vegetable oil castor coconut or whatever kind may be available locally will increase the spreading power of crude oil or liquid fuel to a great extent castor oil being the most effective

Klugler¹⁰⁸ in Palestine found that the most effective mixture was 1 part of crude oil to 4 parts of kerosene with from 0.1 to 0.2 per cent of castor oil added.

Williamson and Rajamoney³⁸³ suggested the use of a mixture containing equal parts of crude oil solar oil and rubber oil. Rubber oil does not increase the spreading power of the other two but

vegetable oil, etc., to be added. The presence of grease or soap explains why an oil which may give excellent results on natural waters will often lie in patches when applied to water containing drainage from dwellings.

Williamson and Rajamoney³⁸⁵ lay stress on the fact that *very* great spreading power is not always an advantage being only permissible in limited areas of pure water, and in the presence of weed or other factors which retard film formation. A very thin film of oil is an ineffectual destroyer of mosquito larvæ.

The choice of oil

There are many grades of oil on the market which may be used for mosquito destruction, ranging from the very light oils such as kerosene to the heavier forms known as crude oils. The choice of an oil will depend very largely on local conditions, thus, with high temperatures a thick oil is required, whilst in the presence of vegetation an oil with great spreading power is necessary. Again in the case of still water a heavy non-toxic oil applied in sufficient quantity to form a complete film will after some hours kill all the larvæ, whilst in the case of moving water a thin layer of rapidly spreading oil with a high toxicity is indicated.

As already stated the poisonous action of mineral oils is related to several factors—viscosity, volatility, and the presence of aromatic hydrocarbons or of certain natural poisons. A thin oil will enter the breathing tubes of the larvæ quicker than a thick, whilst as we have seen, a high volatility and a high percentage of aromatic hydrocarbons increase the lethal effect. Unrefined oils, such as fuel oil or diesel oil, contain small quantities of substances (organic acids, nitrogen and sulphur compounds) which appear to exert a poisonous effect. These are partly removed by refining processes, and a highly refined lubricating or paraffin oil will therefore prove less poisonous than a black fuel oil of the same thickness.

Barnes³²⁸ has found a mixture of 9 parts of crude oil with 1 of pine oil effective. The latter, which contains various alcohols and ketones, has a powerful soporific or paralysing effect on the larvæ, and also increases the spreading power of the crude oil.

(c) *Liquid paraffin* (liquid vaseline) has been used by Swellengrebel and his co workers³⁹³ in Holland. It is an almost colourless oil, costing about £1 per cwt. It is claimed for this oil that it does not need the destruction of vegetation, that its effects are not vitiated by wind or rain, that it does not evaporate and therefore has a lasting effect and that it does not prevent mosquitoes from depositing their eggs. It is also non toxic to fish. Although it costs five times as much as Paris green, its use in the smaller canals in Holland (where manual labour is very expensive) entails less expense, because it is so easily and quickly applied. Five or six c.c. of the preparation are used per square metre of water surface.

(d) 'Leron' a mixture of various mineral oils, has been used with success in Italy and elsewhere. It is claimed for this preparation that it is harmless to fish and cattle and that it gives satisfactory results in the proportion of 10 c.c. to the square metre of water surface, even in the presence of dense vegetation, it being sufficient to move the reeds, etc. a little to ensure uniform spread (Gratch³⁴⁴).

(e) *Petrol*, in the proportion of 24 ozs. for every 80 square feet of surface, has been recommended for use in wells, tanks etc. It is claimed that larvæ are immediately destroyed by this method, and that in 2 hours there is no smell and in 3 hours no taste of petrol in the water. It is suggested that the petrol should be applied in the evening³⁷¹.

Methods of applying oil

(i) *Spraying* by means of a spray can or knapsack sprayer. By this means the oil can be distributed to a distance of 20 to 30 feet from the operator. The sprayer when filled must not be too heavy, a pattern with about 2½ gallons capacity being suitable

greatly increases their toxicity. Solar oil, a heavy fuel oil derived from aromatic crude oils, probably imparts continuity to the films whilst the inclusion of crude oil is justified on account of its low cost, and the fact that it evaporates less quickly than either of the others thereby prolonging their continuance on the water surface.

Anti malarial oil (A M M) is a mixture containing diesel oil, solar oil and kerosene put up by the Asiatic Petroleum Company. Diesel oil is unrefined and non volatile at ordinary temperatures and is added to give a lasting effect. Poisonous substances also occur naturally in this oil³²⁵.

Pesterine (M D B) prepared by the Burmah Shell Oil Company, is another mixture of oils recommended for anti mosquito work. Its present price is Rs. 32.13 per 42 gallon drum. It is claimed that this is slightly superior to A M M.

Feegrade³³⁷ experimenting with various crude oil products of the Burma Oil Company, Rangoon found 'Bauxite extract' to be the most effective.

Walch and Bonne Wepster³⁸⁰ recommend the use of 'sludge' a waste product from the Batavian Petroleum Company, used in the proportion of $7\frac{1}{2}$ to 10 c.c. per square metre of water surface.

Waste motor oil ('crank case oil') has been tried extensively in America. Peterson and Ginsburg³⁶⁷ note that this usually needs straining and does not spread on water but may be made to do so by adding a tar acid containing 25 per cent cresylic acid or by adjusting the specific gravity to 32 or 34 Be. by adding light petroleum distillates. They state that the lasting quality of the oil mixture is 2 to 4 weeks twice as long as that of fuel oil and that the cost is only half that of the latter. Walch and Bonne Wepster³⁸⁰ also used waste motor oil and found it effective when mixed with 10 per cent kerosene and 1 to 2 per cent coconut or castor oil 10 to 20 c.c. of the mixture being applied per square metre of surface. They also obtained good results with a mixture of 3 parts waste oil and 1 part solar oil. Waste crank case oil destroys vegetation.

PLATE I



A O Sprayer with Double Action Pump (Ladywood No 28)



B High Pressure Oil Sprayer Four Oaks Kent

for use by an Indian coolie. There are two principal types of sprayer on the market: one fitted with a double acting pump which has to be worked continuously by hand (e.g. the Ladywood Sprayer No. 28) and the high pressure type (e.g. the 'Four Oaks' High Pressure Sprayer and the Misto Gem Sprayer No. 8) in which the pressure is pumped up to a standard figure marked on a gauge before use (Plate I). The latter type though more expensive is less fatiguing for the operator and leaves one hand free a great advantage in rough country. In America the Myers 3 gallon compressed air sprayer fitted with a 2½ feet extension and a graduating vermorel nozzle has been found satisfactory.

The sprayers should preferably be fitted with leather or fibre valves and flexible metallic hose as petroleum oils cause ordinary rubber to perish. In any case since leaks at the base of the hose are common it is advisable to obtain a spare hose as well as a spare set of washers when ordering the machine. The choice of nozzle is important. Senior White⁷⁸ recommends a pattern which will deliver a cone spray about 18 inches in diameter 4 feet in front of the nozzle. He finds the 'Marvellous' nozzle and the I.T. 1 attachment with 'Marvellous' nozzles of the 'Four Oaks' Company the most suitable. A defect of many nozzles is that they deliver too much oil per minute so that the operator must either proceed at a run or waste oil.

The oil should be filtered into the sprayer through a sieve (which is usually supplied with the machine) and after each day's use the apparatus should be washed out by pouring a pint or so of kerosene into the container and blowing it out through the nozzle by pressure. All spraying requires supervision by some intelligent individual whose duty it is to see that the jets do not get blocked and to re-pack joints and valves⁷⁹.

(ii) A garden watering-pot may be used for sprinkling oil on small pools, drains etc.

(iii) Used cotton waste may be employed economically where there is much machinery. This may be made into swabs and

if they are not already sufficiently oily, these can be soaked in crude oil and wrung out before use. Senior White²⁸ states that they are ideal for use in small rock springs and that by their use running water is more economically oiled than by a machine. They may be weighted down with a brick or stone or pegged into the ground with small sticks. Bits of old sacking may also be used in this way. After about a week the swabs may be re-soaked and used again.

(iv) A long stick with a bundle of old sacking tied on the end may be used to distribute oil over small pools.

(v) *Balls or guddas* (pillows) may be made of tow or sacking weighted with stones and thrown into pools etc. after being soaked in heavy oil. The oil oozes out gradually and comes to the surface. A better plan is to have them attached to a float (made by sealing up an empty tin) so that they do not sink to the bottom (Fig 10). This prevents the collection of silt round




FIG 10
Oil ball and float

them which eventually makes them ineffective. They may be placed in drains or tethered along the edges of streams, irrigation channels etc. They should be opened out and dried at intervals of 7 to 10 days and re-soaked in oil. It is also well to shake them every 2 days or so without removing them from the water. A 'gudda' shaped like an ordinary pillow, 28 inches long and

The amount of oil allowed to drip suggested by Le Prince³⁶⁹ is 10 to 20 drips per minute for a water surface one foot wide

The disadvantages of oil drip-cans are that they do not give good service without proper attention and may get clogged or be washed away by floods or be stolen

(x) In Panama a *barrel pump* mounted on a rowing boat is used. Hot oil is employed because it is found that this flows better through the pipes and nozzles³³⁹

Amount of oil to be applied

It will be evident from what has been said above that the amount of oil necessary to control mosquito breeding will vary very greatly according to the local conditions obtaining i.e. climatic conditions presence or absence of vegetation kind of oil or mixture of oils employed etc. However as the question is so frequently asked in connection with the probable cost of anti larval measures it may be laid down that half an ounce of oil per square yard of water surface or 15 gallons per acre is usually an ample estimate. It is claimed for some of the proprietary preparations that very much less than this is needed. The exact amount of oil required in any particular case can only be ascertained by actual trial in the field.

Frequency of oiling

It is usually said that oiling should be repeated every 10 days but in practice it will be found more effective to oil at intervals of 7 days. This is not because such a practice is essential to control breeding but because unless it is definitely laid down in the programme that such and such breeding places are to be oiled on a specific day of each week it is inevitable that certain places will be missed. Like everything else in connection with malaria prevention the success of oiling depends on the thoroughness with which it is carried out. Oiling should not be attempted in drains or streams during heavy rain. Should a heavy downpour occur within an hour after oiling the process may have to be repeated³⁹

54 MEASURES DIRECTED AGAINST LARVÆ OF MOSQUITOES

16 inches broad made of sacking will soak up about 2 gallons of oil at first and about 1 gallon when re soaked. It will last about 3 months. Oil balls may also be dragged along drains like mops thus clearing the drain and at the same time oiling it.

Lumps of tow or sacking soaked in oil and weighted with a brick are useful in treating water under culverts.

(ii) Oil may be mixed with *sawdust* and thrown after the manner of sowing grain. The mixture must not be wet to the touch and the particles must separate cleanly when thrown on to the water. The best results are obtained when the mixture is allowed to stand for 24 hours before use (see p 47). This method has been advocated for use among vegetation. The proportions to be used are roughly 30 gallons of oil with 10 bushels of sawdust per acre³⁵⁸

(iii) A conical can divided into a lower chamber to contain sand and an upper chamber for oil has been devised by Wetmore³⁵⁹. A tube soldered to the outside opens into the oil chamber at the bottom. The apex of the oil chamber has a wick to permit the oil to escape. The can is thrown into the water and sinks air escaping from the apex followed by oil. Water is drawn through the tube into the lowest portion of the oil chamber replacing the oil. This device is said to work well in water covered with vegetation.

(iv) Standing rain pools on grass may be dealt with by means of an ordinary sweeper's broom dipped into oil and brushed round the edges.

(v) *Drip cans* may be used for the automatic oiling of streams and drains. They may be used as follows.

(a) A nail may be knocked through the bottom of the receptacle and a piece of wool wrapped round the head of the nail. By pulling on the latter the flow may be regulated.

(b) A tap may be fitted to the receptacle.

(c) A thick lamp wick may be stuffed into a hole near the bottom of the receptacle.

- (vi) It renders water unfit for drinking purposes
- (vii) It is likely to deter mosquitoes from depositing their eggs thus driving them to other uncoiled collections of water
- (viii) It is likely to be stolen by unscrupulous employees

Advantages of oiling

- (i) Oil kills the ova and pupæ of mosquitoes as well as the larvæ
- (ii) It kills culicine as well as anopheline larvæ
- (iii) It is easily obtainable everywhere
- (iv) It is easy to see whether it has been properly applied
- (v) No elaborate apparatus is required for its application
- (vi) Its use though requiring supervision does not involve the amount of supervision as does for instance that of paris green

(See *Bibliography* Nos 374-385)

4 Chemical larvicides (a) *Paris green* (copper aceto-arsenite)—Following the publication by Roubaud³⁸³ of the results of his researches as to the value of paraformaldehyde powder as a larvicide Barber and Hayne³⁸⁵ in 1921 carried out experiments with various compounds of arsenic of which paris green proved the most efficient. Since then it has been used widely in many countries and it appears to come nearer to fulfilling all the requirements of an anopheles larvicide than any other preparation at present available.

Paris green (Schweinfurt green Imperial green Emerald green Vitis green) is a micro-crystalline powder of emerald green colour practically insoluble in water but soluble in ammonia and in concentrated acids. Under United States law the name paris green is restricted to powders containing a minimum of 50 per cent arsenious oxide but in Europe the name is sometimes applied to substances containing green aniline dyes which are valueless for larvicidal purposes.

Tests of thorough oiling

Apart from the absence of living larvæ, two other tests of the efficiency of oiling may be mentioned --

- (i) The marginal vegetation along a properly sprayed drain should be burned brown for a foot on each side and after a few applications bare earth margins should become exposed
- (ii) In regularly oiled drains *Spirogyra* will disappear and be replaced by a bottom growth of a matted Cyanophylaceous alga, which as pointed out by Watson⁴ is correlated with the absence of mosquito larvæ.

Where oiling is applicable

Oil may be applied in the case of pools, drains and streams and to the edges of deep ponds lakes and rivers, and wherever it is necessary to destroy culicine as well as anopheline larvæ provided that the water is not to be used for drinking or domestic purposes. It is especially applicable to temporary collections of water which are too numerous to drain or to control breeding places while steps are being taken to deal with them by permanent measures

Disadvantages of oiling

(i) Oil will not easily penetrate a barrier of grass, and to make it thoroughly effective all vegetation and floating débris should be removed

(ii) Wind will break up an oil film and will carry the oil to one side of a sheet of water

(iii) During rainy periods the value of oiling is decreased. Showers of rain wash away the oil

(iv) Oil is heavy for transportation

(v) It may kill fish, and renders them unfit for human consumption

- (ii) It renders water unfit for drinking purposes
- (iii) It is likely to deter mosquitoes from depositing their eggs, thus driving them to other unboiled collections of water
- (iiii) It is likely to be stolen by unscrupulous employees

Advantages of oiling

- (i) Oil kills the ova and pupæ of mosquitoes as well as the larvæ
- (ii) It kills culicine as well as anopheline larvæ
- (iii) It is easily obtainable everywhere
- (iv) It is easy to see whether it has been properly applied
- (v) No elaborate apparatus is required for its application
- (vi) Its use though requiring supervision does not involve the amount of supervision as does for instance that of paris green

(See *Bibliography* Nos 324-385)

4 Chemical larvicides. (a) *Paris green* (copper aceto arsenite) —Following the publication by Roubaud³⁸⁹ of the results of his researches as to the value of paraformaldehyde powder as a larvicide Barber and Hayne³⁹⁰ in 1921 carried out experiments with various compounds of arsenic of which paris green proved the most efficient. Since then it has been used widely in many countries and it appears to come nearer to fulfilling all the requirements of an anopheles larvicide than any other preparation at present available.

Paris green (Schweinfurt green Imperial green Emerald green, Mitis green) is a micro-crystalline powder of emerald green colour practically insoluble in water but soluble in ammonia and in concentrated acids. Under United States law the name paris green is restricted to powders containing a minimum of 50 per cent arsenious oxide but in Europe the name is sometimes applied to substances containing green aniline dyes which are valueless for larvicidal purposes.

Method of action of paris green

The particles are eaten by the anopheline larvæ, which feed on the surface of the water, and the chemical acts as a direct poison. It is therefore important that the particles of paris green should remain floating on the surface for as long a period as possible. This is largely determined by the surface tension of the water. The more vegetation that is present in the water the greater is the surface tension, the longer do the particles float, and hence the greater is the efficiency of the paris green as an anopheles larvicide. Experiments are being carried out by various firms with the object of producing a form of the chemical which will remain on the surface for the longest possible time.

Paris green has no effect on the pupæ of mosquitoes, because these do not feed, nor, when applied in the usual way, does it affect the larvæ of culicines because these are not surface feeders. The very young (first day) larvæ of anophelines are also usually unaffected, presumably because the particles of paris green are too large for them to eat. It is possible that this latter difficulty may be overcome by the production of an extremely fine form of the chemical.

Requirements of a good paris green

- (i) It should be of a deep green colour.
- (ii) It should contain over 50 per cent arsenious oxide, a good quality averages 55 per cent.
- (iii) Its solubility in water should not exceed 3 per cent.
- (iv) It should be capable of being passed through bolting cloth of at least 200 preferably 300 mesh.
- (v) Each batch should pass a practical field test before acceptance for it has been found by experience that samples apparently identical vary greatly in toxicity to larvæ.

Choice of diluent

For use paris green is mixed in varying proportions with a dry powder of some description which acts as a vehicle for its

distribution * A very large number of substances have been used for this purpose amongst which the following may be mentioned —

| | | |
|---------------------------------|------------------|-------------|
| Pond dust | Sulphur | Cement dust |
| Slaked lime | Spoiled flour | † Trass |
| Powdered soapstone or softstone | Ped earth (Java) | Sawdust |
| Ashes | French chalk | Fine sand |
| Cork dust | Coir dust | China clay |

The choice of diluent will depend on the conditions under which the larvicide is to be used. Other things being equal the cheapest suitable diluent locally available will naturally be selected.

(1) For dusting an extensive area such as a swamp or large sheet of water a light vehicle which will carry the paris green with it for a long distance in the wind is required. Soapstone or 'soft stone' (which consists chiefly of silicates of lime and magnesium) and slaked lime have been found suitable for this purpose in India. The former may be obtained in the form of a powder (for sources and prices see Appendix). Slaked lime is prepared from quick lime which can usually be obtained locally. The lime should be of good quality and the slaking done with care, a small quantity of water being added at a time. The process of slaking should never be carried to completion. During the process of screening (see below) which should be done shortly before use the unslaked portions will be removed and should be put back with the unslaked lime for future use. In moist climates simple exposure to the air is sufficient.

Road dust varies in composition in every locality and very frequently contains a considerable quantity of sand which makes it unsuitable for dusting large areas.

* Kerosene oil or a mixture of kerosene and waste motor oil has also been used as a vehicle for distributing paris green in order to obviate the necessity for storing dust during rainy seasons.

† Trass is a volcanic substance largely composed of pumaceous dust and is worked for mortar.

(u) In the case of small streams, ditches, irrigation channels etc., a heavier diluent is required, as, if there is any considerable wind a light vehicle such as powdered soapstone or slaked lime is likely to be carried beyond the breeding place. This is a point of very great practical importance indeed, with a very light diluent if the wind is at all high, the attempt to dust such breeding places may be absolutely ineffective. The local road dust will frequently be found suitable for this purpose, or fine sand alone may be used.

Swellengrebel and his co-workers³⁰³ in Holland first used ashes as a diluent, with 1 per cent of paris green. They found that some of the larvæ survived after the first day, and concluded that this was because some of them swallowed large quantities of ashes besides, or even instead of, paris green. They therefore sought for a heavier dust, which would sink, leaving the paris green floating on the surface of the water and finally selected trass as being more suitable.

Williams⁴³³ and other American workers on the other hand, who wish to dust very extensive areas by means of electric blowers and aeroplanes and who use paris green in dilutions of from 5 to 33 per cent prefer a diluent which will float for the longest possible time such as soapstone or slaked lime.

Recent experiments by Dolloff⁴¹² have shown that mixtures of hydrated lime with the stearates of calcium and aluminium as diluents for paris green prolong the period from the time of dusting to the re-appearance of anopheline larvæ from an average of 3 days to an average of 5 to 6 days, in spite of intervening showers of rain.

Screening

Whichever diluent is employed, it is usually necessary to pass it first through a screen or sieve, in order to get rid of the larger particles. For this purpose it is customary to use an apparatus consisting of a double cylinder of wire gauze, the inner one through which the dust first passes) being of coarse mesh (about 2 to the linear inch) the outer of a finer mesh (about 30 to the

linear inch) The cylinder is mounted within a wooden box on a stand and is operated by turning a handle* (Fig 11)




FIG 11

Screener for dust to be used as diluent for paraffin green (after Missirol)

A simpler type consists of a rectangular wooden tray about 2 feet long by 1½ feet broad with a fine wire gauze bottom on the

* The following dimensions for this screener are given by Missirol —

| | | |
|--------|-------------------------------|------------|
| Box | Length | 58 cm |
| | breadth | 31 |
| | height | 40 |
| Hopper | radius | 21 |
| | length | 57 |
| | slope | 4 |
| Tresle | diameter of internal cylinder | 18 cm |
| | mesh of gauze of | 7 to 1 cm |
| | diameter of external cylinder | 24 cm |
| | mesh of gauze of | 15 to 1 cm |
| Tresle | length | 60 cm |
| | breadth | 33 |
| | height | 40 |
| Bucket | diameter of mouth | 15 cm |
| | base | 37 " |
| | height | 47 cm |

top of which fits a second tray with a bottom of coarse mesh wire. The screener is suspended from the roof by cords attached to its four corners and is agitated by hand, after the manner of sifting grain. The larger particles are removed by the upper tray. This apparatus which may be made locally at very small cost (about Rs 5 in India) has been found suitable for screening road dust. It has the great advantage that there are no complicated parts to get out of order⁴⁰¹

Mixing

This must be thorough, so that the resulting mixture is uniform. A suitable apparatus may be constructed from a rectangular box of galvanized iron suspended on two pillars from diagonally opposite corners and turned by a handle (Fig 12). A




FIG 12

Mixer for use with paris green (after Missirol⁴²¹)



PLATE II



A Haversack for the application of paris green by hand



B Rotary Blower (Peerless Dust Gun) for Paris green
dusting

convenient size is 33×48 cm (about 13×19 inches) Or, an oil drum or small barrel with an iron rod running diagonally through it may be used About 200 turns are necessary for satisfactory mixing

Methods of application

(i) *By hand*—The mixture may be thrown by hand after the manner of sowing grain from a bag resting on the hip and suspended from the opposite shoulder This method has the advantage of dispensing with mechanical blowers, which so frequently get out of order in the hands of the average coolie, and is probably the best way of applying paris green to irrigation channels, drains, etc., in the tropics (Plate II, A)

(ii) *Hand bellows* can be made locally in any bazaar for about Rs 10, and have been recommended for treating small streams irrigation channels ditches, etc Their use is, however, fatiguing to the operator, and they are soon broken by the average coolie, so that in the tropics distribution by hand is usually preferable

(iii) *Knapsack sprayers* are used extensively in Italy, where the local peasant is accustomed to employ the same apparatus for spraying his vines with sulphur etc For use with road dust in India they have been generally found unsuitable owing to the frequency with which jamming occurs They may be used for lighter dusts, such as slaked lime though even then they are apt to get clogged at times when this happens it can often be remedied by inverting the machine and shaking it vigorously

(iv) *Rotary blowers*—There are many of these on the market, and though more expensive than the knapsack sprayers they are much more effective in dealing with large areas The container for the dust and that for the mechanism should be completely separated It is also an advantage for the dust container to be placed behind the operator (as in the case of the 'Peerless Dust Gun') and the gear box and fan in front, the blower is then better balanced, and is less fatiguing to operate than if the whole apparatus is carried in front of the body (Plate II, B) The gears and fanshaft should preferably run on ball bearings.

(v) *Electric blowers*—A dusting unit consisting of a power generator, an electric blower and a small dust hopper has been used with success in America in dusting large sheets of water from a boat. With this machine using a 15 per cent mixture of paris green with slaked lime a lethal path 525 feet wide has been obtained with a moderate breeze (7 or 8 miles per hour).

(vi) *Dusting from aeroplanes*—Insecticidal dusts were first applied to crops from aircraft in the U. S. A. in 1921 and King and Bradley⁴⁹³ employed this method for dusting with paris green in experiments conducted in 1922-24. It has been found an effective means of controlling anopheline breeding in large swamps both in the United States and in Italy. A 'venturi tube' some 2½ feet long by 8 inches square at the mouth, 7½ inches in the centre and 12 inches at the outlet in the rear, is fixed beneath the fuselage of the plane. The dust is fed by gravity from a V-shaped hopper, 30 inches high, into the constriction of the venturi tube, the lower aperture of the hopper being controlled by a sliding door which can be operated by the pilot. When the aeroplane is travelling at high speed (about 65 miles per hour) a high velocity of air current is created at the narrowest part of the venturi tube and a partial vacuum at its outlet. The dust is well broken up as it enters the partial vacuum and is blown out of the tube in an even cloud. By this method a lethal path some 200 yards wide is obtained. Cook and Williams⁴⁹⁹ state that a properly equipped plane flying 4 hours a day for 5 days a week could deal effectively with an area of 20 square miles.

*Methods of determining toxicity of paris green, suitability
of diluent, etc.*

In carrying out preliminary tests to estimate the toxicity of a batch of paris green, the distribution and concentration of the mixture and the suitability of the diluent, the following methods may be used—

(1) Various points in the area to be dusted are selected and a specified number of dips are made in each breeding place with a ladle or dish, the average number of larvæ obtained per dip being

PARIS GREEN

counted. The places where the dipping was done are marked by stakes. About 3 hours after dusting the process of dipping is repeated and the results compared.

It should be borne in mind that the length of time in which larvae are killed depends not only on the degree of concentration of the paris green but also on the temperature of the water. In warm weather the larvae will feed much more actively and will thus be killed more quickly whilst when the temperature is very low they may not be feeding at all in which case any brand of paris green will be ineffective. It is therefore important when comparing the efficiency of different samples of the chemical to carry out the tests under similar climatic conditions.

(ii) Dishes containing known numbers of living larvae are placed at various points over the area to be dusted and the numbers killed or remaining alive after dusting noted. This method though useful does not give as conclusive evidence of the mortality rate as does (i).

(iii) Glass microscopic slides with a one inch square ruled on each are placed at various points. It is convenient to place these across the corners of the dishes used in (ii). After dusting the slides are collected and the number of paris green particles per square inch on each is counted under the two-thirds objective of a microscope. For effective dusting there should be at least 12 particles per square inch on each slide. This is a very useful method for estimating the distribution and concentration of the paris green.

Strength of paris green to be used

For dusting narrow streams, drains, small pools etc. a one per cent dilution of paris green is sufficient. For use over larger areas from 2½ to 5 per cent or even more will be found more effective. For the electric blower a dilution of 15 per cent is recommended whilst for distribution from aircraft Cook and

Williams⁴⁰⁹ have found that 33 per cent is the most suitable for all conditions of wind *

Quantity of paris green to be used

Hackett⁴¹⁷ uses 0.1 c.c. (0.125 gm., 1.9 grains) to the square metre of surface. Thus a litre (1,250 gm.) should control breeding over 10,000 square metres of surface, or along 10 kilometres of water edge. This works out at approximately 1 lb. of paris green per acre, which is the amount usually recommended †

If carefully applied in high dilution with a hand blower it is possible to control breeding with less than half this amount but in practice it is found that such extreme care in application is not economical from a labour point of view.

The presence of scums or dense vegetation on the water surface necessitates the use of somewhat larger quantities of paris green per unit area.

Frequency of dusting

As in the case of oiling this will depend very largely on local conditions, such as intensity of breeding and variations in climate. As a general rule where breeding is profuse paris green should be applied once a week. Sometimes, where the period of development of the larvæ is shortened owing to specially favourable climatic conditions, it will be found necessary to reduce this interval to 5 or 6 days for a while. Again, as the weather becomes colder and the period of larval development becomes lengthened it may only be necessary to dust at intervals of 10 days or so. Even in this case, however, a seven day interval is to be

* Some workers measure the dilution by weight, others by volume. As some diluents are much lighter than others and may be less than half as heavy as the paris green the two methods of calculating the dilution may give widely different results. It is considered advisable always to make the dilution by volume. In any case, in recording results, it should invariably be stated which method has been used in calculating the dilution.

† Barber and Hayne³⁹⁸ recommended 10 c.c. (12 grams, 0.43 oz., 0.6 cubic inch, or 2 level teaspoonfuls) to 90 square metres (1,000 square feet).

recommended, as if it is laid down that each breeding place is to be dusted regularly on a certain fixed day each week there is much less likelihood that any particular breeding place will be overlooked

*Summary of recommendations as to the methods of applying
paris green in the tropics*

From the results of experiments carried out by the Malaria Survey of India up to the present date the following procedure is recommended —

- (i) For dusting large areas a 2 to 5 per cent dilution of paris green by volume in slaked lime or powdered soap-stone (or soft stone') applied by means of a rotary blower $\frac{1}{2}$ to 1 lb of paris green to the acre
- (ii) For treating narrow irrigation channels ditches, etc a 1 per cent dilution of paris green in road dust, or fine sand applied by hand from a haversack

Cost of application

This naturally varies considerably according to the local conditions. As a rough guide it may be mentioned that the cost of applying a 5 per cent paris green mixture with slaked lime at Karnal Punjab by means of a rotary blower, using $\frac{1}{2}$ lb of paris green per acre works out at about Rs 12 per acre. This includes the labour for slaking the lime but not the wages of the coolie distributing the mixture. No allowance is made here for the cost of the flower.

Effects of paris green on other forms of life, etc

Workers with paris green are practically unanimous in stating that in the quantities in which it is used as a larvicide this chemical is harmless to man and domestic animals who may drink the water so treated or to fish or any other forms of aquatic life except mosquito larvae.

Hackett⁴¹⁷ quotes Davis³⁹⁸ as mentioning the poisoning of some ducks and chickens with paris green on one occasion in Brazil. In a private communication Dr Davis informs me that on this occasion some dead chickens and ducks were found on the day following the application of paris green to some pools, but that subsequent inquiries elicited the fact that chickens were also dying in other parts of the town, in which paris green had not been applied, presumably from some epidemic such as fowl cholera. It is unfortunate that this report should have found its way into the literature. In point of fact, birds are particularly resistant to the effects of arsenic.

There is on record an observation by Herrmann Kolosow and Lipin⁴¹⁸ that one out of three rabbits to which drinking water which had been sprayed with paris green was given over a prolonged period developed a paralysis of the hind foot which suggested the possibility that paris green dusting was not so harmless as had been supposed. The investigations of other workers however have tended to show that under natural conditions there is no accumulation of arsenic in solution in water to which paris green has been applied over long periods. Hackett⁴¹⁷ using *Penicillium brevicaulis* a mould which gives off a characteristic garlic odour in the presence of a medium containing one part in 100 000 of arsenic, found that samples taken daily from shallow pools of standing water gave strongly positive tests 24 hours after treatment with paris green, but were always negative after 48 hours. He concluded that arsenic does not accumulate in such collections of water, but is rapidly eliminated, 'perhaps through volatilization by the microflora of the water'.

The Schuurmans⁴⁴⁶ using a 2 per cent dilution of paris green on fish ponds made various analyses of the water treated, and of fish caught in the water. They found a minute percentage of arsenic in each case, but quantities as large were found at times in the case of the fish and water in untreated fish ponds. They calculated that one would have to drink 27 gallons of the water, or consume 50 pounds of fish in order to obtain a dose of 5 mg of arsenic.

Paris green has also no ill effect on rice or other crops. Nicholls⁴³¹ grew rice and certain other plants in tubs in which 2 gm of 2 per cent paris green in coir dust was sprayed over each square yard of water surface without any deleterious effect.

As regards the effects on those working with paris green occasional complaints of headache and diarrhoea have been received in the case of men distributing it by hand. There have also been cases of dermatitis when mixing has been done by hand. Otherwise there appears to be no danger of ill effects and the only precaution necessary is to advise workers to take ordinary care to keep to windward of the dust cloud to change their outer clothing after the day's work and to wash their hands before eating.

In this connection there is an advantage in using hydrated lime as a diluent. This is somewhat irritating to the throat and nose and workers with it will of their own volition keep to the windward of the cloud as much as possible without any instructions being given to that effect.

The question of the effect of paris green on boilers has been raised in connection with the treatment of mill ponds. It is difficult to imagine that such minute quantities of the chemical as are used in antilarval work could have any deleterious results. In America it has been satisfactorily shown that no ill effects need be feared and the owners of one mill in that country offered to put half a pound of undiluted paris green into one of their boilers as a demonstration of its harmlessness⁴⁴³.

Advantages of paris green

- (i) Its low cost
- (ii) Its high toxicity for anopheline larvae
- (iii) Its portability
- (iv) Its ease of distribution by wind
- (v) There is no need to remove vegetation however dense

(vi) There is no ill effect on domestic animals fish or other natural enemies of larvæ or on crops such as rice

(vii) The water treated is not rendered unfit for domestic purposes

(viii) Being useless for other purposes it is not likely to be stolen

(ix) It is quickly applied

(x) It does not prevent mosquitoes from depositing their eggs the water treated thus acting as a trap

(xi) It is equally effective in fresh or brackish waters

Disadvantages of paris green

(i) It has no effect on the eggs or pupæ of mosquitoes and sometimes does not kill the very young larvæ of anophelines

(ii) It has no effect on culicine larvæ when applied in the usual way This is a great disadvantage of paris green for although as far as the control of malaria is concerned it is only necessary to destroy anophelines yet the great majority of the general public believe that all mosquitoes carry malaria and if the culicines remain as numerous as ever the campaign is likely to become discredited Moreover there are many places where the 'mosquito nuisance' (usually mainly due to the bites of culicines) is so great that it is desirable to destroy these also if only for the personal comfort of the inhabitants, whilst in dealing with diseases transmitted by culicines the use of paris green is of course valueless

In this connection it may be mentioned that Griffiths⁴⁴ has reported some success against culicine larvæ by mixing paris green with moist sand so that it is carried to the bottom of the water at once

(iii) It is impossible for inspectors to see whether it has been applied properly or not (except of course by dipping for larvæ) Gater⁶⁴ has made experiments with various commercial dyes incorporated in the paris green mixture with the object of giving

visible traces of dusting but the results on the whole have been disappointing

(1r) It requires special apparatus for distribution and for screening the diluent and for mixing

(1r) Its use requires constant supervision

(11) It is stated by some authors that it is ineffective in showery weather Nicholls⁴³, who used a 1 per cent mixture with coar dust stated that it was found almost impossible to destroy larvæ during rain On the other hand Swellengrebel and his co-workers³¹² using 1 per cent paris green with ash, obtained counts of 32 granules per field within the first 24 hours 'a shower of rain or even three of them making no difference' Dolloff⁴¹², using hydrated lime mixed with stearates as a diluent, also found paris green effective in spite of showers It is difficult to reconcile these statements, and further investigation on the subject is necessary * A similar difference of opinion exists in relation to the effect of the presence of dew on the vegetation, some authors stating that dusting is less effective until the sun has dried it up whilst others (e.g. Williams⁴) state that this makes no difference to the lethal effect It is possible that the choice of diluent may have some influence in both these cases

(See Bibliography Nos 395-455)

(b) *Paraformaldehyde (trioxymethylene)* This substance was recommended by Roubaud³⁸¹ as a larvicide in 1920 The dry powder used alone or mixed with some vehicle such as flour or powdered chalk was spread in the form of a dust cloud on the surface of the water where it was ingested by the larva More recently the same author has described a larvicide having the trade name of stoxal the active principle of which is paraformaldehyde³¹¹

✓ B—For prices and sources of paris green and the various apparatus used in its application see Appendix

* Barber⁴⁰⁸ notes that the action of paris green is most effective immediately after a shower of rain when scums which tend to interfere with the surface spread of the dust particles, are beaten down

During the last few years several workers have tested this substance among whom may be mentioned Barber and Komp³⁸⁶ Mertens³⁸⁸ Ginsburg⁶⁸ the Schuurmans³⁹⁰ Swellengrebel and de Rook³⁷⁷ All these have reported unfavourably on stoxal as compared with paris green * and its chief interest lies in the fact that the work of Roubaud with paraformaldehyde and other powders directly inspired the investigations that led to the discovery of the properties of paris green as a larvicide

(See *Bibliography* Nos 386-394)

(c) *Cresol*—In reading the accounts of experiments by various workers with this substance as a larvicide one is at once struck by the discordant views expressed as to the strength in which it must be used The reasons for this are that different samples vary in composition that dilutions of saponified cresol deteriorate in toxicity on standing and that the composition of the diluting water also has an effect Harold¹⁰⁶ noted that an acceleration of the larvicidal action occurs with higher temperatures and that the presence of organic silt leads to a reduction of larvicidal power He also showed that the resistance of larvæ to cresol could be enhanced by placing them in non lethal dilutions overnight and transferring them to lethal ones next day a point to be taken into consideration when collections of water are intermittently treated Stewart¹¹⁶ notes that the germicidal coefficient of a coal tar disinfectant is no sure guide to its larvicidal power and that the hydrocarbon oils probably play a much larger part than might be thought in comparison with the phenoloids

Mayne and Jackson¹¹⁵ stated that a final dilution of 1 part of cresol in 1 000 000 was effective for *Culex* larvæ and 1 in 100 000 for *Anopheles* but other observers have found that much larger amounts are necessary † Delmege¹⁰³ stated that a dilution of 1 in 100 000 was effective in standing water and 1 in 1 000 in

* It should be mentioned however that Banvallet³ has reported good results with stoxal

† Hehr²⁰ states that 1 in 10 000 000 is sufficient but this is evidently due to a misinterpretation of the article by Mayne and Jackson

slow flowing streams, whilst spraying with a dilution of 1 in 500 delayed the growth of vegetation Strickland and Roy¹¹³ found that a dilution of 1 in 50 000 was the greatest to be of any practical utility, Senior White²⁸ recommends 1 in 40 000 and Marshall¹¹⁴ 1 in 28 000 whilst Klügler¹⁰⁸ found a dilution of 1 in 10 000 to be necessary Harold found that of two different samples of cresol one killed *Anopheles bifurcatus* larvæ in a dilution of 1 in 10 000 whilst the other had no effect in this dilution

Klügler¹⁰⁹ found this larvicide very useful in Palestine in the case of gravelly marshes along the sides of lakes and river banks where the larvæ are hidden under the pebbles A dilution of 1 part cresol in 100 parts of water sprayed at the upper end of the marsh so that the cresol filters slowly through with the stream either kills the larvæ or drives them out into the stream where they are destroyed

Marshall¹¹⁴ reckoning that cresol cost about 4 times as much as kerosene oil and that the dilution necessary for the former was 1 in 28 000 calculated that cresol was the more economical method for water with a depth of less than 5 inches For deeper waters the cost of cresol is prohibitive

In the case of mosquito breeding in the water used for keeping cement concrete wet whilst it is setting cresol is a most useful larvicide as it has no ill effect on the concrete In Bombay *A. stephensi* breeds freely in such water and the use of cresol has been recommended to deal with this^{2,6} In practice it is found convenient to add just sufficient cresol to make the water milky This is a much stronger dilution than is necessary to kill the larvæ but it makes it easy for the inspecting staff to see at a glance whether the larvicide has been applied

A disadvantage of the use of cresol in natural water is that it kills fish Senior White²⁸ notes that fish killed by cresol are violently poisonous when eaten by man

(See Bibliography Nos 104-116)

(d) *Commercial Cyllin* was used with effect by James⁷³ in Ceylon. One teaspoonful in a gallon of water kills larvæ and pupæ but in practice it was found best to use sufficient to make the water milky after it had been well stirred with a stick.

(e) *Sanitas Olo!* a proprietary emulsion which appears to contain a large proportion of phenols and allied compounds was favourably reported on by Ross and Edie¹⁰¹ who found that it was a powerful larvicide effective in dilutions up to 1 in 10 000 at which strength it was non poisonous. Mayne and Jackson¹¹⁵ however found it inferior to cresol.

(f) *Panama Canal Larvicide*²⁵⁷—This is made as follows—150 gallons of carbolic acid are heated in a steel tank to 212°F. Then 200 lbs. of powdered resin are poured in and stirred until completely dissolved. 30 lbs. of caustic soda dissolved in 6 gallons of water are then added and the temperature kept at 212°F. for about 5 minutes. A sample is withdrawn and poured into water if complete and rapid emulsion occurs the larvicide is now ready. It is claimed for this preparation that—

- (i) It is highly toxic to anopheline larvæ a 1 in 1 000 emulsion killing them in 3 to 10 minutes
- (ii) It is concentrated and therefore only a small quantity need be carried
- (iii) Its toxicity is uniform
- (iv) Its composition is simple
- (v) Its toxicity to higher animals is low
- (vi) Its cost is low
- (vii) There is no danger from fire
- (viii) It is highly toxic to protozoa and algæ

Its disadvantages are that in brackish water it is inert and does not emulsify that it kills fish that it deteriorates on exposure and that it rapidly loses its toxicity after mixing with water containing algæ and other organic matter³⁴

This larvicide was used largely during the construction of the Panama canal. Le Prince³⁰⁹ notes that it was very effective when added to crude oil in the dilution of 5 to 10 per cent. In the

Canal Zone at present, however, only crude petroleum (fuel oil) is used¹⁰

(g) *Crude Carbolic Acid*—The use of this substance as a larvicide was advocated by Wise and Minett²⁹¹ in British Guiana, for the treatment of small depressions holding temporary water. It was used in a final dilution of 1 in 16 000. The crude acid, containing cresol, resolic acid and various oily and tarry substances, was found much more effective than the purified product. It was not, however, found suitable for use with a sprayer because some of the constituents are not soluble.

(h) *Creosote* in the proportion of 1 in 4 million parts of water, is stated to be an efficient larvicide and has been used in Italy²⁹², and in the U.S.A.³¹

(i) *Commercial Borax* in the strength of 0.3 ozs. to 10 pints of water is recommended by Matheson and Hinman²⁹³ as a cheap and practical larvicide for use in cisterns, rain water barrels and any container for water used for other purposes than drinking and in pools where the plant and animal life is of little importance.

(j) *Lime*—Borel²⁹⁴ found that mosquitoes did not deposit their eggs in water receptacles coated inside and out with milk of lime though the pH value and calcium carbonate content of the water was not inhibitory to the development of eggs nor was the microscopical algae necessary for the development of the larvae reduced. He concluded that the factor which repelled the adult female was solely the white colour of the containers. These should be lined at least once a month in order to keep them quite white.

(k) *Soap* This is an effective larvicide its great disadvantage being its high cost. Bilsour^{*} suggested that marine soap which is manufactured in large quantities in Marseilles might prove of practical utility. Strickland²⁹⁵ has experimented with this but has found that in laboratory experiments the treatment of 4 square feet of water 1 foot deep would be 18 times more costly than by kerosene whilst the treatment of a pool with 48 square feet of

^{*} *Trans. Roy. Soc. Trop. Med. and Hyg.* XVIII, 4, pp. 153-156 (1924)

surface and only one inch deep would still be 4 times as costly with soap as with kerosene. He found also that in field experiments the soap was less effective than in the laboratory.

There is one way, however, in which the larvicidal properties of soap may be used in the tropics viz, 'dhobies' may be encouraged to wash clothes at the head of streams harbouring anopheline larvæ.

(l) '*Parisian*' is a proprietary preparation which has been put forward as an efficient and cheap larvicide. The Schuurmans³³² however, reported unfavourably on this substance, finding it ineffective even in quantities 6 times as great as those prescribed.

(m) '*Isotex*' is another proprietary larvicide, recommended for use in swamps polluted pools, etc. It is a dark brown fluid, forming a thick film on the surface of water, adhering to any surface even glass and staining clothes. Feegrade³⁰¹ found that in natural breeding places of about 72 square feet of water surface 1 to 2 feet deep, with much aquatic vegetation at the margins 30 c.c. of the larvicide destroyed all aquatic life in 75 minutes. He concluded that it is a powerful larvicide, but unsuitable for general use because it is fatal to fish.

(n) '*Empranin*' is a proprietary larvicide, which has been used with success in Bengal. It is claimed that 6 ounces of concentrated Empranin No. 2 (special quality) will suffice for a water surface of 2 000 square feet. Clyde²²² used this larvicide in anti-malarial work during the construction of the Sarda Canal Headworks United Provinces, India, but its use was discontinued on account of its greater expense, as compared with that of fuel oil.

(o) '*Emulsified Necrosene*' is a proprietary preparation containing approximately 90 per cent of oils, including creosote, turpentine and kerosene, and 5 to 8 per cent of soap. It is claimed for this preparation that its lethal effect on larvæ will persist for four to six months following a single application in a final dilution of from 1/15 000 to 1/25 000.

(p) *Hycol* a preparation derived from coal tar and closely resembling *creolin*, has been used with success in the Belgian Congo and in South Africa. *Sal hycol* is another product of the same firm designed for use in brackish or hard waters.

A very large number of other chemicals have been used for larvicidal purposes, amongst which may be mentioned various fluorine compounds potassium permanganate, mercurous and mercuric chloride, lithium chloride oxalic acid, salicylic acid, potassium cyanide naphthalene arsenic trichloride, copper, iron and sodium arsenate saprol waste sulphite liquor from paper mills waste products of sugar cane crushing etc. Mention may also be made of experiments by Williamson²⁹⁰ as to the effects of gases and vapours on mosquito larvæ in wells (formaldehyde, sulphur dioxide chloropicrin chlorine, steam etc.)

(See *Bibliography*, Nos 249-291.)

5. Sulphate of Copper.—This has been used for many years for inhibiting the growth of algae in reservoirs. For this purpose it is usually employed in a final dilution of 0.1 to 0.25 parts per million of water though this is not sufficient to destroy all forms of algae. Senior White⁸ has shown that certain species will grow even in a dilution of 1 in 10 000 of copper sulphate. As an anti-mosquito measure this chemical is usually used in a final dilution of approximately 1 in 500 000 with the object of destroying algae and thus diminishing the food supply of the larvæ. This amount was found satisfactory by Kligler⁹⁹ in Palestine.

The method of application is to place the required quantity of ordinary commercial crystals of 'blue vitriol' in a gunny bag or wire basket attached to a rope and draw it backwards and forwards in the water at the stern of a rowing boat. Owing to the poisonous nature of the chemical its use cannot be left to the ordinary cooche. It needs intelligent and careful supervision. Small fish may be killed by copper sulphate in dilutions of less than 1 part per million of water¹⁰⁰.

A disadvantage which is sometimes met with is the odour produced by the decaying algae after they have been killed, which

- (ii) For paddy fields wells small ponds etc

*Chela**Haplochilus**Polyacanthus*

- (iii) For salt or brackish water ponds etc

*Therapon**Polyacanthus*

Most observers in India appear to agree that the various species of *Haplochilus* are the most useful destroyers of mosquito larvæ. *H. panchax* (Piku) and *Anabas scandens* (Khajura the climbing perch) have been used in the wells in Bombay for many years. The former was the species tried by Aitken in some of the Bombay fountains and called by him *scooties*. Sewell and Choudhuri¹⁴ stated that none of the species of *Chela* are surface feeders and that *Rasbora daniconius* is a purely vegetable feeder but Wilson¹⁷ says that both these statements are incorrect.

As regards the area of surface that can be kept free from larvæ by fish Sewell and Choudhuri¹⁴ state that in the case of *Haplochilus* three specimens were sufficient in captivity to prevent larvæ developing in a small aquarium having a surface area of 14 by 21 feet.

Requirements for larvicidal fish

(i) They must be small so that they can get about in shallow water among weeds etc.

(ii) They must be hardy and flourish in both deep and shallow waters.

(iii) They must be able to breed freely in confined water areas.

(iv) They must be able to stand transport and handling.

(v) They must be difficult to catch and able to escape their natural enemies including man.

(vi) They must be absolutely worthless and insignificant as food.

(vii) They must be top feeders and carnivorous.

Limitations of fish in the control of mosquito breeding

There are certain disadvantages in the use of fish as a control measure —

- (i) Fish are only effective if present in sufficient numbers
- (ii) They are only completely effective in the absence of all weed and floating débris
- (iii) Small boys can be relied upon to catch them if they get the opportunity
- (iv) Over zealous persons are apt to introduce other and larger species which may prey on them
- (v) If there is not a sufficient food supply for them e.g., in wells they will eat their own young so that breeding places must be periodically re stocked
- (vi) Constant inspection is needed to see that the fish are flourishing and are in sufficient numbers and that the water is free from horizontal vegetation and floatage
- (vii) In order to carry out these inspections and to keep up a sufficient stock of fish for the various breeding places a special staff is necessary

For these reasons the use of fish as a measure of malaria control though useful in certain conditions e.g. in places where for some reason it is impossible to employ any other measure, cannot be said to hold a position of very great importance in India

(See *Bibliography* Nos 196-217)

8 Introduction of Deterrent Aquatic Plants.—(a) Those which act by covering the water so thickly as to prevent the respiration of larvæ e.g. the *Lemnaceæ* Adie¹¹⁷ in India recommended the use of *Lemna minor* for this purpose. Certain species of *Azolla* a duckweed with a fern like leaf have been cited as a means of preventing mosquito breeding by MacGregor¹⁴¹, Mühlens¹⁴⁹ França¹³⁵ and others though experiments by Smith¹⁰² in America suggest that this plant is of little value Bentley¹²²

stated that he had found both *Lemna* and *Azolla* of little value, but that *Wolffia arhiza*, which occurs in the form of small bright green round grains, without stem, roots or leaves, appears to be a useful preventive. *Anacharis alsinastrum*, or American duckweed, and *Trapa bispinosa* or water nut plant (*singhara*), have also been said to inhibit larval development.

The use of the water hyacinth (*Piaropsis crassipes*) has been advocated by Viosca¹⁵³ in America. Barber and Hayne¹²¹, however, came to the conclusion that it was of little value, and in any case on account of the disastrous effects of this plant in choking waterways it may be dismissed as a practical method of mosquito control.

(b) Those which act by entrapping larvæ, e.g., the aquatic carnivorous plants such as the *Utricularia*, or bladderworts. The bladders of these plants have a valve like door, through which mosquito larvæ and other aquatic animals enter when searching for food or seeking to escape from their natural enemies. It is said also that a member of the sundew family (*Aldrovanda vesiculosa*) which grows submerged in water, captures mosquito larvæ²⁴.

(c) Those which are actively poisonous to larvæ. In 1919, Caballero¹³⁰ reported that the presence of *Chara fatida* in collections of water prevented the breeding of mosquitoes. Subsequent experiments by various workers with different species of *Chara* have led to conflicting results, the majority (e.g., MacGregor¹⁴², Hacker³⁵⁰, Fisher³³⁹, Barber¹²⁰, Blow¹²³, Buxton⁶⁰ and Hamlyn Harris⁶³) being unfavourable. Matheson and Hinman¹⁴⁵, however, found that pools in which *Chara fragilis* was growing were free from mosquito larvæ, and are of opinion that its introduction in temporary and permanent pools, slow flowing streams, marshy areas etc., might prove of practical value in anti mosquito work. As to the method by which *Chara* produces its inhibitory effect, Caballero¹³⁰ thought that it was due to asphyxiation by an oily substance exuding from the plant on to the surface of the water, whilst Matheson and Hinman¹⁴⁶ suggest that it may be due to the presence of large amounts of oxygen given off by the

plant and oxidizing organic substance in solution thereby making this type of food unavailable for larvæ. They also found that dried *Chara* showed a marked lethal action, indicating the presence of some toxic substance.

Another member of the *Characeæ* *Antella phauloteles*, was stated by Buhol¹²⁹ in Queensland to prevent mosquito breeding, but Hamlyn Harris¹³⁷ found 3 species of mosquitoes breeding freely in water in which this plant was growing. The latter observer found that a species of alga *Cladophora holmii* inhibits larval growth and suggests that this may be due to the production of a narcotic poisonous gas.

Boyd⁶ states that in Brazil he never found mosquito larva in marshy areas in which the swamp lily or wild ginger (*Hedychium coronarium*) grew. Hildebrand¹³⁸ notes that in the U.S.A. anopheline larvæ are rarely found in water where smart weed (*Polygonum* sp.) is growing in abundance whilst Purdy⁷³ attributed the absence of anopheline larvæ in a rice-field in California to the presence of a blue green alga of the genus *Tolypothrix*.

With the possible exception of the duckweeds which might be useful in the case of disused wells etc. it cannot be said that at present there appears to be any indication that the introduction of deterrent plants will prove of much value as a practical measure of mosquito control.*

(See Bibliography Nos 117-153)

9. Biological control of mosquito breeding.—In 1911

Watson¹³⁹ drew attention to the fact that in certain localities Nature has carried out in the most successful way a method of so altering the composition of the waters of the hills that the carriers of malaria no longer find it suitable for their existence and malaria practically does not exist. He suggested that researches into this question might reveal some way of

* The question of planting shade-giving plants and trees as an anti-larval measure in the case of species of anophelines which prefer sunlit breeding places has been discussed on p. 36.

opened at intervals, thus altering the level of the water and flushing out the lake

Where permanent measures cannot be carried out, the best way of dealing with such breeding places is by dusting them with paris green by means of a rotary blower, or from an aeroplane, in the case of very extensive swamps

2. Salt-water Swamps.—As far as India is concerned the problem in this case is limited to the coasts of Burma, Bengal and the Andamans, where *Anopheles ludlowi* occurs. This species breeds in pools which are reached by the spring tides, but are not washed out daily by the neap tides. It is in places where artificial bunds have been constructed that most breeding occurs. The ideal solution is to reclaim the swamp by dredging or hand filling and where a comparatively small swamp exists in the vicinity of valuable land, e g, near a town, this may be a paying proposition. To reclaim an extensive swamp with the sole object of protecting a small village is usually out of the question for financial reasons, and here the only alternative is often to remove the village to another site.

Where there is sufficient tide, the installation of automatic sluice gates (such as the Calco gate) in the bunds may be effective¹⁷⁷

In Panama it has been found that the larvæ and pupæ of the local malaria carrying anophelines will cease to develop if strongly saline water is introduced into their breeding places. By digging parallel ditches at about 200 feet intervals, and providing a free opening into arms of the sea at each end of the ditches, it has been found possible to eliminate nearly all the breeding in the vicinity of Colon and the adjacent military stations. The ditches are made of such depth that they contain water even at low tide (Chamberlain¹⁶³)

As in the case of fresh water swamps and marshes, if permanent measures are impossible on account of the cost, as is often the case, the best method available is dusting with paris green.

In the case of the great salt marsh districts in America, it has been suggested that mosquitoes may be attacked by the use of poison gas mixed with smoke clouds. The practical utility of such a method is very doubtful on account of the possible danger to human and animal life involved by its use.

3. Lakes and Reservoirs.—It will probably be found impossible to drain lakes in most cases though if this is done valuable land may be made available for cultivation. The margins should be kept free from vegetation or may be treated with paris green from a boat by means of a rotary or electric blower. If there is a narrow outlet to the lake it may be possible to install a dam with a sluice by which the level of the water may be raised and lowered at intervals. Good results have been recorded in the U. S. A. by the use of this method in the case of impounded waters.

The level of the water is kept as high as possible during the non breeding season in order to kill the land vegetation up to this level. The drift and floatage accumulated during this period is chiefly collected along the margins of the reservoir. At the commencement of the breeding season the level of the water is lowered as much as possible to strand the floatage and drift. This kills the larvæ in the floatage and leaves a clean water edge round the reservoir. After this raising and lowering of the water level is carried out at frequent intervals (less than two weeks) throughout the breeding season. Drugging operations must supplement fluctuation where heavy central floatage is present.

(See Bibliography Nos 161 162 166 169, 185, 191)

4 Ponds and Tanks.—It may be impossible to drain or fill these on account of the cost or because they form the only water supply. The following measures may be tried —

- (a) The pond may be regularly dusted with paris green.
- (b) Vegetation and overhanging branches may be removed from the margins

- (c) Copper sulphate may be used, to get rid of algæ (see p. 77).
- (d) For bigger weeds a sub aqueous saw may be used. This is a long band of steel with a double saw edge, weighted with lead, and with a rope attached at each end. A man on either side of the pond pulls the rope, thus cutting the weeds.²⁶⁹
- (e) One end of the pond may be deepened, so that when the water falls there will be a good place for fish, with which the pond may be stocked, if they are not already present. It is a good plan to sink a barrel in the deepest part of the pond to furnish a retreat for the fish during droughts.
- (f) If the outlet can be controlled, the water level may be raised and lowered as in (3).

Of the above methods the use of paris green is probably the most effective, most economical, and most easy to apply.

5. Large River Beds.—These are usually very difficult to deal with. Good may be done, however, by draining pools into the main stream, filling in pot holes and clearing vegetation. Pools left behind after the subsidence of floods may be treated with cresol, or with paris green.

6. Small Streams.—The ideal to be aimed at is that the stream should have steep banks directly above and below the flow line, uniform grade and width, and a straight course, and be free from grass, sticks, stones, or other obstructions which would interfere with the current. With these objects in view the course may be straightened out where possible and the bottom regraded in places, so as to confine the water within narrow banks and increase the rate of flow. In places where there are embayments a channel may be constructed with a line of stones or stakes, the space behind being filled with earth and stones rammed down. Vegetation along the bank should be cut away, and the banks themselves sloped to an angle of about 45 degrees. Pot holes in

stream beds may be filled with stones and pools emptied by draining them into the main stream. Where the bottom of the stream is soft so that there is a tendency for the formation of large pockets in the channel stone may be rammed into place to prevent further excavation.¹⁸⁴

In certain cases dams may be installed with sluices which may be opened at intervals to flush out the larvae. Watson⁴⁵ in Malaya treated hill streams with success both by subsoil drainage and by oiling. Dusting with paris green may also be practised.

7. Hill foot Seepages. These are best treated by constructing a system of hill foot contour drains to catch the seepage at the point where it arises (Fig 13). These may be either open or subsoil drains if open regular oiling will be necessary. Occasionally it may be possible to deal with seepages on the banks of a ravine stream by damming the ravine and submerging the springs.⁴⁶

8 Rice-fields.—These are in themselves rarely dangerous except in hilly tracts where water is in continuous motion through the fields. The cultivation of rice in the beds of streams in such districts is making small artificial bunds is especially dangerous. The problem is a difficult one and the only really satisfactory solution would appear to be a change in the method of cultivation brought about by improved education in agriculture or by law. Drying off of the fields throughout a tract on one day of each week has been generally employed in Java. As regards temporary measures the employment of paris green which does no harm to the crop is the most useful at our disposal. Uncultivated plots in terraced fields which are allowed to become flooded are especially dangerous and may be treated by oiling.⁴⁷

It has also been suggested that the problem of mosquito breeding in rice fields may be open to attack by biological methods and investigations in this direction have been made by Williams⁴⁸ Purdy⁴⁹ and others (see p. 87).

9 Irrigation Channels and Ditches.—These are important sources of anopheline breeding, unless they are properly looked

after. No leakage should be allowed from them and their margins should be kept free from vegetation. It should also be seen that no pot holes are present in the beds of such channels as these will form breeding places when water is not flowing through them. It is however, difficult to enforce regulations to this effect, and the best way of dealing with them in most cases is probably regular treatment with paris green.

10. Irrigation and Water-logging.—The term 'water-logging' as defined by Gill¹⁷⁰, is confined to areas where owing to the height of the subsoil water the superimposed soil is kept permanently damp by moisture derived from the subsoil, as contrasted with 'false water logging' which refers to dampness of the soil caused by the presence of an impermeable stratum preventing for a considerable time the downward percolation of rain water.

The question of the influence of irrigation and water logging on the incidence of malaria is a complex one, and its aspects differ widely in different localities. In India, in the Punjab it has been repeatedly shown that where canal irrigation gives rise to water logging a grave degree of endemic malaria associated with a constantly high spleen rate results¹⁷¹. Here, irrigation if unaccompanied by adequate drainage schemes will tend to raise the level of the subsoil water and favour the formation of swamps, whilst by raising the relative humidity of the atmosphere it will create conditions favourable for mosquito life. The conditions in Bengal on the other hand are entirely different. Here the country is low lying and it has been observed that areas which are regularly flooded are the least malarious whilst those protected from flooding by embankments suffer most. Bentley¹⁵⁸ holds that the adoption of a comprehensive scheme of inundation irrigation from the main rivers would be an effective method of controlling the incidence of malaria in this part of India.

11. Depressions in the Ground should be filled in where possible. For this purpose town rubbish may be used, covered with a layer of earth. Care must be taken that no potential water

containers such as old cans etc., are left exposed. The work should be done preferably in the cold weather to avoid the breeding of flies. Where water is present the rubbish may be shot into it at any season provided it is not allowed to project above the surface, for dangerous anophelines will practically never breed in water thus fouled. Otherwise in summer the rubbish should be dried and burnt and the ash used to fill in the depression. Cinders, sawdust and waste products from factories may be used when available.

12. Borrow-pits.—These are usually harmless when first excavated except where they tap seepage springs. It cannot be seen whether this is the case after rain, and therefore in foot hill localities even fresh pits should be treated as dangerous. Old borrow pits containing vegetation frequently form dangerous breeding places. A row of borrow pits may be drained into one pit so that there remains only a single pool to deal with. This may then be treated with oil or paris green or may be drained by driving a vertical shaft through an impervious stratum. In the case of borrow pits alongside railway lines those near stations may be filled with ashes etc. None should be allowed within distant signals of stations (Senior White⁸³). In certain cases it might be possible to treat a row of borrow pits by the side of a railway line by running a trolley along it and dusting them with paris green.

13. Small Pools formed by showers of rain may often be brushed out with a broom dipped in oil. If too large for this, they may be treated with oil balls or paris green.

14. Holes in Rocks and Trees may be filled with earth or gravel or with cement. In the case of trees the cement may crack and it is probably preferable to use asphalt.

15. Wells.—The only completely satisfactory method of dealing with wells, short of actually filling them in is to provide them with a cement-concrete cover with, if desired the provision of a pump. Trap-doors should be avoided as even if they are

mosquito proof when first installed they soon become ineffective from the effects of constant use and warping. Moreover they are very frequently left open at night⁵⁶

The use of petrol in wells was recommended by Ryles and Majumdar³⁷¹ in the strength of 24 ozs per 80 square feet of water surface (see p 51). Liquid paraffin might also be employed the water being withdrawn by means of a pump.

Hackett⁴¹⁷ suggested the use of paris green stating that in the strength of 1 gm to 90 square metres of surface its use is quite safe. Experiments in this direction have been recently carried out in Bangalore. There are however objections to placing even a harmlessly minute amount of an arsenical preparation into drinking water apart from the fact that in India there is frequently violent opposition to any interference with wells on religious grounds.

Experiments have also been made by Williamson²⁹⁰ as to the larvicidal effect of various gases and vapours in wells.

In the case of disused wells *Lemna* or *Hoffia* may be introduced with the object of forming a dense growth on the surface and preventing the respiration of the larvæ.

Wells may also be stocked with larvæ eating fish (see p 79).

16 Cisterns—These should be covered either with sheet iron or cement concrete. Corrugated iron covers are unsatisfactory and should not be permitted. The man hole should be circular and provided with a cap cover fitting over a rim round the opening. The cover of the man hole should be kept locked. Inlet pipes should be fitted with check nuts and should never enter the cistern through the man hole. Overflow pipes should be provided with a perforated cap of brass or copper. The use of valve boxes should be discouraged where possible as their covers are rarely if ever mosquito proof. The work of inspecting cisterns to see that they are in a mosquito proof condition is greatly aided if each cistern is clearly numbered with white painted figures at least one foot high⁵⁸

17 Fire buckets may be treated with cresol or lime

18 Rain water gutters along the eaves of houses are often sources of mosquito breeding. A gradient of at least one inch ⁱⁿ 24 feet is sufficient for large buildings provided the slope is a permanent one which is ensured by making the gutter with a gradually increasing depth towards the outlet³⁵. If there is any part in which the water stagnates it may be dealt with temporarily by boring a hole at the most dependent point. The ideal solution and that usually adopted in yellow fever campaigns is the total prohibition of all rain water gutters.

19 Water used in Building Construction—Mention has already been made of the amount of anopheline breeding which may occur in the water which is poured on to the surface of cement-concrete whilst it is setting. This may be treated with cresol (see p. 73). Breeding may also occur in the water receptacles used for soaking bricks used in building. Unless it can be ensured that these receptacles are emptied daily after use cresol may be applied in this case also³⁶.

20 Stand pipes and Hydrants—If these cannot be connected directly with the main drainage system a useful plan is to construct soakpits two to three feet in diameter and about one foot in depth round each. The pits may be filled with stones or broken brick covered with a layer of gravel. Another method is by alternation of flow. Two drains are dug leading away from each stand pipe the water being finally allowed to flow over the surface of the soil or to irrigate a plot of ground. Each drain is used for a few days at a time whilst the area supplied from the other is allowed to dry up (Sergeant and Sergeant⁴¹).

21 Leaking Sluice valves—These may be dealt with by filling the valve chambers with sand to the requisite level.

PART IV.

RECENT ADVANCES IN MALARIA CONTROL

In the following pages an account is given of the more important developments in malaria control which have been introduced during the period which has elapsed since this book was first published. The various measures discussed are dealt with in the same order as in the original edition.

A—PROTECTION AGAINST BITES OF MOSQUITOES

1 Repellents—In the U S A the New Jersey pyrethrum larvicide has been extensively used as a spray applied to grass bushes etc. for the protection of outdoor gatherings against the attacks of adult mosquitoes. The formula for the stock solution is given on p 110. This is diluted with 10 to 12 parts of water before spraying and is said to afford protection from mosquito annoyance for a period of several hours (Ginsburg¹⁰¹¹, Vannote¹⁰¹⁰).

The anti mosquito pomade referred to on p 4 as Dover's formula is now widely used in India and has been adopted in a slightly modified form by the Army for use under field conditions.

(See *Bibliography* Nos 1011–1020)

2. Screening—A number of papers have been published regarding the maximum size of mesh permissible for the exclusion of mosquitoes in different parts of the world. Mulligan and Majid¹⁰³⁵ concluded that 14 mesh and 28–30 S W G (aperture 0.055 to 0.057 inch) was the optimum for general use in India. Hargreaves⁶⁴⁵ found that 14 mesh and 30 S W G (aperture 0.059

inch) excluded *A. funestus* in Uganda, but Davey and Gordon¹⁰²⁶, working in West Africa, recorded that both this species and *A. gambiae* could pass through this size of opening, and that some specimens of *A. funestus* escaped through apertures as small as 0.050 inch. They considered that the maximum size of aperture advisable for screening houses in West Africa against anophelines was 0.047 inch, represented by a standard screen cloth of 16 mesh and 28 S. W. G. The same authors described the method by which mosquitoes make their way through narrow apertures, and concluded that their ability to do so is limited by the dorso-ventral diameter of the thorax, confirming in the main the original observations of Russell and Nono¹⁰³⁷.

(See Bibliography, Nos 1021-1039)

B—MEASURES DIRECTED AGAINST ADULT MOSQUITOES

1. **Traps.**—Traps are rarely used as a measure of mosquito control, but a combined method of fumigation and trapping has been employed for this purpose in barracks in India (James⁹⁰⁸). A piece of black cloth, with a central opening 2 feet in diameter, is stretched over a window. The mouth of a muslin bag 6 feet long is sewn round the margin of the opening and the bag itself is passed through the window and attached to the verandah posts, or other available support. The door and the remaining windows are closed and curtained, and the room fumigated by burning pyrethrum coils. The mosquitoes fly towards the opening in the cloth, which is now the only source of light, and are caught in the bag.

Mechanical traps, with light as the source of attraction, fitted with electrically operated suction fans, have been used extensively in the U. S. A. for experimental purposes. A similar apparatus, equipped with a mercury vapour lamp, has been widely advertised in India in recent years for the control of mosquitoes. Several papers recording observations with this machine have been published, but the authors are generally agreed that its high cost

and limited sphere of action renders it ineffective as a method of malaria prevention (Wats and Bilderbeck¹¹⁰⁷, Senior White¹¹⁰⁴, Syddiq¹¹⁰⁶)

(See *Bibliography* Nos 1084-1108)

2 Sprays—The spray killing of adult mosquitoes is now recognized as a major control measure and in the opinion of the author the development of this method represents the most important advance which has been made in malaria prevention in recent years

Swellengrebel¹⁰ * and his co workers in Holland have used the method for many years but it was not adopted extensively in the tropics until the publication of the remarkable results achieved with pyrethrum sprays in rural areas in South Africa from 1932 onwards (Ross¹⁰⁶⁶, de Meillon¹⁰⁶⁰ Booker¹⁰⁴⁴) It has been widely used for controlling malaria in native huts in Natal and Zululand and for the protection of railway and other employees and has been found especially effective in combating outbreaks of malaria in epidemic form Thornton¹⁰⁷⁵ has drawn attention to the fact that the effects are produced not by a diminution in the number of anophelines but by a reduction in their rate of infection which may be very marked a fact which has been confirmed by our experiences in India

Spray killing of adult mosquitoes was brought into use in Delhi in 1936 when the quarters occupied by four communities of Government employees living in particularly malarious sections of the urban area were regularly sprayed throughout the malaria season (Covell Mulgan and Afridi¹⁰⁴⁸) The results were so striking that the method was immediately recommended for use throughout India as being particularly suitable for personnel such as police railway forest or other Government employees and labour forces on estates mills and other industrial enterprises where housed in permanent quarters It was at first thought that the usefulness of this method would be limited to such conditions but in 1937 it was introduced as an experimental measure

in two villages on the outskirts of Delhi. The results were so encouraging that it has since been extended to a number of other villages in Delhi Province.

Reports on the efficacy of the spray killing of adult mosquitoes from employers of labour, railway authorities and Government officials from all parts of India have been universally favourable, and Russell and Kripe¹⁹⁰⁸ have reported that this method has had a marked effect in reducing malaria transmission in a typical South Indian village.

Further tests on the various locally prepared mixtures which have been recommended as sprays and on a number of the proprietary preparations now on the market have shown that those which contain pyrethrum are invariably far more efficacious than the others. A very effective spray can be prepared by merely soaking one pound of dried pyrethrum flowers, previously powdered in 1 gallon of kerosene oil for 72 hours provided that the flowers are of good quality. It may however be found more convenient to use one of the trade extracts of pyrethrum (e.g. *Pyagra*, *Pyroicide*, *Pyefly*, *Pyrentol* etc.) which are diluted with kerosene oil before use.

Pyrethrum cultivation on a commercial scale was formerly almost entirely confined to Dalmatia and Japan but in recent years it has become an important industry in Kenya. It is now being grown in several localities in India (Kashmir, North West Frontier Province, Punjab, Nilgiri Palni and Cardamom Hills) and flowers from all these sources have been shown to possess excellent mosquitocidal properties. It is expected that an ample supply of Indian grown pyrethrum will shortly be on the market so that an effective mosquitocide will be available throughout the country at a comparatively low price (Covell¹⁹¹⁷).

The efficacy of the method is in direct proportion to the frequency of its application. In areas where the infection rate among the carrier species of anopheline is low, good results have been recorded by spraying once a week only. Where the infection rate

is high it is advisable to spray twice or even three times a week throughout the malaria season. It is seldom possible to spray more often than this on a large scale but in very malarious localities residents should be encouraged to spray their living rooms and those of their servants daily while transmission is at its height especially when a case of malaria has actually occurred on the premises or in the immediate vicinity. Particular attention should be directed to the spraying of rooms occupied by young children and infants who constitute the chief source of infection for mosquitoes.

The spray should be mixed by a responsible person to minimize the risk of theft. All apertures should be closed as far as possible before spraying and should preferably remain closed for 20 minutes thereafter. Pieces of sacking, old blankets, sheets etc. are useful for this purpose. If there are any cowsheds, stables or other attractive shelters for mosquitoes nearby these should be sprayed also.

The amount to be sprayed in any particular case can only be determined by actual trial. For a room in the usual type of menial quarter or village in India (capacity about 1000 cubic feet) half an ounce of the diluted mixture is usually sufficient but this may have to be increased if any apertures remain unclosed. The fact that rooms cannot be completely sealed is by no means an indication that spraying will be ineffective and excellent results have been achieved under such circumstances by using rather more of the spray than would otherwise be required.

Regular spraying should be commenced when the local carrier species of mosquito begins to enter the houses at least a fortnight before the malaria season is expected to start and should be continued throughout the transmission period. The most suitable time of day for spraying is in the early morning before cooking has commenced. The personnel employed should be recruited from among the local residents and it is best to engage for this purpose boys of under 14 years of age as there is not

likely to be any objection to them entering rooms occupied by women. The exercise of tact is necessary in introducing the method but once the routine has been established, it is invariably found to be extremely popular among the persons concerned. After a while it may be found possible to leave the actual spraying to the local inhabitants but periodical supervisory visits are essential for it is the universal experience of sanitarians that the individual if left entirely to himself will seldom lift a finger to safeguard his own health, or that of his family.

Hitherto the chief difficulty in applying the method has been the lack of a suitable type of sprayer. For small rooms the hand sprayers or atomizers obtainable everywhere at a trifling cost are useful but they are not very durable and are uneconomical in use. Far the most effective apparatus at present available is an ordinary power spray painting machine driven either by an electric motor or a small petrol engine. These are especially valuable for use in large buildings such as barracks.

As regards costs the expenditure in the Delhi villages in 1937 worked out at one anna per head per month. This does not include the salary of the officer in charge who would normally exercise supervision over a number of villages. The costs given by Russell and Knipe for 7 months spraying in a Madras village amounted to Re 0.15.3 per head. In South Africa Thornton reported the cost of spraying for 6 months at £ 0.2.5 per hut. De Meillon states that the spray killing of adult mosquitoes proved only one third as costly as that of larval control measures conducted over the same area of operations.*

Two great advantages of spray killing of adult mosquitoes are (1) it is the only anti-malaria measure which is universally

* This method is sometimes employed to destroy potentially infective mosquitoes during the winter months. This has no appreciable effect on the numerical prevalence of mosquitoes during the following summer, a finding which is in agreement with R. I. James' original observations regarding winter larval control in Manipal (see *Memorandum Govt. Ind.*, No. 6, 1943). Cold weather control of mosquito breeding has however been a featured in recent years by Pice in Assam.

is high it is advisable to spray twice or even three times a week throughout the malaria season. It is seldom possible to spray more often than this on a large scale but in very malarious localities residents should be encouraged to spray their living rooms and those of their servants daily while transmission is at its height especially when a case of malaria has actually occurred on the premises or in the immediate vicinity. Particular attention should be directed to the spraying of rooms occupied by young children and infants who constitute the chief source of infection for mosquitoes.

The spray should be mixed by a responsible person to minimize the risk of theft. All apertures should be closed as far as possible before spraying and should preferably remain closed for 20 minutes thereafter. Pieces of sacking, old blankets, sheets etc. are useful for this purpose. If there are any cowsheds, stables or other attractive shelters for mosquitoes nearby these should be sprayed also.

The amount to be sprayed in any particular case can only be determined by actual trial. For a room in the usual type of menial quarter or village in India (capacity about 1000 cubic feet) half an ounce of the diluted mixture is usually sufficient but this may have to be increased if any apertures remain unclosed. The fact that rooms cannot be completely sealed is by no means an indication that spraying will be ineffective and excellent results have been achieved under such circumstances by using rather more of the spray than would otherwise be required.

Regular spraying should be commenced when the local carrier species of mosquito begins to enter the houses at least a fortnight before the malaria season is expected to start and should be continued throughout the transmission period. The most suitable time of day for spraying is in the early morning before cooking has commenced. The personnel employed should be recruited from among the local residents and it is best to engage for this purpose boys of under 14 years of age as there is not

likely to be any objection to them entering rooms occupied by women. The exercise of tact is necessary in introducing the method but once the routine has been established, it is invariably found to be extremely popular among the persons concerned. After a while it may be found possible to leave the actual spraying to the local inhabitants but periodical supervisory visits are essential for it is the universal experience of sanitarians that the individual if left entirely to himself will seldom lift a finger to safeguard his own health or that of his family.

Hitherto the chief difficulty in applying the method has been the lack of a suitable type of sprayer. For small rooms the hand sprayers or atomizers obtainable everywhere at a trifling cost are useful but they are not very durable and are uneconomical in use. But the most effective apparatus at present available is an ordinary power spray painting machine driven either by an electric motor or a small petrol engine. These are especially valuable for use in large buildings such as barracks.

As regards costs the expenditure in the Delhi villages in 1937 worked out at one anna per head per month. This does not include the salary of the officer in charge who would normally exercise supervision over a number of villages. The costs given by Russell and Knipe for 7 months spraying in a Madras village amounted to Re. 0.15.3 per head. In South Africa Thornton reported the cost of spraying for 6 months at £ 0.3.5 per hut. De Meillon states that the spray killing of adult mosquitoes proved only one-third as costly as that of larval control measures conducted over the same area of operations.*

Two great advantages of spray killing of adult mosquitoes are (1) it is the only anti-malaria measure which is universally

* This method is a method employed to destroy potentially infective mosquitoes during the winter months. This has no appreciable effect on the numerical prevalence of mosquitoes during the full winter summer, a finding which is in agreement with Sir James's original observations regarding winter larval control in Mian Mir (See *Memoirs of the Entomological Society of India* No. 6, 1932). Cold weather control of mosquito breeding has however been advocated in recent years by Sir J. C. G. in Assam.

popular among the persons whom it is designed to protect, and (ii) it is the only anti-malaria measure which can exert an immediate effect on the course of an epidemic. In India, the fact that pyrethrum of high quality can be grown in a number of different localities makes it certain that the cost of the spray will eventually be greatly reduced, and that its use will be widely extended in future years.

The spraying of aircraft with pyrethrum insecticides has been practised as a routine measure for some years as a yellow fever control measure in America, Africa and India (Whitfield¹⁰⁸⁰).

(See Bibliography, Nos. 1040-1083)

3 Zoophylaxis—A number of puzzling problems connected with this subject have been clarified in recent years by the discovery that a single species of anopheline may be made up of a number of varieties, or subspecies which, though apparently identical are biologically distinct differing widely in their habits and food preferences. In Europe, *A. maculipennis* is now known to comprise at least six forms, two of which, *typicus* and *melanoon* have but little contact with man whereas two others, *labranchiae* and *sacharovi* habitually enter houses, whilst the remaining two *messee* and *atroparvus*, are easily diverted by domestic animals but overflow into human habitations in search of food under certain circumstances. *Typicus*, *messee* and *melanoon* are associated with malaria only in exceptional circumstances, *labranchiae* and *sacharovi* are invariably dangerous vectors, whilst *atroparvus* occupies an intermediate position being as a rule not dangerous but capable under certain conditions of becoming a carrier of considerable importance (Hackett and Missiroli¹¹¹⁵). * The various varieties, or subspecies are distinguishable by certain differences in the pattern of the egg.

* It should be noted that the term *anthropophilous* as applied to biological races of mosquitoes in Europe denotes those which attack man and animals indiscriminately. *zoophilous* indicates races which attack animals by preference.

In India *A. fluviatilis* in the Wynad district of Madras Presidency is a most dangerous malaria carrier feeding freely on man, whilst an apparently identical species in the Terni area of the United Provinces is almost exclusively a cattle feeder and is relatively harmless. Sweet and Rao¹¹³⁸ have described a variety of *A. stephensi* distinguishable from the type form by the characters of the e.g. and probably biologically distinct. It appears likely, on epidemiological evidence that other species e.g., *A. culicifacies* may likewise include more than one biologically distinct variety or subspecies (Senior White¹¹³⁹).

These observations have explained certain problems which were previously obscure as for instance why one locality is malarious whilst another where the anopheline fauna is apparently the same is non malarious and it has been suggested that further study of the various biological strains may lead to a method of species control whereby relatively harmless races may be substituted for dangerous ones. Hitherto however there has been little indication that animal prophylaxis is likely to become an important measure of malaria prevention.

(See Bibliography Nos 1109-1148)

4 Housing and Malaria—The suggestion has sometimes been made that the incidence of malaria might be diminished if human sleeping quarters were rendered unattractive as resting places for anophelines but the investigations of Barber and Rice⁶⁴ lend no support to this view. There is no evidence that the occupants of a house suffer from malaria in proportion to its attractiveness for mosquitoes for the daytime resting places of these insects afford no indication as to where they are likely to be found at night. Christophers and Missiroli⁶⁵, in a comprehensive review of the subject concluded that the only effective method of controlling malaria through housing was by

popular among the persons whom it is designed to protect, and (ii) it is the only anti malaria measure which can exert an immediate effect on the course of an epidemic. In India, the fact that pyrethrum of high quality can be grown in a number of different localities makes it certain that the cost of the spray will eventually be greatly reduced, and that its use will be widely extended in future years.

The spraying of aircraft with pyrethrum insecticides has been practised as a routine measure for some years as a yellow fever control measure in America, Africa and India (Whitfield¹⁰⁹⁰).

(See Bibliography, Nos 1040-1083)

3 Zoophylaxis.—A number of puzzling problems connected with this subject have been clarified in recent years by the discovery that a single species of anopheline may be made up of a number of varieties or subspecies which, though apparently identical, are biologically distinct, differing widely in their habits and food preferences. In Europe, *A. maculipennis* is now known to comprise at least six forms, two of which *typicus* and *melanoon* have but little contact with man, whereas two others *labranchiae* and *sacharovi* habitually enter houses, whilst the remaining two *messex* and *atroparvus*, are easily diverted by domestic animals but overflow into human habitations in search of food under certain circumstances. *Typicus*, *messex* and *melanoon* are associated with malaria only in exceptional circumstances, *labranchiae* and *sacharovi* are invariably dangerous vectors, whilst *atroparvus* occupies an intermediate position being as a rule not dangerous, but capable under certain conditions of becoming a carrier of considerable importance (Hackett and Missiroli¹¹¹⁵). * The various varieties, or subspecies, are distinguishable by certain differences in the pattern of the egg.

* It should be noted that the term *anthropophilous* as applied to biological races of mosquitoes in Europe, denotes those which attack man and animals indiscriminately. *zoophilous* indicates races which attack animals by preference.

In India *A. fluviatilis* in the Wynad district of Madras Presidency is a most dangerous malaria carrier feeding freely on man, whilst an apparently identical species in the Terai area of the United Provinces is almost exclusively a cattle feeder and is relatively harmless. Sweet and Rao¹¹²⁶ have described a variety of *A. stephensi* distinguishable from the type form by the characters of the egg and probably biologically distinct. It appears likely, on epidemiological evidence, that other species, e.g. *A. culicifacies* may likewise include more than one biologically distinct variety or subspecies (Senior-White¹¹²⁷).

These observations have explained certain problems which were previously obscure, as, for instance why one locality is malarious whilst another where the anopheline fauna is apparently the same is non malarious, and it has been suggested that further study of the various biological strains may lead to a method of species control whereby relatively harmless races may be substituted for dangerous ones. Hitherto however there has been little indication that animal prophylaxis is likely to become an important measure of malaria prevention.

(See Bibliography Nos 1109-1148)

4 Housing and Malaria.—The suggestion has sometimes been made that the incidence of malaria might be diminished if human sleeping quarters were rendered unattractive as resting places for anophelines but the investigations of Barber and Rice⁶² lend no support to this view. There is no evidence that the occupants of a house suffer from malaria in proportion to its attractiveness for mosquitoes for the daytime resting places of these insects afford no indication as to where they are likely to be found at night. Christophers and Missiroli⁶³, in a comprehensive review of the subject concluded that the only effective method of controlling malaria through human habit is screening.

C—MEASURES DIRECTED AGAINST LARVÆ OF MOSQUITOES

1. **Drainage**—In the history of malaria control there are numerous examples of the truism that a method suitable for one set of conditions may be ineffective or even deleterious in another. The divergent views expressed by malaria workers regarding the lay out of drainage systems are thus largely explicable by the existence of local differences in physiography and climate. For instance Le Prince and Orenstein⁵⁴ as the result of their experiences in Panama stated that the outlet of subsoil drains must be well above the body of water into which they discharge their contents. This recommendation which was endorsed in the first edition of this book has been stigmatized by Schirff⁵⁵ in Malaya as misleading and likely to lead to mistakes in design since the placing of pipes too near the ground surface is a frequent fault. Again whilst in Malaya great stress is rightly laid on the importance of securing the lowest possible outfall for drains in general in Delhi the outfall of storm water drains is now placed at the highest possible level consistent with the provision of a satisfactory flow in order to minimize the heading up of water when the River Jumna is in flood.

In recent years the choking of subsoil drains by the roots of trees and grasses has become a serious problem on rubber estates in Malaya and in some instances a reversion has been made to the former system of open drains and oiling. Barrowman⁵⁶ recommends the provision of a double line of pipes in such circumstances laid one above the other. The roots grow into the lower line first and though the upper one too may ultimately become choked it is claimed that the effective life of the system is thereby prolonged by at least sixfold.

A development of subsoiling which has yielded strikingly good results in Malaya is that known as *fascine drainage* originally introduced by Scott⁶⁰. A layer of sticks each about 6 feet long and 2 inches in diameter is placed in the bottom of the drain or stream which has been previously graded and straightened the

thickness of the layer depending on the depth of the drain and the amount of water to be disposed of. A series of layers of coarse grass (*balling*) each overlapping the next below it like thatch on a roof is placed over the sticks. Over this, the trench is filled with heavy earth. A modification of this method is to lay the sticks against the lower 6 inches or so of the sides of the trench fixing them in place by vertical pegs. Other short sticks are laid crosswise supported by the top layer of the lengthwise sticks. A layer of long sticks is now applied lengthwise as a roof to the channel these being supported in their turn by the crosswise sticks. The work is then completed as already described*. It was originally expected that fascine drains would last only for a few months but many of them are still functioning perfectly after a lapse of several years. It appears that in Malaya not only sticks which are continuously in water but also those laid in drains with only a seasonal flow, have astonishing powers of survival.

Subsoil drains with the channel made of flat stones as figured on p. 40 are also in common use in Malaya e.g. for dealing with seepage outcrops in quarry pits under the designation of *stone packing*. This method has also been found useful under similar conditions in the Ridge quarry pits in Delhi.

A method of drainage which has met with great success in Delhi is that known as the *hilo* system which is employed when for financial or other reasons it is found impossible to remove water completely from a particular area. The entrance of extraneous water into the area in the form of river flooding or storm water is excluded by bunds if necessary and the area thus protected is so dressed that all rain-water falling on it is directed into a single depression or *hilo* which is deepened and widened for the purpose. In this way large numbers of breeding places are abolished, and the single remaining water collection can easily be kept

* This procedure is sometimes locally referred to as *kerbaya corer*, a term which should properly be reserved for the method described on p. 11*

clean weeded and effectively treated with larvicides or stocked with fish. An important feature of this method is that whilst every advantage is taken of the natural slope of the ground no formal drains are constructed so that maintenance charges are practically non-existent. The method is most effective in areas where the annual rainfall is moderate in amount and it can be carried out for a fraction of the cost entailed by filling operations (Covell and Afridi⁵⁸¹).

Before leaving the subject of drainage particular attention is drawn to the value of turfing the sides of drains to minimize the effects of scour. A special technique for planting and growing grass for use in anti malarial drainage projects has been described by Scharff⁷⁶.

(See *Bibliography* Nos 732-764)

2 Oiling—A number of important researches have been carried out in recent years regarding the method of action of oil on mosquito larvæ and the essential qualities required for effective larval control. These have led to some modification of the generally accepted ideas on the subject. Allen⁶⁰⁵ sums up the present position as follows—

For a long time we have known that we want an oil combining high toxicity with low volatility (which may occur together but are not necessarily related) low viscosity and good spreading properties. Recent studies the most important by Murray⁶⁰⁰ seem to establish the following additional important points—

- (i) Reasonable toxic properties are sufficient, the highest possible toxicity is neither necessary nor desirable
- (ii) A perfect film is not absolutely necessary to ensure good results. Small breaks in a film of good durability and toxicity do not permit the escape of larvæ
- (iii) Volatility low enough to ensure a good film for a few hours is sufficient. It is not necessary that the film should remain intact for several days

- (ii) The aromatic vapour pressure and the aromatic-aliphatic ratio should be below the critical figure for film instability
- (i) Spreading pressure should be between 11 and 35 dynes. A greater spreading pressure would be expected to produce too thin a film to be practical
- (iv) The addition of special 'spreaders' to mosquito oils is in general not justified

In India *Malariaol* a proprietary preparation specifically designed for anti-larval purposes is now widely used and has been adopted by the Army authorities. It is claimed that this oil has good spreading powers, is highly toxic to larva (killing them within 15 minutes of application) and burns up the vegetation and grass amongst which the larva shelter along the banks of breeding places to which it is applied. It is however somewhat more expensive than the diesel oil-cresol mixtures formerly in general use and opinions differ as to whether the advantages claimed for it compensate for the increased cost. One decided advantage of this preparation is that it is not likely to be stolen by unscrupulous employees.

In Malaya much experimental work has been done to determine the relative efficacy of various mixtures of oils classed as heavy, solar and light. Heavy oils are those employed for diesel oil engines, solar oils for lighter oil engines and light oils for lighting and cleaning purposes. The effect of adding pyrethrum to such oils has also been tested and it has been shown that this has the effect of shortening the time in which the larva are killed in all cases and that in the case of an oil with high boiling range (Diesel II) it also greatly increases the death rate.

As regards methods of applying oil mention may be made of the brush spray mop routine introduced by Quaise* in Malaya for the treatment of streams and drains. A thin line of oil is sprayed on the water surface for four paces, and this is vigorously

distributed with a brush for 100 yards or more the coolies brushing rapidly with the flow of the water and taking care that all side pockets are treated. The workers then proceed to the next oiling point 100 yards upstream and repeat the performance. The spray coolie deals with any seepage or side channels encountered and finally all small ponds pot holes etc. are treated by a mop dipped in oil. For the sprayer the finest nozzle is used and work is abandoned if heavy rain occurs. This method is claimed to be both economical and effective but constant skilled and intimate supervision is essential for its success.

One of the most difficult problems in anti malaria work in India is the control of *larval drift* in streams and irrigation channels. In Delhi for instance the Western Jumna Canal Tail Distributary is a prolific source of breeding of *A. culicifacies* the larvæ of which were formerly carried down into the urban area in enormous numbers. Oil booms of various designs have proved very effective in dealing with situations of this kind. A considerable degree of ingenuity is required in designing such booms for whilst all larvæ must be held up by them the flow of water must not be obstructed. Moreover they must be able to function equally well at varying water levels. In the case mentioned above a boom has been erected across the canal a short distance above the point where it enters the urban area. It consists of three sections each made of *sarkanda* (moonj) grass in the form of a chick which can be raised or lowered as desired the stems of the grass being parallel with the water surface. When the water in the canal is low, only the central section is used whilst when it is full all three sections are brought into operation. The boom is supported by stakes driven into the bed of the canal and *bhoosa* (chaff) soaked in oil is thrown on to the water above it so that the larvæ trapped by the boom are killed by the oil exuding from the *bhoosa*. It is necessary to provide some arrangement for holding up objects which may float down the stream and damage the booms. A single rope stretched across from bank to bank is usually sufficient for this purpose (Covell and Afridi³¹).

All oil booms need a good deal of attention, and, wherever they are used some one responsible must be constantly on the watch, to prevent unauthorized persons from tampering with them. Put their use leads to economy in oil and they probably constitute the most effective method of dealing with the difficult problem of larval drift. The type of boom suitable in any particular case will depend on local conditions.

As regards sprayers the type found most generally useful in India is the knapsack sprayer (Vermorel I chur or Four Oaks) operated by a hand lever located immediately beneath the right elbow. This pattern is lighter, less costly and less likely to get out of order than high pressure sprayers, and the coolie has no difficulty in regulating the amount of oil ejected from the nozzle.

(See Bibliography Nos. 905-911)

3 Chemical Larvicides *Paris green*—The use of this larvicide has been continued and extended in many European countries and in the U. S. A. but in the tropics it is not generally employed as widely as had been expected when it was first introduced. There are several reasons for this, two of the most cogent being (i) that its application requires a good deal more intelligence and supervision than that of oil and (ii) that it has no effect on the larva of culicines, the control of which is usually demanded by those who provide the necessary funds. Consequently in most tropical countries the application of paris green is limited to situations where local conditions preclude the use of oil. In Delhi, for example, it is found extremely useful for bringing under control stretches of low lying ground which have been subjected to river flooding, due perhaps to a breach in some embankment. Other water collections are for the most part dealt with by oiling as a routine measure.

The application of paris green from aircraft continues to be practised in the U. S. A. to deal with mosquito breeding in impounded waters, notably in the Tennessee Valley, where the

shoreline will eventually exceed 7 000 miles one third of which will need protection (Watson and Bishop⁷⁰⁹) This method is also employed very extensively in Russia where in 1937 breeding places extending over an area of 12 000 square miles were treated It has been found particularly effective for dealing with peat bogs reed beds ricefields and cotton growing areas intersected by a network of irrigation ditches (Nabokov⁹⁷⁸) It has also been used in Madagascar (Legendre⁹⁶⁹ Boyé⁹⁵¹) Morocco (Sicault⁹⁰⁹) and Indo China (Morin and Martin⁹⁷⁴) In Delhi experimental dusting of paris green from aircraft was carried out in 1936 when it was thought likely that this might prove to be the only practicable method of controlling mosquito breeding in the low lying tract bordering the Jumna River The method proved effective but expensive and it was ultimately found possible to deal with breeding in this area with greater thoroughness and at much less cost by other means (Covell and Afridi⁵⁸¹)

Barber and his colleagues⁹⁴⁷ in Greece have described a dustless method of applying paris green which has also been used in India (Russell and Jacob⁹³⁰) in Cyprus (Aziz⁹⁴) and in Russia (Keiris and Klokov⁹⁶⁵ Yurchak and Bozhenko⁸⁰⁸) A stock mixture is prepared composed of kerosene oil 400 c c paris green 200 c c and dry powdered egg albumin 1 gramme* which are poured into a Winchester quart bottle in that order The bottle is vigorously shaken a procedure which is repeated whenever any of the mixture is poured out A number of corked vials are provided into each of which 25 c c of the stock solution is poured before the cooler sets out on his round The vials are carried in a cartridge belt which accommodates 20 to 25 vials each in a separate pouch The cooler also carries a tin can of 1 litre capacity which when not in use is suspended from the belt and a funnel provided with a wire gauze sieve

* In Delhi a mixture of kerosene oil $\frac{1}{2}$ gallon paris green $2\frac{1}{2}$ pounds castor oil 1 ounce and the whites of 4 to 6 eggs shaken up in a petrol can and used in the dilution of 1 ounce to 1 gallon of water has been found effective

On arrival at the scene of operations the coolie pours 2 litres of water into the sprayer from his tin can, filtering this through the funnel to remove débris. He then adds the contents of one vial, after vigorously shaking it, washing the last remnants of the stock mixture into the sprayer with a little more water. Three more cans full of water are added, making 5 litres in all to 25 cc of stock mixture. The tank of the sprayer is agitated from time to time by a sway of the hips during the application of the spray. One vial (i.e. 25 cc) of the stock mixture is sufficient for treating 700 to 900 square feet of water surface and an average Indian coolie uses the contents of 20 to 25 vials per working day (Russell and Jacob⁹⁰⁰, Russell, Knipe and Rao⁹¹¹).

The advantages claimed for this method are that there is no need to store or transport the diluent, that very little paris green is lost by sinking as it is kept afloat by the kerosene, and that there is no dust left on the water surface to block the feeding of the larvæ, which last consideration may explain why so little paris green is found necessary as compared with other methods of application. It seems likely that this technique may be found of great value in special circumstances as for instance during the course of military operations.

De Benedetti⁹¹⁰ working in Italy claims that a practically unsinkable diluent for paris green can be prepared by mixing intimately a small quantity of mineral oil with any siliceous or calcareous dust (except calcium sulphate), and heating it to 250°C to evaporate the excess oil. Paris green applied by means of this vehicle is said to persist on stagnant water and to retain its larvicidal power for 24 to 25 days.

(See *Bibliography*, Nos. 912-1010.)

4. Vegetable Larvicides—*Pyrethrum* has been extensively used in recent years in the U. S. A. in the form of an emulsion usually known as the New Jersey Pyrethrum Larvicide (Ginsburg⁸⁴¹, Vannote and Ginsburg⁹⁰⁰, Smith⁸⁴⁸). Allusion has

shoreline will eventually exceed 7 000 miles one third of which will need protection (Watson and Bishop⁷⁰⁹) This method is also employed very extensively in Russia where in 1937 breeding places extending over an area of 12 000 square miles were treated It has been found particularly effective for dealing with peat bogs reed beds ricefields and cotton growing areas intersected by a network of irrigation ditches (Nabokov⁹⁷⁸) It has also been used in Madagascar (Legendre⁹⁶⁹ Boyc⁹⁵¹) Morocco (Sicault⁹⁹⁹) and Indo China (Morin and Martin⁹⁷⁴) In Delhi experimental dusting of paris green from aircraft was carried out in 1936 when it was thought likely that this might prove to be the only practical method of controlling mosquito breeding in the low lying tract bordering the Jumna River The method proved effective but expensive and it was ultimately found possible to deal with breeding in this area with greater thoroughness and at much less cost by other means (Covell and Afridi⁵⁸¹)

Barber and his colleagues⁹⁴³ in Greece have described a dustless method of applying paris green which has also been used in India (Russell and Jacob⁹⁹⁰) in Cyprus (Aziz⁹⁴) and in Russia (Kenis and Klovov⁹⁶⁰ Yurchak and Bozhenko⁹⁹⁸) A stock mixture is prepared composed of kerosene oil 400 c c paris green 200 c c and dry powdered egg albumin 1 gramme* which are poured into a Winchester quart bottle in that order The bottle is vigorously shaken a procedure which is repeated whenever any of the mixture is poured out A number of corked vials are provided into each of which 25 c c of the stock solution is poured before the coolie sets out on his round The vials are carried in a cartridge belt which accommodates 20 to 25 vials each in a separate pouch The coolie also carries a tin can of 1 litre capacity which when not in use is suspended from the belt and a funnel provided with a wire gauze sieve

* In Delhi a mixture of kerosene oil $\frac{1}{2}$ gallon paris green $\frac{1}{4}$ pounds castor oil 1 ounce and the whites of 4 to 6 eggs shaken up in a petrol can and used in the dilution of 100 to 1 gallon of water has been found effective

On arrival at the scene of operations, the coolie pours 2 litres of water into the sprayer from his tin can, filtering this through the funnel to remove debris. He then adds the contents of one vial, after vigorously shaking it, washing the last remnants of the stock mixture into the sprayer with a little more water. Three more cans full of water are added, making 5 litres in all to 25 cc of stock mixture. The tank of the sprayer is agitated from time to time by a sway of the hips during the application of the spray. One vial (i.e. 25 cc) of the stock mixture is sufficient for treating 700 to 900 square feet of water surface, and an average Indian coolie uses the contents of 20 to 25 vials per working day (Russell and Jacob⁹⁹⁰, Russell, Knipe and Rao⁹⁹¹).

The advantages claimed for this method are that there is no need to store or transport the diluent; that very little paris green is lost by sinking, as it is kept afloat by the kerosene, and that there is no dust left on the water surface to block the feeding of the larvae, which last consideration may explain why so little paris green is found necessary as compared with other methods of application. It seems likely that this technique may be found of great value in special circumstances, as for instance during the course of military operations.

De Benedetti⁹¹⁶ working in Italy claims that a practically unsinkable diluent for paris green can be prepared by mixing intimately a small quantity of mineral oil with any siliceous or calcareous dust (except calcium sulphate), and heating it to 250°C to evaporate the excess oil. Paris green applied by means of this vehicle is said to persist on stagnant water and to retain its larvicidal power for 24 to 25 days.

(See *Bibliography*, Nos. 942-1010)

4. Vegetable Larvicides—*Pyrethrum* has been extensively used in recent years in the U. S. A. in the form of an emulsion, usually known as the New Jersey Pyrethrum Larvicide (Ginsburg⁸⁴¹, Vannote and Ginsburg⁸⁴⁰, Smith⁸⁴⁸). Allusion has

already been made to its employment as a repellent for outdoor meetings (p. 94)

A stock mixture is prepared as follows. Four gallons of soap solution made by dissolving 16 pounds country soap in hot water and adding 48 ounces methylated spirit, are agitated in 30 gallons of water till a foam is produced. An oil mixture composed of pyrethrum extract (e.g., *Pyrocide* 20) 3 gallons, pine oil 3 gallons and kerosene oil 60 gallons previously prepared and thoroughly mixed is added to the soap mixture with vigorous agitation till an emulsion is formed. Before use the stock mixture is diluted with ten times its volume of water taken from the breeding place. The sprayer must be agitated from time to time in the course of application.

The advantages claimed for this larvicide are that it is cheaper and quicker in action than oil, easily transportable, clean and easy to handle, kills larvæ and pupæ of both anopheline and culicine mosquitoes, is harmless to fish and vegetation, and does not rot rubber hose and washers of sprayers as quickly as does oil. A serious disadvantage is that the larvicidal properties of the stock mixture rapidly deteriorate and it has to be made up fresh each day in order to obtain optimum results.

More recently, Ginsburg⁸⁴ has evolved an improved emulsion for use on fresh or salt water. The formula is 100 U.S. gallons of kerosene containing pyrethrum extract equivalent to 100 pounds of flowers, 50 U.S. gallons of water, and 6 pounds of Gardinol W. A concentrated (sodium laurel sulphate). To avoid excessive foaming 2 to 3 pounds of wool grease (Degras) may be dissolved in the kerosene before it is added to the mixture of water and emulsifier.

The use of cashew nut shell oil as a larvicide has been advocated by Wats and Bharucha⁸⁵. This is a viscous tarry oil obtained from the charred pericarp of the cashew nut, which is a part of the fruit of *Anacardium occidentale*. About 32 000 tons of raw cashew nuts are roasted every year in India and the oil is obtained as a by-product which is used as a preservative by

fishermen for their boats and nets. For use as a larvicide it is mixed with kerosene or with diesel oil in dilutions of from 2 to 5 per cent. The addition of 1 per cent stearic acid in kerosene oil is said to increase the stability of the film. Disadvantages are that the mixture does not retain its efficacy and must be freshly prepared before use; that oils obtained from different sources differ in their toxicity, presumably due to adulteration; and that the oil has an irritant effect when applied to the skin. The authors, however, believe that these disadvantages can be largely overcome and consider that the oil has distinct possibilities as a larvicide on account of its high toxicity to larvæ, rapid action and low cost (Rs. 1.40 per gallon).

5 Naturalistic Control Measures—Under this heading are included those measures by which changes are effected in the natural conditions under which malaria-carrying mosquitoes exist, rendering them unfavourable to the life and activities of these insects, either in the aquatic or adult stages. These have been conveniently classed as chemical, physical and biological measures by Hackett and his colleagues⁶⁵ in their report on an enquiry into the use of such measures, instigated by the League of Nations Malaria Commission, from which much of the information given in the following pages has been extracted.

(a) CHEMICAL MEASURES

(1) *Pollution of water*—Earlier attempts to check the breeding of malarial vectors by this means have already been discussed (p. 83). In Russia, mosquito breeding in peat bog pits has been dealt with by adding sulphuric acid to change the reaction of the water. This was successful where the bed was of clay, but not in the case of a sedge peat bed, where the action of the acid was rapidly neutralized by the alkaline reaction of the peat (Pavlova⁶⁶).

Certain experiments carried out by Williamson⁷¹² in the Cameron Highlands, Malaya, are of great interest. In one locality,

a small stream in which *A. maculatus* had been found breeding in large numbers was converted into a series of ponds in some of which control was effected by the trampling in of cut vegetation whilst in others larvivorous fish were introduced. In another case white clay puddled into the sides and beds of pools and of a stream in which there had previously been a considerable amount of breeding was effective. Williamson's best known method of larval control is that known as *herbage packing* or *herbage cover* in which fresh cut green vegetation is packed into stagnant or slowly running shallow water to a height of 12 to 18 inches and then trampled in wooden stakes being driven in at the lower end of the packed section in the case of streams and drains to prevent the herbage from being washed away by storm water. When packing running water it is essential that there shall be a complete and thick cover of packing above the water level. By this means light is excluded from the water a mechanical barrier to ovipositing mosquitoes is provided and the water is polluted with organic matter the pH in most cases falling to about 6.0 the last mentioned factor probably being the most important. The method has been employed in India with complete success as regards the larval control of malaria carrying mosquitoes but frequent inspection is necessary to ensure that the packing is repaired or replaced in case of damage due to storm water (Senior White⁶³³ Covell and Harbhagwan⁶³⁴ Covell⁶³⁴).

(ii) *Changing the salt content of water*—A method of dealing with brackish water breeders which include some of the most dangerous malaria carriers known is to increase or diminish the degree of salinity in their breeding places by the admission or exclusion of sea water. Near Durazzo Albania a marsh in which *A. saccharovi* formerly bred in profusion was converted into a sea water lagoon by installing automatic tide gates with their usual aspect reversed at the mouth of a channel connecting it with the Adriatic (Hackett⁶⁵⁰). A similar result was achieved in Jamaica by reopening a channel and admitting sea water into a large permanent swamp in which the local vector *A. albimanus* was breeding

In this locality, which was originally non malarious, a virulent epidemic had followed the closing of the above mentioned channel (Washburn⁷⁰⁷)

In the Netherlands the process of reclamation which has been in progress for some years has reduced the breeding of *A. atroparvus* by the washing out of salt from the soil due to abundant irrigation brackish water thus being replaced by fresh. It is confidently expected that the malaria problem in this area will thus be solved (Swellengrebel and de Buck⁶¹²)

(b) PHYSICAL MEASURES

(i) *Filling by silting*—As a general rule, this is a method which can only be carried out successfully by engineers with experience in this type of work, for the results depend upon an appreciation of the volume of silt in suspension, the reduction in velocity necessary to cause this to be precipitated and the disposal of the residual water. On the other hand in the absence of advice from a malarialogist there is a risk that such operations may actually favour the breeding of dangerous malaria carrying mosquitoes especially whilst the work is actually in progress. A marsh can sometimes be filled by diverting a silt bearing stream into it, a method which has been used with success by Strickland and Murphy⁷⁰⁰ in Assam. Loops in streams can sometimes be silted up by the use of groins made with jungle stakes interwoven with withies placed at varying angles a technique which has been extensively practised for river training by Worth⁷¹⁶ in Ceylon.

(ii) *Sluicing**—The principle underlying this method is the sudden release of water confined in the reservoir created by the

* The earliest example of the operation of sluice gates as an anti mosquito measure is probably the procedure originally recommended by Graham at Lucknow in 1913 which has been carried out continuously since 1916 (Raja Ram⁶⁹⁹). The gates of the weir in the River Gamti are lowered once a week from November to May from 5 a.m. to 12 noon, the purpose being to flush out the drains discharging into the river above the weir and to wash out larvae sheltering under the banks into the middle of the stream (Banerjee⁶⁹⁷).

damming of a stream or drain, in order to flush the channel below the dam. A number of larvæ are drowned by the turbulence of the flush itself, and many others, together with eggs and pupæ are stranded on the banks as the wave passes downstream. Under favourable conditions, mosquito breeding can sometimes be controlled throughout several hundred yards of stream by the operation of each sluice. It is usually found necessary to adopt subsidiary control measures above the sluice.

Excellent results can often be secured in small streams and tea garden drains by the use of kuchcha dams with wooden sluice gates, which can be erected at a very low cost. In larger streams, concrete dams are sometimes installed, with gates of wood, or iron which can be raised by means of levers. Another type of gate is provided with hinges at the bottom, and held upright by a wire cable, attached to a ring set in concrete in the bed immediately above the dam. The gate can be made to fall flat on to the stream bed by operating a release clip attached to the cable. Gates of this type are easily operated by a single person and enable the whole contents of the reservoir to be released with extreme suddenness.

In Malaya (Williamson and Scharff⁷¹⁵) and Ceylon (Macdonald⁶⁶⁵, Worth and Subrahmaniam⁷¹⁷), and in some localities in India (Ramsay and Anderson⁶⁸⁴), automatic siphon sluices are in use for this purpose, but these usually need frequent attention and inspection, not only for maintenance purposes, but also because they are liable to interference by unauthorized persons. Moreover, it is not an uncommon experience that whilst an automatic sluice may function admirably when there is a normal flow of water in the stream, it is apt to fail when this is reduced to a minimum by abnormal drought, which is the very time when its effective operation is most needed. It is the author's opinion that hand operated sluices are more suitable for general use, at any rate in India. These are less costly than automatic sluices, are easily constructed,

are not likely to be interfered with require very little attention and can be operated at any desired moment

(iii) *Flooding*—In Bengal the annual flooding of the delta by the rise of water level in the great rivers provides a striking example of natural control of malaria. The silty flood waters are inimical to the breeding of the local vectors the extent of breeding is reduced and the raising of the general water level above that of the aquatic vegetation exposes the larvæ to the attacks of fish and other natural enemies. The highly malarious condition of the western part of the delta is largely attributable to the exclusion of flooding by road and railway embankments and the embanking of tidal rivers to prevent salt water from spilling over reclaimed land. There are obvious grave difficulties in removing existing embankments but it is the declared policy of the Bengal Irrigation Department to abolish embankments step by step wherever practicable and to forbid the erection of any new ones (Curry⁶³⁷ Griffin⁶⁴⁷). Bentley has pointed out that in Bengal flush irrigation is not practised until October by which time most of the season's anopheline carriers have already been produced. He recommends that canals be allowed to run full throughout the flood season so that the largest possible area may be flushed out with silt laden water.

(iv) *Fluctuating water level*—Allusion has already been made to the practice of this method in impounded waters in the U S A. It has been applied in the Tiber delta in the case of a large marsh which has been surrounded with an embankment and is periodically filled from the river by pumping the primary object being to irrigate 1 000 acres of land (Hackett⁶⁵⁰). In Singapore the water level of a large artificial lake in the Botanical Gardens is raised and lowered periodically with complete success as regards mosquito breeding. In Savantvadi south west India the weekly opening of a sluice gate at one end of a lake in the capital of the State serves the double purpose of lowering the water level of the lake and of flushing the stream into which the water is discharged.

(v) *Intermittent drying*—This method originally employed by James in India* has been practised in recent years in irrigated tracts in a number of countries e.g. Algeria (Sergeant and Sergeant⁴¹) Bulgaria (Konsuloff⁶⁶¹) Armenia (Ananyan⁶¹⁹) Russia (Prosolupov⁶⁵² Gnukolopov⁶⁴) Portugal (Hill⁶⁵⁴) Dutch East Indies (Smalt⁶⁵⁰) and Indo China (Morrin⁶⁷¹) In India field experiments have been conducted in Madras by Russell and Rao⁶⁹¹ The principle is first to flood the fields with water and then cut off the supply for a number of days (usually 3 or 4) in order to minimize mosquito breeding in the fields and small irrigation channels In some cases this procedure appears to increase the yield of the crop whilst in others a slight decrease in yield has been reported

(vi) *Agitation of the water surface*—This method is used in Penang where pools on a hill side are fed by water carried from streams by conduits of split bamboo the terminal section being supported at a height of 4 feet or more above each pool Mosquito breeding is prevented by the consequent splashing and rippling of the water surface (Scharff⁶⁹⁰)

In the U S A reduction in mosquito breeding in large bodies of water has been effected by the action of waves created by driving fast launches round the margins of certain lakes (Libbey¹⁵⁶)

(vii) *Stagnating or Ponding*—These terms denote the conversion of a stream into a series of pools by means of dams a method first used by Williamson⁷¹⁰ for the control of *A. maculatus* in Malaya The dams were constructed of earth and timber and pond plants and larvivorous fish were introduced This method has been successfully employed by Ejercito⁶⁴¹ in the control of *A. minimus* var *flaviventris* in the Philippines

(viii) *Muddying*—This is a term used to indicate the introduction of silty water into breeding places not with the object

* James S P (1903) First report of the anti malarial operations at Malacca. *Sci. Memos Govt Ind.* No 6

of filling them as above described but because silt in suspension is in certain cases fatal to malaria-carrying anophelines. This is the basis of the *flood flush* method practised in the Bengal delta by Griffin⁶⁴⁷, whereby the silty waters of the rivers are introduced into drainage channels and borrow-pits by a system of sluices. This is a method which requires considerable engineering skill, and should not be attempted unless this is available.

(12) *Shading* —On p. 36 brief mention was made of the earlier experiments of Ramsay and his co-workers with the growing of shade giving plants and trees over tea garden drains, small streams and swamps in Assam*. This method has been greatly developed in subsequent years and has been practised more recently in tea estates in southern India also. It is estimated that at least 2 000 miles of channels have been rendered free from the breeding of dangerous mosquitoes during the last 10 years. Although shading has proved such a conspicuous success in estates where the population is under strict disciplinary control it is not applicable in uncontrolled rural areas for any break in the shade such as may be produced by cattle or goats or by villagers cutting trees or bushes for fuel will provide a favourable anopheline breeding place.

Ramsay⁶⁴⁸ has also advocated the control of mosquito breeding in tanks by shading their margins with broad sheets of matting whilst in Malaya the shading of seepage wells by huts of thatch is widely practised and small drains are covered with coconut fronds or with a thatch of coarse grass placed over bundles of twigs with a similar purpose (Scharif⁶⁴⁹). Experiments in artificial shading have also been conducted in Mauritius (Birk⁶⁵⁰) Argentina (Alvarado⁶⁵¹) Indo China (Morin⁶⁵²) and the Dutch East Indies (Overbeel and Stoker⁶⁵³).

(c) BIOLOGICAL MEASURES

(1) *Iarvivorous fish*—In the original edition of this book the use of indigenous fish was recommended in localities where species which prey on mosquito larvæ were available. Experience gained during the past 10 years with *Gambusia* has however led to a modification of this view. Hackett⁶⁵⁰ has pointed out that the local fish, after the lapse of centuries have reached a balance with regard to their natural enemies, and their numbers cannot be artificially increased and maintained at an abnormally high level of density, whereas, if an exotic fish can be acclimatized, it will proliferate and maintain its numbers to an extraordinary degree. For instance, *Gambusia* were introduced into Italy in 1922 and have undergone a prodigious multiplication reaching a density far greater than that attained in their native country, contrary to expectation. After the lapse of 18 years, there is no sign of any diminution in their numbers.

Gambusia have been introduced into practically every country where anti-malaria work is in progress. They have been established in many localities in India, and are employed for the control of mosquito breeding in wells in Bangalore and Bombay, and in ornamental waters in Delhi. As has been already explained (p. 81), the utility of fish in the control of malaria is as a rule, strictly limited. It is claimed however, that along the northern coast of Dalmatia, including the peninsula of Istria anopheline density has been reduced below the threshold of malaria transmission by the agency of *Gambusia* alone (Hackett⁶⁵⁰).

(ii) *Changing the fauna and flora*—The best known example of this type of control is the treatment of fish ponds in the Dutch East Indies (Overbeek and Stoker⁶⁷⁷). The beds of the ponds are dried for a couple of days at least once a month, after which they are filled with fresh sea water. During the drying period, the fish remain in ditches dug for the purpose in the beds of the ponds. The top algæ are killed by drying, and the development of blue

algæ which grow on the bed of the pond and on which the fish live is promoted. The algæ rise to the surface in broad sheets which no larvæ can penetrate. Any that may be present in the pond are destroyed by the introduction of larvivorous fish. These measures have resulted in a great reduction in the incidence of malaria in the neighbourhood of Batavia.

(See *Bibliography* Nos 617-719)

D—GENERAL CONDUCT OF ANTI MALARIA CAMPAIGNS

The provision of a special checking staff of insect collectors entirely separate from the executive staff is of immense value in assessing the efficacy of control measures and of ascertaining the lines on which these may need modification. In Delhi where operations are in progress over an area of some 50 square miles there is a staff of seven insect collectors to each of whom are assigned certain catching stations. If an unexplained local increase occurs in the catch of adults of the vector species the whole of the checking staff is concentrated in that particular section until the source of breeding is discovered. The checking staff also make house to house enquiries regarding the occurrence of fever cases in various parts of the urban area (Covell and Afridi⁵⁸¹).

The maintenance of a spirit of enthusiasm among all members of the staff is of vital importance and this depends very largely on the personality of the officer directing the campaign. Most if not all of the measures above described depend for their efficacy on the existence of adequate and unremitting supervision. For this reason one of the most important factors for the furtherance of anti malaria work in the tropics is the development of communications whereby the cost of supervision can be spread over larger areas. Another important factor is the instruction not only of engineers but also of other government officials and employers of labour in the elementary principles of anti malaria sanitation. The institution of courses in malariology for engineers by the Health Organization of the League of Nations and more recei

by the Government of India marks an important advance in this direction

It is only by the diffusion of knowledge among officials as well as the public that we can hope to counteract the natural tendency to curtail expenditure on preventive measures as soon as the incidence of malaria begins to fall and the stimulus to action diminishes particularly in times of financial stringency. There are many instances on record where the very success of anti malaria campaigns has led to the reduction of expenditure to a level at which the scheme can no longer operate effectively resulting in a disastrous recrudescence of the disease. This is well illustrated by the history of malaria in Bombay (Covell⁵⁶)

Most of the advances which have been made in the development of anti malaria measures in recent years owe their origin to an intensive study of the bionomics of malaria carrying anophelines. In the author's opinion our further progress largely depends on a still more intimate association of both field and laboratory researches with the actual prosecution of control measures.

It is perhaps fortunate that in anti malaria work perfection is so seldom attainable. There is always scope for improving existing conditions and for devising new and more effective methods of control, and therein lies the great attraction of this particular branch of public health activity.

BIBLIOGRAPHY *

A GENERAL

- 1 BARNES M F and RUSSELL A programme for the control of malaria in
P F (19 6) Ceylon *Ceyl Soc Paper* No IX
- 2 BLANCHARD R (1904) Les moustiques Paris
- 3 BOYCE P W (1910) Mosquitoes and Man? London
- 4 BOYD J S K (1924) The principles of the prophylaxis of malaria
Jl P i M C XLIII " and 3
pp 95-100 and 182-91
- 5 BOYD M F (19 6) Studies of the epidemiology of malaria in
the coastal lowlands of Brazil Amer
Jl Hyg Monograph Ser No V
- 6 BOYD M F (19 0) An introduction to malariaology Can
berline Ma sa huettis
- 7 BYAM W and ARBAID Irritation of medicine in the tropics I
P G (19 1 ") pp 388-410 II pp 160-16 0 London
- 8 CARTER H F RUSTOMJEE Report on malaria and anopheline mos
K J and SARAVANA quitoes *Ceyl Soc Paper* No VII
MUTTU E T (1927)
- 9 CELLI A (1901) Malaria according to the new researches
London
- 10 CHAMBERLAIN W P (19 9) Twenty five years of American medical
activity on the Isthmus of Panama
1904 19 9 Panama Canal Press
- 11 CRAIG C F (1909) The malarial fevers pp 31-309 New
York
- 12 CRAWFORD J A and Mosquito relation and malaria preven
CHALAM B S (19 6) tion London
- 13 DARLING S T (1926) Mosquito species control of malaria
1 er *J Trop Med* VI 3 pp 16
178
- 14 DEADERCK W H (1909) A practical study of malaria Philadelphia
and London
- 15 DUNHAM C C (1928) Methods of reducing the cost of anti mos
quito measures in the control of malaria
M I Surg XLII 1 pp 33-48

* R4LB = *Review of Applied Entomology* Series B.
TDB = *Tropical Diseases Bulletin*

- 16 FOWLER, C E P (1926) Malaria in Palestine A summary of anti malarial work carried out with the 54th Division in the Mulebbis Area April--June 1918 *Jl R A M C XVI* 4 pp 241-248
- 17 GOSIO, B (1926) Five lectures on malaria control Rome
- 18 HARDENBURG, W E (1922) Mosquito eradication New York and London
Les moustiques Ed 2 Brussels
- 19 HEOH, E (1921) Malaria in India London
- 20 HENIR, P (1927) Malaria etiology, pathology, diagnosis, prophylaxis and treatment London
- 21 HENSON, G E (1913) The prevention of malaria with a Division in the Field *Jl R A M C, VII* 4 pp 259-265
- 22 HOLDEN, C W (1924) How the cotton belt railway cut malaria rate 97 per cent in 9 years *Railway Engineering and Maintenance XXII*, 10, pp 382-390 Abst in *U S A Pub Hlth Rpts*, XLII, 13 pp. 878-879
- 23 HOVENBURG, H W VAN (1926) The mosquitoes of North and Central America and the West Indies, I, pp 354-444 Washington
- 24 HOWARD, L O, DYAR, H G, and KNAB, F (1912) Malarial fevers Ed 3 Govt Press, Calcutta
- 25 JAMES, S. P (1909) Malaria at home and abroad London
- 26 JAMES S P (1920) The epidemiology and control of malaria in Palestine Chicago
- 27 KLIJLER I J (1930) Malaria its investigation and control Calcutta
- 28 KNOWLES, R, and SENIOR WHITE, R A (1927) *Traité du paludisme Ed 2 Paris*
- 29 LAVERAN, A (1907) *Paludisme Paris*
- 30 MARCHOUX, E (1926) Some aspects of malaria control through mosquito eradication *U S A Pub Hlth Rpts*, XXXIV, 5 pp 167-183
- 31 METZ, C W (1919) The organization of minor anti anopheline field work *Ceyl Jl Sci*, D II, 1, pp 31-36
- 32 NICHOLLS, L (1927) Can we get better Anopheles control and more malaria control at less cost? *Proc Inter Conf Hlth Prob Trop Amer* (United Fruit Co Pbn) pp 157-164
- 33 PRINCE, J A LE (1924) Mosquito control in Panama New York and London
- 34 PRINCE, J A LE and ORENSTEIN, A J (1916) Some experiments with fish and chemicals against mosquito larvae *Dept Ind and Ag Hand Suriname RAEB XIII*, p 6
- 35 REYNE, A (1924)

- 36 ROSE P (1909) Mosquito brigades and how to organize them London
- 37 ROSE P (1911) The prevention of malaria Ed ° London
- 38 ROSE P (1906) Malaria control in Ceylon plantations
Jl Trop Med Hyg XXIV pp 150-155
- 39 SEGUY E (1904) Les moustiques de l'Afrique Mineure de l'Egypte et de la Syrie Paris
- 40 SERGEANT ED and SERGEANT ET (1905) Survey of anti malaria methods *Arch Inst Past Algér* III pp 359-388
- 41 SERGEANT ED and SERGEANT ET (1908) Vingt cinq années d'étude et de prophylaxie du paludisme en Algérie *Arch Inst Past Algér* VI ° and 3 pp 117-134
- 42 SIMPSON W J (1909) Principles of hygiene as applied to tropical and sub tropical climates London
- 43 VISBECK and LACAZE H (1904) Le paludisme aux armées et la lutte antipaludique Paris
- 43a WARBLE R A and BUCKLE P (1903) The principles of insect control London
- 44 WATSON M (1915) Rural sanitation in the tropics London
- 45 WATSON M (1911) The prevention of malaria in the Federated Malay States Ed ° London
- 46 WILSON H C (1912) Some notes on larvices and natural enemies of mosquitoes in Southern India G C Press Simla
- 47 WILSON H C (1914) A note on the treatment of swamps etc., with a view to the destruction of mosquito larva *Ind Jl Med Res* I 4 pp 691-701

B SPECIAL SUBJECTS

Bats

- 48 CAMPBELL, C A P (1913) The eradication of mosquitoes by the cultivation of bats *Monthly Bull Agric Intell and Plant Dis* Rome IV, pp 1175-1181 *RAEB* I pp 16-177
- 49 HOWARD L O (1900) Mosquitoes and bats *U S A Pub Illus Rpts* XXXV 31 pp 189-195
- 50 MIYAKOTO S (1930) On the destruction of mosquitoes by *P. p. etellus armatus* in Tahiti *Jl Med Ass Formosa* No 301 p 19
TDB XXVII p 903
- 51 NELSON E W (1906) Bats in relation to the production of guano and the destruction of insects
U S A Dept Agric Bull No 1395
- 52 STORER T I (1906) Bats bat towers and mosquitoes *Jl Mammal* VII pp 8-90
Summ Bull Hyg II 7 p 593

- 53 VITTORIO, V (1926) Auxiliary measures in the campaign against malaria *Ann Med Lav Colon*, II, pp 273-292 *Summ Bull Hyg*, II, 7, p 589
- 54 VLASSOV, J (1927) On the question of mutual relation between bats and mosquitoes *Rev Microb and Epidem*, VI, 2, Eng Summ., pp 260-261

Biological Control.

- 55 BARBER, M A (1927) The food of *Anopheles* larva.—Food organisms in pure culture *U S A Pub Hlth Rpts*, XLII, 22 pp 1494-1510
- 56 BARBER M A and KOMR W H W (1933) The hydrogen ion concentration of the soil and natural waters in relation to *Anopheles* production *U S A Pub Hlth Bull* No 137, pp 11-12
- 57 BARBER M A, KOMR W H W, and HAYNE T B (1926) Malaria in the prairie rice regions of Louisiana and Arkansas *U S A Pub Hlth Rpts*, XII, 45, pp 2527-2549
- 58 BATSCHINSKY, P F (1936) The biology of larva of *Anopheles* and *Culex*. Attempts to infect the larva with spores of mould fungi *Hyg Epidem Leningrad* pp 33-44 *RAEB*, XV, p 153
- 59 BEATTIE, M V F (1930) Physico-chemical factors in relation to mosquito prevalence in ponds *Jl Ecology* XVIII, 1, pp 67-89
- 60 BUXTON P A (1934) Applied entomology of Palestine *Bull Ent Res*, XIV, 3, pp 289-340
- 61 COGGESHALL, L T (1926) Relationship of plankton to *Anopheles* larva *Amer Jl Hyg*, VI 4 pp 556-559
- 62 GATER B A R (1930) The asthenobiosis theory and larva surveys *Malayan Med Jl* V, 4 pp 126-128
- 63 HAMLYN HARRIS R (1928) The relation of certain algae to breeding places of mosquitoes in Queensland *Bull Ent Res*, XVIII, 4 pp 377-389
- 64 HAMLYN HARRIS, R (1930) The consideration of certain factors as potentialities in mosquito control in Australia *Proc Roy Soc Queensland*, XLII, 10, pp 86-105
- 65 HANCOCK G L R (1930) Some records of Uganda mosquitoes and the ecological associations of their larva *Bull Soc Roy Ent Egypte* 1, pp 38-56 *RAEB*, XVIII, p 241

- 66 HINMAN F H (1939) A study of the food of mosquito larvae (Culexidae) *Amer Jl Hyg*, XII, 1, pp 239-250 (5 pp refs)
- 67 HOWLAND L J (1930) The nutrition of mosquito larvae with special reference to their algal food *Bull Ent Res*, XXI, 4 pp 431-440
- 68 HOWLAND L J (1930) Bionomical investigation of English mosquito larvae with special reference to their algal food *Jl Ecology*, XVIII, 1, pp 67-80
- 69 IYENGAR M C T (1930) Dissolved oxygen in relation to *Anopheles* breeding *Ind Jl Med Res* XVII, 4 pp 1171-1183
- 70 LAMBORN W A (1922) Some problems of the breeding places of the Anophelines of Malaya *Bull Ent Res* XIII, 1, pp 1-23
- 71 LAMBORN W A (1922) The bionomics of some Malayan Anophelines *Ibid* XIII, 2 pp 129-149
- 72 MACGREGOR M E (1931) The influence of the hydrogen ion concentration in the development of mosquito larvae *Parasitology* XIII 4 pp 348-351
- 73 MACGREGOR M F (1924) The significance of the pH in the development of mosquito larvae *Ibid*, XXI, 1-2, pp 132-157
- 74 MEYER C W (1919) Observations on the food of *Anopheles* larvae *U S A Pub Hlth Rpts* XXXIV 32, pp 1783-1791
- 75 NIELSEN J I (1936) Preferential breeding conditions of *Anopheles* in the Philippine Islands *Phil Jl Sci* XLII 1 pp 59-63
- 76 NIKITINSKI A (1932) Life and physico-chemical régime of the larva and conditions of development of *Anopheles maculipennis* Russ *Jl Trop Med* pp 13-15 *RIEB* XIV, p 201
- 77 PATTERSON, N F (1923) The effects of change of hydrogen ion concentration on Culexine mosquito larvae *S Afric Jl Sci*, XXII pp 311-317
- 78 PERDY W C (1920) Biological investigation of Californian rice fields relative to mosquito breeding *U S A Pub Hlth Rpts* XXXI 44 pp 2556-2570
- 79 PERDY W C (1925) Biological investigation of Californian rice fields and attendant waters with reference to mosquito breeding *U S A Pub Hlth Bull*, No 14*

- 80 RAMSAY, G C (1930) The factors which determine the varying degrees of malarial incidence in Assam Tea Estates, and the fundamental principles governing mosquito control of malaria in Assam *Trans Roy Soc Trop Med and Hyg*, XLIII, 5 pp 511-518
- 81 RAMSAY, G C (1930) The malaria problem of Assam *Jl Trop Med and Hyg*, XLVIII, 23, pp 352-357
- 82 RAMSAY, G C, and FRASER, G (1930) Notes on silt and shade in the control of malaria in Assam *Ross Inst Publ* London
- 83 RUDOLFS, W (1925) The food of mosquito larvæ *Procs 12th Ann Mtg N Jers Mosq Ext Ass*, pp 25-33
- 84 RUDOLFS W (1928) Contribution to the causes of mosquito breeding in specific places *Procs 15th Mtg N Jers Mosq Ext Ass*, pp 66-76
- 85 RUDOLFS W, and LACKEY, J B (1929) The composition of water and mosquito breeding *Amer Jl Hyg*, IX, 1, pp 160-180
- 86 SENIOR WHITE, R A (1926) Physical factors in mosquito ecology *Bull Ent Res*, XVI, 3, pp 187-248
- 87 SENIOR WHITE, R A (1928) Algae and the food of Anopheles larvæ *Ind Jl Med Res*, XV, 4 pp 969-988
- 88 SENIOR WHITE R A (1928) On the relationship existing between carbonates and pH and conductivity in natural waters *Ibid*, pp 989-996
- 89 SENIOR WHITE, R A (1928) Physical factors in mosquito ecology Part II *Ibid*, XVI, 1, pp 11-30
- 90 SENIOR WHITE, R A (1929) Progress towards realization of biological control of mosquito breeding *Trans 7th Cong F E 4 T M*, 1927, II, pp 718-722
- 91 SHARMA H N (1921) A preliminary note on the action of acids and alkalis on the development of Culicid eggs and larvæ *Procs 4th Entom Mtg, Pusa*, pp 199-204
- 92 SMORODINZEW, I A, and ADOWA, A N (1930) The role of calcium in the ecology of the larvæ of Anopheles *Bull Soc Path Exot*, XLIII, 1, pp 34-38
- 93 SMORODINZEW, I A, *et al* (1928) Les variations de la réaction réelle de réservoirs d'eau artificiels dans les recherches sur la biologie des Anophèles *C P Soc Biol*, XCIX, 36, pp 1763-1763 *RAEB*, XVII, p 65

- 93a STRICKLAND C (1916) Malaria in the Malay Peninsula *Ind J Med Res*, IV 3 pp 256-267
(Dangers of jungle clearing)
- 94 WATSON M (1911) The prevention of malaria in the Federated Malay States Liv School Med Pbn
- 95 WILLIAMSON K B (1926) Investigations dealing with rice fields *Malay Med J* I, 4 pp 21-27
- 96 WILLIAMSON K B (1928) Mosquito breeding and malaria in relation to the nitrogen cycle *Bull Ent Res*, XVIII, 4 pp 433-439
- 97 WILLIAMSON K B (1929) Chemical factors in relation to Anopheline breeding *Trans 7th Cong F E A T M* 1927 II pp 723-735

Copper Sulphate

- 98 HARGREAVES E (1923) Entomological notes from Taranto *Bull. Ent Res* XIV 2 pp 213-219
- 99 KLIGLER I J (1925) Methods of mosquito control in Palestine *Trans 1st Inter Cong Mal Rome*, pp 149-153
- 100 MILLER D (1920) Mosquito investigation in the North Auckland Peninsula of New Zealand *New Zealand Dept Pub Hlth Pbn* No 3 p 14
- 101 ROSS F and EDIX E S (1911) Some experiments on larvicides. *Ann Trop Med Paras* V 3 pp 385-390
SENIOR WHITE R A (1928) See No 87
- 102 SMITH J B (1904) Report of the New Jersey State Agricultural Experiment Station upon the mosquitoes Trenton N J
- 103 WHIFFLE G C (1914) The microscopy of drinking water, pp 251-260 New York

Cresol (as larvicide).

- 104 BOYD J E M (1925) Some further notes on cresol as a larvicide *Jl R A M C* XLIV 4 pp 285-287
- 105 DELMEGE J A (1919) Some practical notes on the prevention of mosquito breeding *Jl Trop Med Hyg* XXII 19 pp 181-184
- 106 HAROLD C H H (1926) Studies in mosquito bionomics *Jl R A M C* XLVII 2 pp 81-94 and 3 pp 180-187
- 107 HENDLEY M (1916) Report on malaria in the Punjab during the year 1915 Mal Bureau Lahore
HOLDEN C W (1924) See No 22
- 108 KLIGLER I J (1925) Methods of mosquito control in Palestine *Trans 1st Inter Cong Mal Rome*, pp 149-153

- 109 KLIGLER, I J (1926) - The control of malaria in Palestine by anti-anopheline measures *Jl Pres Med*, I, pp 149-183
- 110 KLIGLER, I J, *et al.* (1923) Report of the Malaria Research Unit, Palestine
- 111 KLIGLER, I J, and SHAPIRO, J M. (1925) Organization of areas for mosquito control in Palestine *Harcfuah*, I 4-5, pp 180-188 *RAEB*, XIV, p 57
- 112 KLIGLER, I J, SHAPIRO, J M, and WEITZMAN, L (1924). Malaria in rural settlements in Palestine *Jl Hyg*, XXIII, p 280
- 113 MARSHALL, J F (1925) . The destruction of mosquito larvae in salt and brackish water *Hysling I Moeg. Cont Circ.* No. 4.
- 114 MARSHALL, J F. (1925) -- Some practical notes on mosquito control *Ibid* No 16
- 115 MAYNE, J. F, and JACKSON, W R (1920) Larvicides *Jl R A M C*, XXXIV, 2 pp 112-120
- 116 STRICKLAND C, and ROY, D N (1926) Cresol, saponified, as a larvicide, with a note on the cresols by STEWART, A D, *Jl R A. M C*, XLVII, 3, pp 183-193

Deferrent Plants and Trees.

- 117 ADIE, J R (1904) *Lemna minor* as a preventive against mosquitoes. *Ind Med Ger*, XXXIX, 6, pp 207-208
- 118 ANCONA, H L (1930) *Lemnaceae* and mosquito larvae *Ann. Inst Biol Univ Mexico*, I, 1, pp 33-37 *RAEB*, XVIII, p 212
- 119 AFFELBECK, V (1925) Recherches et observation sur les arthropodes pathogènes de l'homme et des animaux. *Ed Insap Mex San Pub Sarajevo*, No 17 *RAEB*, XVI, pp 178-179
(*Lemna*, *Chara* *Utricularia*, etc.)
120. BARBER, M A (1924) . The effects of *Chara robbinsii* on mosquito larvae *U. S A Pub Hlth Rpts*, XXXIX, 13, pp 611-615
- 121 BARBER, M A, and HAYNE, T B (1925). Water hyacinth and the breeding of *Anopheles* *Ibid*, XL, 47, pp 2557-2562
- 122 BENTLEY, C A (1910) The natural history of Bombay malaria *Jl Dom Nat. Hist Soc*, XX, pp 392-422 (*Lemna*, *Azolla*, *Wolffia*)
- 123 BLOW, T. B. (1927) Observations on the alleged larvicidal qualities of the Charophyta. *Proc Linn Soc*, Sess 139, pp 46-47.
- 124 BOSTIN, S. L. (1917) .. Fucalvptus trees and malaria *Agric Jl India, Agric Res Inst*, Pusa, XII, 3, p 495.

- 125 BOYD M F (1930) Malaria control New Orl Med and Surg JI LXXXIII 3 p 194 (Alfalfa)
- 126 BROCHER F (1927) Capture of Anopheles larvae by Utricularia Ann Pa as Hum and Comp V 1 pp 46-47
- 127 BRECHT E (1925) Capture des larves des Culicidées par les plantes du genre Utricularia Ibid III 4 pp 403-411
- 128 BUCHÉ E DE (1929) The incompatibility of occurrence to either of Anopheles larvae and Lemna Med Pa s Cal II 3 pp 262-270
- 129 BURNOT E W (1927) Effects on mosquito larvae of a Queensland Vella Pro s Roy Soc Queensland XXXVIII pp 59-61
- 130 CARRALLERO A (1919) La Chara fatida A Br y las larvas de Stigmaria Culex y Anopheles Bol R Soc Espan Hist Nat Madrid XIX 8 pp 449-455 RAEB VIII p 61
- 131 CARTER H G (1916) Aquatic plants supposed to possess power to destroy mosquito larvae Ind Med Rec XXVI 1 p 15
- 132 CASTILLO V (1926) Preliminary studies on the insecticidal properties of three species of Drosera in the Philippines Phil Agric XV pp 257-275 RAEB XV p 17
- 133 FEDERCI E (1927) The toxic action of species of Chara on Culicid larvae Red a XVI 1 2 pp 17 23 RAEB XVI p 111
- 134 FRANÇA C (1923) Recherches sur les plantes carnivores II Utricularia vulgaris Bol Soc Boter I 1 pp 11-37 RAEB, XII p 178
- 135 FRANÇA C (1924) L'emploi des plantes dans le combat des moustiques Rev Med Angola III 4 pp 419-4 P4EB VII p 53 (Recommend use of Azolla)
- 136 GUNGA TULSI (1903) Mosquito plants Ind Lancet XVII p 603
- 137 HAMLYN HARRIS R (1927) Notes on the breeding places of two mosquitoes in Queensland Publ Ent P s XVII 4 pp 411-414 (Vella phaula L.)
- 138 HAMLYN HARRIS R (1928) S s No 63 (Characeae Cladophora holostea)
- 139 D'HERELLE F (1921) Immunity to natural infectious diseases pp 18-183 Baltimore (Clover and malaria)

- 139 KELSALL, A, *et al* (1926) *Derris* as an insecticide 16th Ann Rept. Ent Soc Ontario, 1925, pp 24-40
RAEB, XV, p 17
- KLIGLER I J (1925) . See No 108
(*Chara*)
- 140 LANGERON, M (1925) Biologie et ecologie des Utriculaires dans leur rapports avec la prophylaxie du paludisme Ann Paras Hum and Comp, III, 4, pp 412-425
- 141 MACGREGOR, M E (1920) The possible use of *Utricularia filiculoides* as a deterrent to Anopheline breeding JI R A M C, XXXIV, 4 pp 370-372
- 142 MACGREGOR, M E (1924) Tests with *Chara fatida* and *C hispida* on the development of mosquito larva Parasitology, XVI, 4, pp 383-387
- 143 MATHESON, R. (1928) The effect of *Chara fragilis* on mosquito development Procs 15th Mtg A Jerr Mosq Exl iss, p 77
- 144 MATHESON R (1930) The utilization of aquatic plants as aids in mosquito control Amer Nat, LXIV, No 690, pp 86-86 (3 pp refs)
(General review)
- 145 MATHESON, R., and HINMAN, E H (1928) *Chara fragilis* and mosquito development Amer JI Hyg, VIII, 2 pp 279-292
- 146 MATHESON, R., and HINMAN E H (1929) Further studies on *Chara* spp and other aquatic plants in relation to mosquito breeding Amer JI Trop Med IX 4 pp 249-266
(*Chara*, *Phyllotria*, *Spirogyra* *Oedogonium*, *Zygnema*)
- 147 MATHESON, R., and HINMAN F H (1930) A seasonal study of the plankton of a spring fed *Chara* pool versus that of a temporary to semi permanent woodland pool in relation to mosquito breeding Amer JI Hyg, XI, 1, pp 174-188
- 148 MAYNE, B (1930) The effects of coumarin on the life of the mosquito and the malarial parasite Ind JI Med Res, XVII, 3, pp 963-969
(Clover and malaria)
- 149 MUEHLENS, P (1924) Preliminary report on studies in the Argentine Territories of Chaco and Formosa and in Paraguay 4th Dept Hyg, XXX, 2 pp 139-146 RAEB. XV, p 149
(*Utricularia*)
- 150 SANITAS (1903) . Sunflowers and malaria Ind Lancet, XVI, p 1109

51. SIVARAMIAH M N (1916) . Aquatic plants supposed to possess the power to destroy mosquito larvæ *Ind Med Rec*, XXVI, 3, p 60
52. VALERIO, B G, and JONGH, I R DE (1909) Observations on the destruction of the larvæ and pupæ of Culicidæ *Cent f Balt*, XIX, 4 Abst in *Jl R A M C*, XII, pp 702-704 (*Lemna*, *Utricularia*)
53. VIOSCA P (1925) A biological study of the mosquitoes of New Orleans and south eastern Louisiana *Procs 12th Ann Mtg N Jers Mosq Ext Ass*, pp 34-50 (Advocates use of water hyacinth)
54. WILCOCKS W (1927) Why is cultivated Egypt immune from malaria? Nile Mission Press Cairo (Clover and malaria)

Drainage

55. BAILE, J F *et al* (1919) The use of dynamite in malaria control operations *U S A Pub Hlth Bull* No 104, pp 137-162 (For blasting ditches)
56. BENNETT, H L (1930) Anti malarial drainage *Kenya and E Afric Med Jl* VII, 7 pp 190-198
57. BENTLEY, C A (1914) A proposal for experiment in subsoil drainage (MS) Office of D P H, Bengal
58. BENTLEY C A (1925) Malaria and agriculture in Bengal How to reduce malaria in Bengal by irrigation Calcutta
59. BRAHMACHARI B B (19'3) Flood flushing and malaria in the Kumar khali Municipality *Calc Med Jl*, XVIII, 2, pp 342-347
60. BUCK F (1914) Report on the control and utilization of rivers and drainage for the fertilization of land and mitigation of malaria *Procs 3rd 4th Ind San Conf*, I, pp 27-29, 32-35, 51-52 and II, pp 1-26
- 60a. CARTER H I (1915) Impounded waters *U S A Pub Hlth Rpts* XXX, 1, pp 15-33
61. CARTER, H I (1924) The effect of the variation of level of impounded water on the control of *Anopheles* production *South Med Jl*, XVII 8, pp 575-577
62. CARTER H I, PRINCE, J A I R, and GRIFFITHS T, H D (1916) Impounded water *U S A Pub Hlth Bull* No 79

- 163 CHAMBERLAIN, W P (1927) Sanitation of Panama Canal Zone Per
manent drainage and mosquitoes.
Mal Surg, LX, 4, pp 397-422
- 164 CHAMBERLAIN, W P and CURRY, D P (1926-30) Mosquito control *Rpts Hlth Dpt
Panama Canal*, 1925 1926, 19 7
1928
- 165 DAVIS A P, and WILSON, H M (1919) Irrigation engineering Ed 7 New York
and London
- 166 DINE, D L VAN (1922) Impounding water in a bayou to control
breeding of malaria mosquitoes
U S A Dpt Agric Bull No 1098
- 167 ELLIOTT, C G (1912) Engineering for land drainage Ed 2
London
- 168 EVANS, F D (1916) Engineering operations for the prevention
of malaria *Procs Inst Civil Engineers
CC, Session 1914-15, Part II*, pp 1-62
- 169 GAGE, E H (1925) ~ ~ ~ ~ ~
- 2219
- 170 GILL, C A (1917) Report on malaria in Amritsar Govt
Press, Lahore
(Water logging and irrigation)
- 171 GILL, C A (1927) The relationship of canal irrigation and
malaria *Agric JI Ind*, XXII, 3,
pp 168-172 Reprinted in *Rec Mal
Surg Ind*, I, 3, pp 417-421
- 172 GORMAN, A F, STROMQUIST W G, and BISHOP, G H (1919) Vertical drainage *U S A Pub Hlth
Bull* No 104 p 64
- 173 GRASSI B (1925) Drainage, artificial lakes and malaria
Rome
- 174 GRIFFIN, F C (1927) Meenglas anti malarial scheme—Inspection
note Govt Press Calcutta
(Subsoil drainage)
- 175 GURRAY, H A (1917) Surface drainage Ed 2 Calcutta
- 176 HENRY, A K (1919) Destruction of mosquito larvae in streams
Lancet, CXCVI, No 4995 pp 908-909
(Canalization)
- 177 HOME, H (1926) The engineer and the prevention of
malaria London
- 178 HUNTER, P S (1926) A description of the sanitary and anti
malarial work carried out at Gunong
Pulai waterworks, South Johore
(Subsoil drainage of ravine streams,
etc)
- 179 IRVINE, J (1919) Memorandum on the plan and section of
the type ravine drainage (MS) Office
of D P H, Bengal

- 180 JACKSON, L F (1925) Mosquito control in New Jersey during the breeding season of 1924 *Amer JI Pub Hlth*, **XV**, 2 pp 131-133 (Ditch-cutting and ditch cleaning machines)
- 181 KING, F H (1922) Irrigation and drainage Fd 8 New York
- 182 LEEZE T A (1924) Regulation of rivers without embankments as applied in the training works at the head waters of the Rangoon River, Burma London
- METZ C W (1919) See No 31 (Drainage of swamps and seepage outcrops with lines of contour drains)
- 183 MUSHHEAD W A (1912) Practical tropical sanitation Ed 2 London
- 183a PRINCE J A LE (1915) Impounded waters *U S A Pub Hlth Rpts*, **XXX** 7, pp 474-481
- 184 PRINCE J A LE (1915) Malaria control Drainage as an anti-malarial measure *U S A Pub Hlth Rpts* **XXX**, 8 pp 536-545 (Open and subsoil drainage)
- 185 PRINCE J A LE (1927) Mosquito control in relation to impounded water supply *Jl Amer Waterworks Ass* **XVII** 1 pp 31-36 Abat in *U S A Pub Hlth Rpts* **XLII** 25, pp 1713-1714
- 186 PUBLIC WORKS DEPT F M S (1922) Specification for construction of under drainage Reprinted in Knowles and Senior White 1927 (see No 28)
- 187 REELEY F A (1928) A history of salt marsh ditching and ditch cleaning machinery *Procs 15th Mtg N Jers Mosq Ext Ass* p 24
- 188 SANCAR S L (1927) Flood and flush schemes ancient and modern with reference to the ancient city of Gour *Ind Med Gaz*, **LXVI** 4 p 188
- 189 SCOTT J (1920) Draining and embanking Fd 2 London
- 190 SIMMS H (1909) Sub drainage as applied to the anti malaria campaign on the Isthmus of Panama *Ann Trop Med and Paras*, **II**, 4 pp 291-293
- 191 SCH S N (1929) Malaria problem in Bengal Govt Press Calcutta
- 192 WALKER J H C (1930) Anti malaria measures in Malaya *Jl R A M C* **LIV** 2, pp 87-106
- 193 WHEELER W H (1888) The drainage of fens and lowlands London

- 163 CHAMBERLAIN, W P (1927) Sanitation of Panama Canal Zone Per
manent drainage and mosquitoes
Mal Surg, LX, 4, pp 397-422
- 164 CHAMBERLAIN, W P, and CURRY, D P (1926-30) Mosquito control *Rpts Hlth Dpt
Panama Canal*, 1925, 1926, 1927,
1928
- 165 DAVIS, A P, and WILSON, H M (1919) Irrigation engineering. Ed 7 New York
and London
- 166 DINE D L VAN (1922) Impounding water in a bayou to control
breeding of malaria mosquitoes
U S A Dpt Agric Bull No 1098
- 167 ELLIOTT, C G (1912) Engineering for land drainage Ed 2,
London
- 168 EVANS, F D (1916) Engineering operations for the prevention
of malaria *Procs Inst Civil Engineers*,
CC, Session 1914-15, Part II pp 1-62
- 169 GAGE E H (1925) Studies of impounded waters in relation to
malaria *U S A Pub Hlth Rpts*, XL,
28, pp 1357-1375, and 42 pp 2211-
2219
- 170 GILL, C A (1917) Report on malaria in Amritsar Govt
Press, Lahore
(Water logging and irrigation)
- 171 GILL C A (1927) The relationship of canal irrigation and
malaria *Agric Jl Ind*, XXII, 3,
pp 168-172 Reprinted in *Rec Mal
Surg Ind* I, 3, pp 417-421
- 172 GORMAN, A F, STROMQUIST, W G, and BISHOP, G H (1919) Vertical drainage *U S A Pub Hlth
Bull* No 104, p 64
- 173 GRASSI B (1925) .. Drainage, artificial lakes and malaria.
Rome
- 174 GRIFFIN, F C (1927) Meenglas anti malarial scheme—Inspection
note Govt Press, Calcutta
(Subsoil drainage)
- 175 GUBBAY, H A (1917) Surface drainage Ed 2 Calcutta
- 176 HENRY, A K (1919) Destruction of mosquito larvae in streams
Lancet, CXCVI No 1995, pp 908-909
(Canalization)
- 177 HOME, H (1926) The engineer and the prevention of
malaria London
- 178 HUNTER P S (1926) . A description of the sanitary and anti
malarial work carried out at Gunong
Pulai waterworks South Johore
(Subsoil drainage of ravine streames,
etc)
- 179 IRVINE, J (1919) .. Memorandum on the plan and section of
the typo ravine drainage (MS) Office
of D P H, Bengal

- 180 JACKSON L F (1925) Mosquito control in New Jersey during the breeding season of 1924 *Amer JI Pub Hlth* **XX**, 2 pp 131-133 (Ditch-cutting and ditch cleaning machines)
- 181 KIRBY F H (1922) Irrigation and drainage Ed 8 New York
- 182 LEYTE T A (1924) Regulation of rivers without embankments as applied in the training works at the head waters of the Pangoon River Burma London
- METZ C W (1919) See No 31 (Drainage of swamps and seepage outcrops with use of contour drains)
- 183 MUIRHEAD W A (1922) Practical tropical sanitation Ed 2 London
- 183a PRINCE J A LE (1915) Impounded waters *U S A Pub Hlth Rpts* **XXX** 7 pp 473-491
- 184 PRINCE J A LE (1915) Malaria control Drainage as an anti-malarial measure *U S A Pub Hlth Rpts* **XXX** 8 pp 536-545 (Open and subsoil drainage)
- 185 PRINCE J A LE (1927) Mosquito control in relation to impounded water supply *Jl Amer Waterworks Ass* **XXII** 1 pp 31-36 Abat in *U S A Pub Hlth Rpts* **XLII** 2, pp 1713-1714
- 186 PUBLIC WORKS DEPT F M S (1927) Specification for construction of underdrainage Reprinted in Knowles and Senior White 1927 (see No 28)
- 187 REILEY F A (1928) A history of salt marsh ditching and ditch cleaning machinery *Procs 15th Mtg N Jers Mosq Ext Ass* p 24
- 188 SANCAR S L (1927) Flood and flush schemes ancient and modern with reference to the ancient city of Gour *Ind Med Gaz* **LXII** 4 p 183
- 189 SLOTT J (1920) Draining and embanking Fl 2 London
- 190 SIMMS H (1929) Sub drainage as applied to the anti malaria campaign on the Isthmus of Panama *Ann Trop Med and Paras* **II** 4 pp 291-295
- 191 SUE S N (1929) Malaria problem in Bengal Govt Press Calcutta
- 192 WALKER J H C (1930) Anti malaria measures in Malaya *Jl R A M C LIV* 2 pp 87-100
- 193 WHEELER W H (1898) The drainage of fens and lowlands London

- 194 WILLIAMS L I (1905) Effects of pond control on malaria prevalence *U S A Pub Hlth Bull* No 156 pp 56-64
(Fluctuation of levels)
- 195 WRIGHTSON W D (1915) Mosquito eradication and prevention *Am r Jl Trop Dis and Prev Med*, II 12 pp 738-752
(Open and subsoil drainage)
(See also Nos 1 to 47)
- Fish**
- 196 AITKEN E H (1901) Notes on *Anopheles* or malaria mosquito *Jl Bom Nat Hist Soc* XIII, pp 691-695
- 197 CHATTERJEE G C (1909) Role which fish play in the control of malaria *Ind Med Rec* May
- 198 CHAUDHURI B L (1909) Mosquito larvae eating propensities of the Genus *Haplocheilichthys* *Proc Asiat Soc Bengal New Series* V p 36
- 199 CHAUDHURI B L (1912) Fish and mosquito larvae *Calc Med Jl* VI 12 pp 457-470
- 200 DAY F H (1889) Fishes Fauna of British India I and II
- 201 DERANIYAGALA P E P (1930) Some probable Ceylon larvivores *Ceyl Jl Sci D* II 4 pp 191-201
- 202 DONALD C H (1923) List of larvicidal fish found in the Punjab with their vernacular names (MS) Office of Health Officer Delhi
- 203 FRY A B (1912) Indigenous fish and mosquito larvae *Paludism* No 5 pp 71-74
- 204 GRAHAM J D (1913) Note on mosquito larvae destroying fish in the United Provinces Allahabad
- 205 HILDEBRAND S G (1921 1925) A study of the top minnow *Gambusia holbrooki* in relation to malaria *U S A Pub Hlth Bull* Nos 114 and 153
- 206 HORA S L (1927) The use of fishes for the control of malaria *Ind Med Gaz* LXII 4 pp 187-188
- 207 HOWELL G C L (1913) Fish in relation to the malaria problem *1st Rept Dept Fish Punjab* Part IV
- 208 INTERNATIONAL HEALTH BOARD ROCKEFELLER FOUNDATION (1924) The use of fish for mosquito control New York (1917 refs)
- 209 MURPHY W O S (1912) Larvae eating fish of Sind (MS) Office of D P H Bombay
- 210 NICHOLLS L (1910) Mosquitoes and their natural enemies *Ann Rept S Lucia W Indies Comm Bull Ent Res* I pp 213-218

FISH, FUMIGANTS

- 211 O'DONNELL S P. (1912) Reports on the suitability of 'mullions' as destroyers of mosquito larvae in India. Reprinted in *Pec Mal Surv Ind*, 1, 2, pp 137-139 (1930)
- 212 OGBORN W (1906) Fish mosquitoes and lime *Ind Med Gaz* XLI, 12, pp 498-499
- 213 STILLA M (1925) A new means of combating Anophiles in Italy. An account of the acclimatization and progress of *Gambusia*. *Trans 1st Inter Cong Mal Rome*, pp 105-114
- 214 SWEET, R B, and CHURCH, B L (1912) Indian fish of proved utility as mosquito destroyers. *Ind Museum Calcutta*
- 215 SOUTHWELL, T (1930) Fish and mosquito larvae in Bengal, Bihar and Orissa India. *Ann Trop Med and Paras* XIV, 2, pp 181-186
- 216 WILSON H C (1917) Some notes on larvicides and natural enemies of mosquitoes in southern India. *Proc 3rd Mtg Cent Mal Comm, Madras 1912*
- 217 WILSON H C (1918) A note on the treatment of swamps, etc. *Malaya Fisheries Bur Bull* No 11, pp 161-172

Fumigants

- 218 ALEXANDER D (1919) Trial as fumigant *The Medical Officer*, July. Abet in *Ind Med Gaz*, LIII 12, pp 465-466
- 219 BAISFOUR A (1913) A year's anti-malarial work at Khartoum. *Jl Trop Med Hyg* XVI 15 pp 225-232 (Cresol)
- 220 BARRETO I DE B et al (1926) Work in dwellings against adult Anophelines. *Sc Med* IV 12 pp 693-707 *RAFB* VI p 67 (Sulphur dioxide)
- 221 BENYAMINSON F S and NABOKOV V A (1930) The different products of pyrethrum in the campaign against hibernating Anophelines. *Russ Jl Trop Med* VII, pp 476-486 (Summary in French)
- 222 BROMLEY S W (1938) Killing mosquitoes with Cyanogas calcium cyanide. *Jl Econ Entom* XIX 6 p 871
- 223 CLYDE D (1931) Report on the control of malaria during the Banda Canal construction (1920-1931). *Pec Mal Surv Ind*, II 1, pp 49-110 (Sulphur with waste tobacco, pyrethrum cresol)
- 224 DYK and JUMENTIE (1911) Artemisia (*Sinifonica*) used by trappists as fumigant in Japan. *Le Caducée*, No 7

- 224 FERMI, C, and IUMBAO, S (1900) Befreiung einer Stadt von der Mücken
Cent f Balt, I Abt, XXVIII
pp 179-186
(Chlorine from sulphuric acid and
chloride of lime)
- 225 FRYER, J C F, *et al* (1928) English grown pyrethrum as an insecticide
Ann App Biol, XV, pp 423-445
- 226 GILES, G M (1902) A handbook of the gnats or mosquitoes
Ed 2, p 221 London
(Sulphur cones)
- 227 GNADINGER C B, and CORLI
C S (1929) Studies on pyrethrum flowers I The
quantitative determination of the active
principles *Jl Amer Chem Soc, LI*,
pp 3054-3064
- 228 GUITERAS, G M (1909) Mercuric chloride as an insecticide
U S A Pub Hlth Rpts, XXIV, 50,
pp 1859-1861
- 229 ISTHMIAN CANAL COMMISSION (1912) Destruction of mosquitoes by pyrethrum
Rept Dpt San Istn Can Comm
- 230 LEGROUX, R (1917) Presentation du material de prophylaxie
antipaludique destiné à l'armée d'Orient
Bull Soc Path Exot, X, 6, pp 421-427
(Cresol vaporiser)
- 231 McDONNELL, C C, *et al* (1930) Relative insecticidal value of commercial
grades of pyrethrum *U S 1 Dpt
Agric Tech Bull No 198*
- 232 McDONNELL, C C, ROARK,
R C, and KEEVAN, G L
(1920) Insect powder *U S A Dpt Agric Bull
No 824* (299 refs)
- 233 NABOKOV, V A (1929) Fumigation with sulphur dioxide from
sodium sulphite *Russ Jl Trop Med,
VII, 5*, pp 349-354 *R4EB, XVIII*,
p 38
- 234 NABOKOV, V A (1929) New combined apparatus for disinfection
and destruction of insects *Ibid, VII*,
8 pp 530-536 *R4EB, XVIII*, p 80
(Apparatus for fumigation with sulphur
dioxide, tobacco dust or formalde
hyde)
- 235 OLF (1911) Description of a Chinese mosquito lamp
Arch Schiff's Trop Hyg, XVI Abst in
Jl R A M O, XVI, p 584
- 236 PARSONS A C, and BROOK
G R (1919) The mosquito problem in Britain sugges
tions for a winter campaign against the
important mosquitoes, with notes on
insecticides *Jl R A M O, XXXII, 1*,
pp 1-23
(Cresol, potassium cyanide, chlorine,
sulphur dioxide, formalin, etc)

- 237 PARYKHIEV, T A (1935) On the methods of destruction of mosquitoes *Jl Econ Entom* XVIII 6 pp 839-840 *PIEB*, XIV p 28 (Waste tobacco)
- 238 RICHMOND A E., and MENDIS J C (1930) A report on investigations carried out in Pestawar during 1927 in connection with malaria prevention among troops *Pec Mal Surr Ind I* 3 pp 205-290 (Sulphur dioxide cresol)
- 239 SHAPIRO J M (1933) Smoke production as a measure of mosquito control *Trans Roy Soc Trop Med and Hyg* XVII 4 pp 267-264
- 240 SRINGAREV N I (1936) Observations on the control of hibernating mosquitoes *Russ Jl Trop Med* 4 pp 23-25 *RIEB* XVI p 203 (Tobacco dust)
- 241 SMITH J B (1908) The house mosquito a city town and village problem *N Jers Exp Sta Bull* No 218 (*Datura stramonium*)

Miscellaneous Measures against Adult Mosquitoes

- 242 BACOT A W (1914) Naphthalene for the destruction of mosquitoes in covered cisterns and wells *Brit Med Jl* July 4th p 15
- 243 DAVIS N C (1936) Study on the dispersion of resting Anopheles mosquitoes from dwellings in Brazil *Amer Jl Hyg* VI March Suppt pp 23-35 (General review)
- 244 DAVIS N C *et al* (1931) Notes on anti malarial work against adult mosquitoes *Rev Univ Buenos Ayres* XXIV pp. 112-1031 *RIEB* XXII, p 78
- 245 LECHETTI A B (1936) Experiments on the action of toxic substances on adult mosquitoes *Rev Soc Ent Argent Buenos Ayres I* 3 pp 29-32 *RIEB* XV p 110 (Poison baits with honey and mercury bichloride boric acid arsenious acid diethyl potassium arsenate potassium arsenite potassium cyanide)
- 246 PRINCE J A LF (1936) Destroying emerging Anopheles as a malaria control measure *U S J Pub Hlth Rpts*, XLI #3 pp 1230-1236 (Battlers and catching tubes)

- 247 REITLER R. (1930) The role of measures against a full mosquito in the prevention of malaria *Arch Schiff's Trop Hyg.* XXXIV, 4 pp 198-210 *RAFB* XVIII, p 189
(Mosquitoes caught in test tubes wetted inside with kerosene, or woollen garments hung in dark places)
- 248 TAKASHIMA K. (1927) Carbon tetrachloride as applied for the extermination of mosquitoes and flies *Jl Pub Hlth Japan* III, 6 pp 1-9 *RAEB*, XVI, p 12

Miscellaneous Measures against Larvæ of Mosquitoes

- BACOT A W (1914) See No 247
(Naphthalene)
- BALFOUR, A (1913) See No 218a
(Izo izal sanitas okol, creosote)
- 249 BARRIERI A (1928) Fluorescent sensitizing substances as larvicides The photodynamic action of light *Rev Med*, VII 4 pp 456-467
Eng Summ., p 626
- 250 BISHOP, G H (1919) Larvicides *U S A Pub Hlth Bull*, No 104, p 165
(Water gas tar)
- 251 BOREL E (1905) Note sur un procédé simple pour empêcher la ponte des moustiques dans les collections d'eau familiales indigènes *Bull Soc. Path Exot* XVIII 10 pp 779-781
(Lime)
- 252 BOREL, F (1926) Au sujet de l'action empêchante de la couleur blanche sur la ponte des moustiques, dans les jarres indigènes blanchies à la chaux *Ibid.*, XIX 8, p 702
(Lime)
- 253 BUNAT VARILIA, P (1928) La verdunisation dans la lutte contre la fièvre jaune *C R Acad Sci Fr* CLXXVII 22 pp 1005-1006
RAEB, XVII p 69
(Chlorine)
- 254 CAMPBELL F L (1930) A comparison of four methods for estimating the relative toxicity of stomach poison insecticides *Jl Econ Ent.* XXIII 2 pp 357-370 *RAFB*, XVIII p 191
(Acid lead arsenate and sodium fluosilicate)
- 255 CAWSTON F G (1927) Mosquito proof gutterings *Jl R A. M C.* XLIX, 6, p 441

- 256 COVIEL G (1928) Malaria in Bombay, 1928 (Govt Press, Bombay)
(Wells, cisterns hydrants fountains, etc.)
- 257 DARLING S T (1912) A mosquito larvicide disinfectant and the methods of its sterilization *Amer JI Pub Hlth*, Feb
(Panama Canal larvicide)
- 258 DUREN, A N (1930) Les mesures prises contre la fièvre jaune à Matadi en 1928 *Bull Off Int Hyg Pub* XVII, 2, pp 287-300 *RAFB*, XVIII, pp. 169-180
(Chlorine from sodium hypochlorite)
- 259 FLIXINGTON I S C (1912) Mosquito destruction *Brit Med JI*, I of 1912, p 106
(Potassium cyanide)
- 260 FREDERICK F S (1929) Experiments to note the larvicidal effects, if any, of sodium nitrate potassium nitrate and magnesium sulphate in natural waters *Ind JI Med Res*, XVII 1 pp 249-250
- 261 FEEGRADE F S (1923) Experiments with Isotex as a larvicide *Ind Med Gaz*, LXIV 5, p 252
- 262 FERMI, C (1926) Note di anofelinologia *Riv Med* V 3, pp 286-301
(Creosote)
- 263 FLETCHER T B (1924) Carbon bisulphide not efficacious as a mosquito larvicide *Procs 5th Entom Mfg Pans*, 1923 pp 226-227
- 264 GATTELLI, P A R (1929) *Ann Rept Med Dept F U S* 1928, pp 69-74
(Sodium silicofluoride, calcium arsenate, Paris green calcium cyanide)
- 265 GIBSON, A (1928) Waste sulphate liquor from paper mills as a larvicide *Sci American*, June p 552
- 266 GIBSON A (1928) Mosquito suppression in Canada in 1927 *Procs 15th Ann Mtg N Amer Mosq Fzt Ass*, pp 136-146
(Waste sulphate liquor powdered *Derris* root pyrethrum powder as larvicides)
- 267 GIBSON A (1929) Mosquito suppression in Canada in 1928. *Procs 16th Ann Mtg N Amer Mosq Fzt Ass*, pp 102-108
(Pyrethrum and *Derris* dusts)
- 268 GINSBURG I M (1928) Studies of mosquito oils and dust larvicides. *Procs 15th Ann Mtg N Amer Mosq Fzt Ass*, pp 63-65
(Dust consisting of kaolin saturated with 20 per cent fuel oil and 6 per cent high boiling point hydrocarbons obtained from coal tar)

- 269 GORMAN, A E (1919) Sub aqueous saws *U S A. Pub Hlth Bull* No 104, pp 58-63
- 270 GUNASEKARA, S T (1912) Potassium cyanide as a larvicide *Brit. Med JI*, p. 991
- 271 HARDENBURG, W E (1919) Larvicides *U S A Pub Hlth Bull* No 104, p 164
(Creosote, nitre cake)
HARGREAVES F (1923) See No 98
(Cactus paste)
- 272 HORN, A E (1924) A simple device for the mosquito protection of water tanks casks etc, without the use of gauze *Jl Trop Med and Hyg*, XXVII, 12 p 175
(Automatic cover)
- 273 JAMES S P (1914) Summary of a year's mosquito work in Colombo *Ind Jl Med Res*, II, 1, pp 240-241
(Cyllin)
- 274 MACFIE J W S (1923) A note on the action of lithium chloride on mosquito larvæ *Ann Trop Med and Paras* XVII, 1, pp 9-11
- 275 MARCOVITCH, S (1908) Studies on the toxicity of fluorine compounds *Pull Tenn Agric Exp Sta*, No 139 *RAEB* XVI p 213
(Organic acids metals naphthalene, fluorine compounds)
- 276 MATHESON R, and HINMAN, E H (1928) A new larvicide for mosquitoes *Amer Jl Hyg*, VIII 2 pp 293-296
(Borax)
MAYNE J F, and JACKSON, W R (1920) See No 115
(Cresol sanitas okol bleaching powder, kerosene)
- METZ C W (1919) See No 31
(Creosote, nitre cake)
- OSBORN W (1906) See No 212
(Lime)
- 277 PULBERNAU, S DE (1907) Use of chopped up prickly cactus as larvicide *Ann d Hyg Med Colon*, IV Abst in *Jl R A M C* XI p 209
- 278 RICKARD E R, et al (1928) Investigations on the value of the waste products of sugar cane crushing as a good and cheap larvicide 4 *Reun Soc Argent Pat Reg Agric, Bol Inst Clin Quir*, IV, 28-31, pp 153 156
RAEB, XVII, 3, p 60
- ROSS R, and EDIE, E S (1911) See No 101
(Panama larvicide, sanitas okol mercuric chloride, copper sulphate oxalic acid potassium cyanide)

- 79 POURAUD E (1917) I eclos on de l'œuf et les st mutants d'éco-
los on chez le *Stegomyia* de la fièvre
jaune Appli at on à la lute anti lar
vaire *C R Acad S* CLXXXIV
74 pp 1491-1497 *RAEB* XV
p 161
(Sodium hypochlorite)
- 80 SAUTET J (1930) A propos de l'emploi des hypochlorites dans
la lutte contre la fièvre jaune *Bull
Soc Pa A Ext* XXIII 2 pp 207
208
- 81 SEW S K (1915) Observations on the respiration of the
Cul de Ind *Jl Med Res* II 3
pp 681-697
(Effects on larvæ of vapours of various
substances incluin_g benzene and
kerosene)
- 82 SEW S K (1911) A note on the effect of mercurous
chloride on Culex larvæ and discussion
Proc 44 Ent Mty Amer pp 184
191
- 83 SEW OR WHITE R A (1918) Studies on malaria as it affects Indian Rail
ways Govt Press Calcutta
(Borrow pits etc)
- 84 STRICKLAND C (1919) Soap as a mosquito larvicide *Trans Roy
Soc Trop Med and Hyg* XVII 6
pp 509-510
- 85 TH BAULT J H (1918) Vegetable powder as a larvicide in
the fight against mosquitoes *Jl
Imer Med Ass* LXX 1 pp 175-
176
- 86 TICHENKO O D (1918) Emploi de préparations arsenicales dans la
lutte contre les larves d'anophèles
Bull Soc Path Ext XVI 8 pp 655-
656
(Arsenite et chlorure cuivreux and
sodium arsenate)
- 87 TWYNN C R (1919) Mosquito control at Ottawa Ontario 5th
Ann Rept Ent Soc Ontario 1919
pp 17-17
(Oils *Derris*)
- VALERIO B C and JONQU
J R DE (1909) See No 157
(Sulphuretted hydrogen saprol
powder)
- 88 VALLE V (1917) The afforestation of dunes in relation to
ant malarial works *Proc Mal VI*
pp 85-90 Eng Summ p 19
- 89 WAGENHAUS H H (1919) Sub-aqueous saws *U S A Pub H H
Bull* No 104 p 60

- 290 WILLIAMSON H B (1924) The use of gases and vapours for killing mosquitoes breeding in wells *Trans Roy Soc Trop Med and Hyg* XVII 8, pp 485-519
(Formaldehyde sulphur dioxide chloropicrin chlorine, naphthalene steam)
- 291 WISE, K S, and MINERT E P (1912) Experiments with crude carbolic acid as a larvicide in British Guiana *Ann Trop Med and Paras* VI 3B pp 327-330

Natural Enemies *

- APFELBECK V (1925) See No 119
(*Hydrometra*, *Velia*, *Gyrinus*, *Corixa*, *Notonecta*, *Crustacea* the most important)
- 292 ATKINSON J M (1909) A possible natural enemy to the mosquito *Il Trop Med and Hyg* XII 17 pp 255-256.
(*Laepa sinensis*)
- 293 BALFOUR A (1914) Birds and malaria *La cet CLXXXVII* No 4762 p 1326
- 294 BALFOUR A (1923) Mites on mosquitoes *Jl R A M C* XL 2 pp 122-127
- 295 BARBER M A and KING C H (1927) The tadpole of the spadefoot toad as an enemy of mosquito larvae *U S A Pub Hlth Rpts*, XLII 52, pp 3189-3193
- 296 BOYD J E M (1922) On the incidence of mites on mosquitoes *Jl R A M C* XXXVIII 6, pp 459-460
- 297 BRAGINA (1929) A hydra devouring an *Anopheles* larva *Beihfte Arch Schiffs Trop Hyg* XXXIII 3 pp 57-58
- 298 BRITTON W E, and VIERCK H L (1903) Report on mosquito investigations *Conn Exp Sta* for 1904 pp 253-310 In HOWARD DYAR and KNAB 1911 (No 24)
(*Hydrophilus*)
- 299 DEMPWOLFF (1904) In HOWARD DYAR and KNAB 1911 (No 24)
(*Notonecta*)
- 300 DERIVAUX R C (1916) A note on the predaceous habits of *Dytiscus* (Whirligig Beetles) towards *Anopheles* larvae *U S A Pub Hlth Rpts*, XXXI, 20 pp 128-1230

* References to **Bats** (Nos 48-54), **Deterrent Plants and Trees** (Nos 117-154) and **Fish** (Nos 196-217) are given under separate headings

- 301 DIXON, S G (1914) The duck as a preventer of malaria *Jl Amer Med Ass.* LXIII, 14, p 1203
In *Lancet* II of 1914, No 4753, p 1102
- 302 DYE, L (1905) Les parasites des Culicidés *Arch Paras.* p 60.
- 303 DYE, W H (19-4) Parasitic mites on mosquitoes *Jl R A M C.* XLII, 2, pp 87-102.
- 304 EYSELL, A (1903, 1907) In HOWARD, DYAR and KNAB, 1912 (No 24)
(*Nepa*, *Banatra*, *Naucaus*, *Notonecta*, *Corixa*, *Limnoblatta*, *Hydrometra* locusts scorpion flies wasps, robber flies)
- 305 FEARNSIDE C J (1900) Parasites found on mosquitoes *Ind Med Gaz.* XXV, p 123
- 306 FOLLY E H and YVERNAULT, A (1909) Campagne antipaludique dans une oasis Saharienne, à Beni Ounif (Sud Oranais) en 1908 In HOWARD, DYAR and KNAB 1912 (No 24)
(Ephemeroïd larvae)
- 307 HAMLYN-HARRIS, P (1929) The relative value of larval destructors and the part they play in mosquito control in Queensland (33 refs.) *Procs Roy Soc Queensland* XLI 3 pp 23-38
- 308 HAMLYN-HARRIS, P (1930) *Holthorns* (Col) as possible factors in natural control of mosquitoes in Queensland *Bull Ent Res* XXI, 2 p 159
(Small beetles)
- HOWARD, L O DYAR H G, and KNAB F (1912) See No 34 pp 156-178
- 309 ITENGAR, M O T (1929) Parasitic nematodes of Anophelis in Bengal Trans 7th Cong I F A T M, 1927, III, pp 128-133
- 310 ITENGAR, M O T (1929) Microsporidian parasites of Anopheles larvae *Ibid.*, pp 136-142
- 311 LAMBORN R H (1890) Dragon flies versus mosquitoes The Lamborn Prize Essays New York (Essays by ABRAHAM, C B, WEEKS, A C and BRUYENMULLER, W)
- 312 LAMBORN W A (1900) The habits of a dipteran predaceous on mosquitoes in Nyasaland *Bull Ent Res.* XI, 3, pp 278-291
(*Liapt*)
- 313 LAVIEU G (1921) Les parasites invertébrés hématophages, Paris
- 314 MACFIE J W S (1923) A note on a beetle which preys on mosquito larvae *Bull Ent Res.* XIII 4 p 403
(Tiger beetle)

- 315 MACGREGOR, M E (1920) The question of natural enemies *Jl R 4 M C*, XXXIV, 3, pp 248-250
- 316 MATHESON, R, and HINMAN, E H (1929) The vermilion spotted newt (*Desmognathus fuscescens roseni*) as an agent in mosquito control *Amer Jl Hyg*, IX, 1, pp 188-191
- 317 MORISHITA, K (1926) Notes on the natural enemies of mosquitoes *Dept Hyg Govt Res Inst Formosa* Also *Jl Med Ass Formosa*, No 257 Eng Summ, p 5
- 318 PRUTHI H S (1928) Some insect and other enemies of mosquitoes *Ind Jl Med Res*, XVI, 1, pp 153-157
- 319 ROSS, R (1906) Notes on the parasites of mosquitoes found in India between 1895 and 1899 *Jl Hyg* VI pp 101-109
- 320 SHAW, F R (1925) Effects of temperature on aquatic life in cisterns *U S A Pub Hlth Bull* No 156, pp 63-71 (Musk turtle)
- SMITH, J B (1904) See No 102, pp 79-113 (Spiders, frogs toads lizards bats, birds, round worms dragon flies, attacking adult mosquitoes Fish, birds, Dytiscid larvae and adults, *Gyrinus*, *Neptis*, *Ranatra* *Psephenus*, *Corethra* dragon fly larvae, attacking larvae)
- 322 SPEER A I (1927) Compendium of the parasites of mosquitoes *Hyg Lab Bull* No 146 *U S A Pub Hlth Service*
- VALERIO, B G and JONGE, J R DE (1909) See No 152 (Frogs, *Discoglossus*, tritons fish)
- 323 YABLOKOV, N A (1926) Little known enemies of *Anopheles* larvae *Russ Hydrob*, Zs, V, 7^o, pp 194-195 *RAEB*, XIV, p 220 (Hydras, leeches)

Oil.

- 324 ANTONIBOV, A (1923) The action of petroleum on mosquito larvae *Ann d Igine* XXXIII, 11, pp 796-801 *RAEB*, XII p 23
- 325 ASIATIC PETROLEUM CO, SINGAPORE (1929) Anti malarial oil (MS)
- 326 BACOT, A W (1916) Experiments dealing with the destruction of *Stegomyia fasciata* Tell *Fever Comm*, W Africa III, pp 153 155

- 327 BARNES M E (1925) A survey of the turpentine industry for possible larvicidal substances *Amer Jt Hyg*, V. 3, pp 309-314
- 328 BARNES, M F (1925) The toxic action of oil films upon *Anopheles* mosquito larvae with particular reference to pure oil films. *Ibid.*, pp 315-329
- 329 BOYE, K (1927) Malaria control at Birnagar, 1926 *Calc Med Jt* XVI, pp 350-354
- 330 BOYE, K (1929) Proceedings of a conference held at Birnagar (Dental) to discuss the problem of malaria control at Birnagar *Ind Med Gaz* LXIV, 6, pp 323-328
(Oil said to spoil fish for food)
- 331 BOYS F DE (1930) Etude expérimentale sur quelques méthodes employées dans la lutte contre les larves d'*Anopheles* *Bull Soc Path Exot*, XXIII 4 pp 402-427
(Liquide paraffin pétrole huile brûlée)
- 332 CHARRIER, H (1925) Action des huiles et du pétrole sur les larves de moustiques *C P Soc Biol*, XCI 16 pp 1240-1241 *R4FB* XIII, p 116
- 333 CLARKSON L M (1930) Oil mixtures and oil equipment for *Anopheles* larvae control of impounded areas *South Med Jt* XXII 4, pp 397-398
- 334 DARLING, S T (1910) Studies in relation to malaria Vol 2 Panama Commission Ibm
- 335 DELBOEUF, J (1895) La destruction des moustiques *Proc Scient* IV p 729
(Records that he used oil in 1845 as larvicide)
- 336 EDITORIAL (1929) Methods and mechanism of mosquito control *Malayan Med Jt*, III, 1, pp 53-55
- 337 FREUNDLICH, F S (1929) Results of experiments with crude oil products of the Burma Oil Co., Ltd. Rangoon, as larvicide *Ind Med Gaz*, LXIV 5, p 253
- 338 FISHER H C (1922) *Rept Hlth Dept Panama Canal* for 1921
- 339 FISHER H C (1923) *Ibid* for 1922
(Use of barrel pump)
- 340 FREUNDLICH S B, and ATSAAT L F (1914) The effects of petroleum oils on mosquito larvae *Jt Fein Entom*, XI 3 pp 299-308

- 341 GIBSON A (1927) Mosquito investigations in Canada in 1926 *Procs 14th Ann Mtg N Jers Mosq Ext Ass*, pp 110-115
(Addition of caustic soda to increase spreading power)
- 342 GINSBURG, J M (1927) The effects of various chemicals on the spreading and penetration of oils in different mosquito breeding places
Ibid pp 52-64 (17 refs)
- GINSBURG, J M (1928) See No 268
- 343 GINSBURG J M (1929) Relation between toxicity of oil and its penetration into respiratory siphons of mosquito larvae *Ibid*, 16th *Mtg* pp 50-63
- 344 GRATCH, I (1930) Malaria in the commune of Ravenna *Rev Val*, IX, 1 pp 1-34 Eng Summ p 94
(Recommends 'Leron')
- 345 GREEN, H W (1924) The effect of oil upon *Anopheles* mosquito larvae *Amer Jl Hyg*, IV, 1 pp 12-22
- 346 GREEN, H W (1925) The theory of the use of oil for destroying mosquito larvae *Amer Jl Pub Hlth* XV, 3, pp 219-222 *TDB San Suppt*, 1925 pp 110-111
- 347 GRIFFITHS, T H D (1919) Oiling *U S A Pub Hlth Bull* No 104 pp 113 and 121
(L shaped oil booms)
- 348 GRIFFITHS, T H D (1925) Method of applying oil under pressure as a larvicide *Ibid* No 156 pp 15-19
- 349 HACKER H P (1919) Malaria problem at Gemas *Mal Bur Rpts*, F M S, I, pp 53-60
- 350 HACKER H P (1925) How oil kills *Anopheles* larvae *Ibid* III 62 pp
- 351 HARDENBURG W E (1919) Oiling *U S A Pub Hlth Bull* No 104 p 128
(Oil soaked sawdust, sunk in a box in the water)
- 352 HEADLEE T J, et al (1928) Mosquito control *Rept N Jers Agric Exp Sta*, 1926-27, pp. 149-156
(Cresylic acid increases toxicity)
- 353 HOWARD, L O (1932) An experiment against mosquitoes *Insect Life*, V, pp 12-14, 109-110 190
- 354 HOWARD, L O (1901) Mosquitoes New York
- 355 HOWARD, L O (1929) The work with mosquitoes throughout the world in 1928 *Procs 16th Ann Mtg N Jers Mosq Ext Ass* p 13

- 356 HICWOOD, A S (1923) Some observations on the use of oils as a larvicide *Health*, I, 9, pp 255-259
RAFB, XII, p 39
(Mixture of kerosene and crude oil, $\frac{1}{2}$ oz per sq yd in sheltered pools, 1 oz. in open country)
- 357 JATZENKO T (1926) Use of naphtha and petroleum for the destruction of the larvae of *A. maculipennis* *Per Microb and Epidemiol.*, V, 4, pp 279-283 *ALB*, XV, p 31
- 358 JOHNSON, F B (1921) Oiling and larvicides *U S A Pub Health Bull* No 115, pp 100-102
(Oil soaked sawdust)
- 359 JOHNSON, E B (1922) Efficient and economical use of oil in the prevention of malaria *South Med J.*, XV, 5, pp 371-376
(Oil soaked sawdust also sacking and cotton waste soaked in oil)
- 360 KIRBY, C H (1923) Larvicides *U S A Pub Health Bull* No 150, pp 141-142
- KLICLER I J (1925) See No 108
(Kerosene, crude oil, castor oil etc.)
- 361 KLICLER, I J, and WHITE MANN, I (1922) Malaria control demonstrations in Palestine *Rept of Hadasah Med Organization TDR*, XVI, pp 298-299
(Kerosene, crude oil, castor oil)
- 362 LYAK W N (1921) On the improvement of oiling fluids for anti mosquito work *Jl Trop Med and Hyg* XXIV 4 pp 37-40
(Castor and other vegetable oils)
- 363 MCFARLAND, J (1929) The use of oil for the destruction of mosquito larvae and pupae *Jl Amer Med Ass*, XCI 25 p 2014
(Records earliest reference to use of oil as larvicide 1793)
- 364 MACVIE, J W S (1917) The limitations of kerosene as a larvicide *Bull Ent Res* VII, 3, pp 277-293
- 365 MARSHALL, J F. (1927) Principles and practice of mosquito control *Brit Mosq Cont Inst. Hayling Island*
- 366 PETERSON J P (1928) Mechanical oiling equipment utilized in New Jersey mosquito work, and its further possibilities. *Proc 15th Ann. Mtg N Jers Mosq Ext Ass*, pp. 45-55
(Use of crank-case oil)

- 367 PETERSON, J P. and GINSBURG, J M (1929) Two years' study and practical use of crank case waste oil as a mosquito larvicide *Procs 16th Ann Mtg N Jers Mosq Ext. Ass*, pp 92-101
- 368 POCKELS, A (1894) On the spreading of oils on water *Nature*, L, pp 223-224
- 369 PRINCE, J A LE (1915) Control of malaria Oiling as an anti mosquito measure *U S A Pub Hlth Rpts*, XXV, 9, pp 599-608
- 370 ROOK, H DE, and SWELLENGREBEL, N H (1927) Effets comparés du 'vert de Paris' et de la paraffine liquidum, comme larvicides *Bull Soc Path Exot*, XX, 2, pp 123-127
- 371 RYLES, C S, and MAJUMDAR, B C (1927) The use of petrol as a larvicide in wells, tanks and similar collections of water used for drinking or domestic purposes *Malayan Med Jl*, II, 4 p 144
- 372 SAMMIS, P H (1927) Progress of mosquito work in Nassau County, New York, for the year 1926 *Procs 14th Ann Mtg N Jers Mosq Ext Ass*, pp 107-110
(Description of various power sprayers and pumps)
- 373 SCHARFF, J W (1927) Notes on practical measures of malaria prevention from the point of view of mosquito control *Malayan Med Jl*, II, 2, pp 49-56
(Solar oil, kerosene and crude oil, apparatus for dripping oil on streams)
- SEN, S K (1910) See No 281
(Effects of benzene and kerosene vapours on larvæ)
- 374 SENIOR WHITE, R A (1925) The eradication of malaria from a rubber estate *Ind Jl Med Res*, XII, 3, pp 545-551
- 375 SOLLMANN, T (1918) The spreading power of coal oils *Jl Amer Med Ass*, LXXI, p 1553
- 376 SPAULDING, C H (1919) Oiling *U S A Pub Hlth Bull* No 104, p 124
(Drip cans)
- 377 SWELLENGREBEL, N H and ROOK H DE (1928) Effet comparé de quelques larvicides *Bull. Soc. Path Exot*, XXI, 2, pp 109-112
- 378 TAKATSUKI, A (1917) An essential property of petroleum for mosquito control *Kyoto Igaku Zasshi*, XIV, 7, p 84 *RIEB*, VI, p 130
- 379 TAKATSUKI, A (1920) The real action of petroleum on mosquito larvæ *Ibid*, XVII, pp 67-70 *TDB*, XVIII, p 190

- TWINS, C R. (1927) See No 297
(Caustic soda added to improve spread.)
- 380 WALCH, E W, and BOYNE WEPSTER J (1929) A comparative examination of some oils and mixtures of them as larvicides
Genesek Tijd Ned Ind, LXIX, 4, pp 333-343 With Eng Summ
(Recommend studie also crank case oil with kerosene and coconut or castor oil)
- 381 WALLACE R B (1929) Reduction in oiling at certain seasons
Malayan Med Jl IV, 3 pp 92-95
- 382 WALLACE R B (1930) Cessation of oiling in a malarious area during a certain season of the year
Ibid V, 4 pp 117-125
- 383 WETMORE W O (1926) A device for economically oiling ponds and other waterways
Jl Amer Med Ass, LXXXVII, p 1298
- 384 WIGGLESWORTH V B (1930) A theory of tracheal respiration in insects
Proc R Soc B, CVI p 229
- 385 WILLIAMSON, K B (1926) *Ann Rept Mal Bur Inst Med Res*, F M S, for 1925
- 386 WILLIAMSON K B and LAMAMONY I D (1928) Comparative tests on the larvicidal and spreading powers of rubber crude and other oils
Malayan Med Jl, III, 2, pp 89-92

Paraformaldehyde

- 388 BARBER M A and KOMP W H W (1927) Some tests of the larvicide Stoxal
U S A Pub Hlth Rpts, XLII, 31 pp 1937-2004
- 387 BAUVALLEY H (1928) Essai de lutte antilarvaire par l'emploi du poudre larvicide dans une zone inondée du Bas Dahomey
Bull Soc Path Exot XXI 4 pp 323-325
- BAKER F DE (1930) See No 331
- (JENSEN J M (1928) See No 269
- 389 MERTENS W A (1929) My experience of Stoxal as a larvicide
Genesek Tijd Ned Ind LXVIII 7 pp 1010-1014 *RAF B* XVII p 111
- 390 POIRAUD F (1929) Emploi du Trioxyméthylène en poudre pour la destruction des larves d'*Anopheles*
C R Acad Sci, CLXX 25 pp 1521-1522
- 391 POIRAUD F (1930) Mode d'action du Trioxyméthylène en poudre sur la larve d'*Anopheles*
Ibid, CLXXI 1 pp 31-32
- 392 POIRAUD F (1934) Emploi des poudres larvicides légères dans la lutte contre les moustiques
Bull Soc Path Exot XIX 4 pp 287-302

- 392 SCHUURMAN, G J, and Experiments with paris green for the des-
SCHUURMAN TFF BORJEL truction of Anopheles larvae *Middel-*
HUIJINK, A (1928) *Vol. Ned. Ind*, XVII, 2, pp 207-219
- 393 SWELLENGREBEL, N H, *et al* Anti-larval measures around Mideembhek
(1929) (North Holland) in 1928 (Ms) *2nd*
Rept to Inter Hlth Coun, Rockefeller
Foundation
- SWELLENGREBEL, N H, and See No 377
ROOK, H DE (1928)

Paris green

- 394 BARBER, M A (1925) Some further experiments with paris
green as a larvicide for Anopheles
larvae *Trans 1st Inter Cong Med*,
Rome, pp 153-157.
- 395 BARBER, M. A, and HAYNE, Arsenic as a larvicide for Anopheles
T B. (1921). larvae *U S A Pub Hlth Rpts*,
XXVI, 49, pp 3027-3034
- BARBER, M A, KOMP, See No 5
W H W, and HAYNE,
T B (1926)
- 396 BIRN (1930) Observations on the effect of paris green
on the larvae of *A. maculipennis* Meig
Sanitent Byull, I, 3, pp 19-21
RAEB, XVIII, p 255
- 397 BLAGOVESHCHENSKI, D I Paris green as a larvicide in the control of
(1928) malaria mosquitoes *Rev Microb and*
Epidem, Parasit, VII, 2, pp 156-167
RAEB, XVI, pp 200-201
- BOYD, M F (1926) See No 5
- 398 BOYD, M I, and DAVIS Experiments in Brazil Unpublished
N C (1924) Repts to International Health Board,
New York. Quoted by HACKFET 1925
(No 417)
- 399 BUEN, E DE (1928) Paris green as a larvicide *Med Pais*
Cal, I, 1, pp 64-73
- 400 BUEN, F DE (1929) Indicaciones practicas el empleo del verde
de Schweinfürt (verde Paris) *Dir Gen*
San, Madrid
- 401 BUEN, E DE (1929) An experimental study of some larvicidal
substances against Anopheles *Med*
Pais Cal, II, 5-6 pp 401-430,
503-540 (51 refs)
- BUEN, E DE (1930) See No 331
- 402 BUEN, S DE, and BUEN, The first experiments in Spain with paris
E DE (1927) green in anti malarial work *Bol Tec*
Dir Gen San, II, pp 161-168 *Summ*
Bull Hyg, II, pp 582-583

- 403 PURS, S DE and BURY, Trabajo sobre el empleo del verde Paris
F DE (1927) *Ind Gen. San., Madrid*
- 404 BURY N M (1931) A simple and inexpensive portable
screener for paris green diluents *Rec.
Mal Surc Ind II 2*, pp 333-
335
- 405 CHALAM, B S (1927) The possibilities of paris green as an An-
opheline larvicide *Ind Jl Med Res.*
XIV 4, pp 807-803
- 406 CHALAM B S (1930) Further observations on paris green as an
anophelines larvicide *Proc Mal Surc
Ind I 4*, pp 515-522
- 407 CHALAM B S and DALAL
P A (1926) Bombay mill ponds and Anopheline
control *Ind Med Ga.*, LXI 3,
pp 115-116
CLYDE D (1931) See No 220
- 408 COOK S S (1930) Malaria control in Haiti. *South Med
Jl XXIII 5* pp 454-459
- 409 COOK S S and WILLIAMS
L L (1928) Airplanes and paris green in the control
of Anopheles production *Ibid.*, XXI, 9,
pp 754-759
- 410 DALAL P A, and MADON
F F (1926) Bombay mill ponds and Anopheline
control *Ind Med Ga.* LXI 8,
pp 383-387
- 410a DALAL P A, and MADON
F F (1927) A new diluent for paris green *Ibid.*,
LXII 10 pp 554-557
(French chalk)
- 411 DALAL P A and MADON
F F (1928) Report on anti mosquito work in mill
ponds etc (MS) Sept to Messrs F
D Sassoon Bombay
- 412 DOLLOFF A F (1929) The use of stearates (calcium and alu-
minium) as diluents for paris green in
Anopheline control *J S I Pub Hlth
Rpts XIV 43* pp 2588-2594
CATER I A L (1929) See No 264
- 413 GILL, D G, BAKER C C
and SIMS I C (1929) Practical application of paris green as a
larvicide for Anopheles larvae in an
impounded lake in Alabama *South
Med Jl XXII 4* pp 399-404
- 414 GRIFFITHS T H D (1927) Mist and method of applying paris
green for destruction of sub-arctic
feeding mosquito larvae *E S A
Pub Hlth Rpts XIII 44*, pp 2701-
2705
- 414a GRIFFITHS T H D (1930) Malaria control by use of paris green
preliminary report of countermeasures
work in Dougherty County Georgia.
South Med Jl. XXIII 5 pp 428-
436

- 415 GRIFFITHS T H D, *et al* (1923) Discussion on the use of paris green *U S A Pub Hlth Bull* No 137, pp 46-48, 177
(Harmless in boilers)
- 416 GUEIMINO, G (1928) On larvicide powders *Glas Tent Aug Zav*, III, 1-6, pp 4^a-46 *RAFB*, XVIII, p 111
- HACKER, H. P (1925) *See No 350*
- 417 HACKFUT, L W (1925) The importance and uses of paris green as an Anopheles larvicide *Trans 1st Inter Cong Med Rome*, pp 158-166
- 418 HACKFUT L W (1929) Malaria control through anti mosquito measures in Italy *Trans Roy Soc Trop Med and Hyg* XXII 6 pp 477-499
- 419 HAYNE T B (1923) Paris green as a larvicide for Anopheles larvae *U S A Pub Hlth Bull* No 137, pp 41-45
- 420 HERRMANN O, KOLOSSOW, J and IPIRY, N (1926) Paris green as an Anopheles larvicide *Cent F Balk 1st Orig* XCVIII 7-8 pp 547-552 *Samm Bull Hyg*, I p 809
- 421 HERRMANN O, KOLOSSOW J and LIPIN, N (1928) Is paris green a better larvicide than kerosene or liquid paraffin? *Arch Schiffs Trop Hyg*, XXXII 3 pp 140-143
- 422 KING W V (1924) Airplane dusting *Trans 6th Ann Conf Mal Field Workers New Orleans* Quoted by NICHOLLS, 1927 (No 434)
- 423 KING W V and BRADLEY G H (1926) Airplane dusting in the control of malaria mosquitoes *U S A Dept Agric Circ* No 167
- KLIGLER I J (1925) *See No 108*
- KLIGLER, I J (1926) *See No 109*
- 423a KOROSTELEV V F (1928) An experiment with paris green for the control of the larvae of anopheline mosquitoes *Russ Jl Trop Med* VI 6, pp 385-389 *RAFB*, XVII p 6
- 424 KOROSTELEV, V F (1929) Encore sur l'effet larvicide du vert de Paris *Ibid*, VII 4 pp 299-301 *RAFB*, XVII, p 201
- 424a LOPEZ I A (1929) Arsenical dermatitis with case report *Univ Med Jour* 18th Ann Rept pp 189-191
- 425 MALARET, P S (1929) Control of malaria in the Preston Division *Ibid*, pp 74-79
- 426 MALTBERGER, J R (1929) Malarial preventive measures as applied in the Chiqui Land Company *Ibid* pp 79-85

- 427 MARCOVITCH, S (1928) . The relative toxicity of arsenicals and fluorine compounds to various organisms *Jl Econ Ent*, XVI 1 pp 109-114 *R4FB*, XVI p 178
- 428 MARTINI, F (1927) Mode of use of arsenicals as larvicides *Arch Sch Hyg Trop Hyg*, XXXI, 2, p 53 *RAEB*, XV, p 73
- 428a MATTEI G DI (1929) On a dermatosis caused in larval control agents by using paris green *Riv Malariol*, VIII 6 pp 669-673 *Eng Summ* p 757
- 429 McLENEY H F (1927) The types of breeding place used by *Anopheles hyrcanus* in North and Central China *China Med Jl* XLI 4 pp 347-350
- 43 MURDOCH K I (1927) Effects of paris green on tabanic larvae *Jl Trop Med and Hyg* XXX, 15 pp 197-198
- 431 MINSIROLI A (1927-1930) The prevention of malaria in practice *Riv Mal* VI 3 pp 501-572 VII 4, pp 413-455 IX 6 pp 667-700
- 432 MINSIROLI A (1929) Prevention of malaria by paris green *Seuchenbekämpfung* VI 3, and 4, pp 155-161 238-244 *TDB* XXVII pp 219-220
- 433 NABRO C IVL (1927) Paris green as a larvicide *Arch Hyg Dept Hlth Brazil* 1 2 pp 143-158 *Abst in U S A Pub Hlth Rpts* XVIII 7 p 410
- 434 NICHOLLS L (1927) The use of copper acetoarsenite (paris green) as an *Anopheles* larvicide *Ceyl Jl Sci D II* 1 pp 21-30
- 435 OTTOLENGHI D *et al* (1929) Conditions relating to malaria and anti-malarial work in the anti-malarial campaign in Ferrara *Int Mal* VIII 2 pp 113-119 *Eng Summ* p 271
- 436 PARVUTSEV I A (1926) On the question of controlling mosquito larvae *Defence des Etats* III 2 1 pp 274-276 *R4FB* XIV 3 215
- 437 LAVLOVSKII F N *et al* (1929) Animal parasites and some parasitic diseases of man in Tadzhikistan *Leningrad R4FB* XVIII p 5
- 438 PRINCE J A LE *et al* (1927) Mosquito control by airplane memorandum on the distribution of paris green by airplane in the control of *Anopheles* production in uncultured pond near Ramber South Carolina Sept 8th 1927 *U S A Pub Hlth Rpts* XII 39 pp. 233-238

- 439 PRINCE J A LE and JOHNSON H A (1929) A new method for macul at dusta-
bution of paris green as an Anopheline
larvicide preliminary report *South
Med JI*, XVII 4, pp 373 374
(Electric blower)
- 440 PRINCE, J A LE and JOHNSON H A (1929) Development of a power dusting appliance
for applying paris green as an An-
opheline larvicide *U S 4 Pub Hlth
Rpts*, XLIV, 17, pp 1001 1017
- 441 QUARF W T (1938) A trial of paris green as a larvicide in
moving water in Selangor F M S
With a summary of previous work on
paris green *Malayan Med JI* V,
1 pp 26-29
REYNE A (1934) See No 35
- 442 RICKARD E R and PEREZ A (1928) Larvicides in practical work Exper-
iments with petroleum and paris green
*Reun Soc Argent Pat Reg Norte (in)
Bol Inst Clin* IV, 28-31, pp 166-
171 *RAEB* XVII p 61
- 443 ROOK H DE (1927) Paris green as an insecticide for Ano-
pheline larvae Thesis Univ Am-
sterdam (46 refs) *P4FB* XV p 73
ROOK, H DE and SWEL See No 370
LENGKEBEL, N H (1927)
- 443a RIKAVISINIKOV B I (1930) Essays of the control of *Anopheles*
maculipennis Mg *Plant Protection* VI
3-6 pp 775 776 *RAEB* XIX p 21
- 444 SALISBURY E I and CORRIAN J A (1929) Control of malaria in Costa Rica *United
Fruit Co 18th Ann Rept* pp 88 97
- 445 SCHARFF J W (1926) An experiment with paris green as an
A ludlowi larvicide *Malayan Med
J*, I, 1 p 14
SCHUURMAN G J and See No 392
SCHUURMAN TEN BOKKFL
HUIJINK A (1928)
- 446 SCHUURMAN C J and Appl ation of paris green as an *Ano-*
pheline larvicide (2nd Rept) *Meded
Volled Ind* XVIII 1 pp 143-150
HUIJINK A (1929)
- 447 SELLA M (1929) *Cambusia* and paris green in the anti-
malarial campaign at Novigno *P:
Med*, VIII 4 pp 327 329
- 448 SHAPIRO J M (1923) Note on paris green *Procs 8th Mtg
Int'nal Adm Comm, Palestine
TDR San Suppl* 2 pp 81 82 1924
- 449 SEMIC C (1929) Can one protect Skopje from malaria
by an anti larval campaign with paris
green? *Glas Tsent Ak g Zor* IV
1-3 pp 29 37 *RAEB* XVIII p 3

- 420 BURR H. and PARKER H. Paris green as an Anopheles larvicide
(1929) *Ind Med Gaz.* LVII, ~ PP. 376-378
See No 393
- 421 SWELLENGREBEL A. H. et al Influence de la substance diluante sur l'action larvicide du vert de Paris.
(1929) *J r Mal.* VIII 2 p. 313
- 432 TIERHAGEN W. and FOWD (1927) Malaria in the Philippines *Jl Prev Med.* I 3 pp 200-234
- 433 WILLIAMS L. L. and COOK Paris green applied by airplane in the control of Anopheles production.
N. S. A. Pub. HHA 11th VII, 7, p. 409-480
- 434 WILLIAMS L. L. and FARRAR A. F. (1927) Length of life of *quadrinaculatus* after beginning of control of production
South Med Jl XVI 9 pp 32-33
- 435 WILLIAMS L. L. and FARRAR A. F. (1927) The life of *quadrinaculatus* Bureau of M. S.,
for 1927 *Ann. Rept. M. S. Dept.* 1927 pp 2-4
(Filtrate from suspension of Paris green toxic to larvae)
- 436 YARBOROUGH and SMITH A. N. (1929) A preliminary report on the suitability of Paris green as an Anopheles larvicide as applied to Luning control na.
Ind Med Gaz. LXX - PP. 81-84

Repellents.

- 437 BAILEY A. W. and TAYLOR H. (1919) The comparative effectiveness of certain culicids under laboratory conditions
Parasitology XI - PP. 221-230
- 438 BINKER C. W. and HIRSCHFELD A. H. (1923) Mosquito repellents
Amer Jl Trop Med. V 3 PP. 297-307
- 439 CECIL A. (1921) A preliminary report on the use of remote oil as a mosquito repellent
J S A Pub. HHA 11th VIII 10, 11 43-44 Also in *J S A Pub. HHA Bull.* No 127 PP. 11-15
- 440 COOPER C. I. (1923) Mosquito repellents and mosquito control
J S A Pub. HHA Bull. V 10 PP. 136-137
- 441 COOPER C. I. (1923) An improved mosquito repellent
Ind Med Gaz. LXXI 3 p. 601

- 461 DUNN M B (1920) Methods of protection from mosquitoes etc Canada Dept Agric Pamphlet n o No 55
(Mosquito nets fly proof tents, smudges, repellents)
- FERMI C (1906) See No 262
(Creosote, painted on walls etc)
- FERMI C and JUMBHO S (1900) See No 224
- FOWLER C E P (1926) See No 16
- 462 FREEBORN S B (1924) Observations on the control of Sierran *Aedes* Pan Pacific Ent, IV 4, pp 177-181 RAEB, XVI p 214
(Formulae for 2 repellents)
- 463 GIVEN D H (1922) The campaign against mosquitoes on board H M S Cadmus in 1912 Jl Roy Nat Med Serv VIII 4 pp 261-269
(Citronella)
- 464 HOWARD L O (1916) Remedies and preventives against mosquitoes U S A Dept Agric Farmers Bull No 444
- LAVERAN A (1907) See No 99 pp 578-579
- 465 MITCHELL E G (1907) Mosquito life p 10 New York and London
- 466 RUDOLFS W (1925) Studies of mosquito repellents Rept N Jers Agric Exper Sta 1923-24 pp 350-353
- 467 RUDOLFS W (1926) Investigations of mosquito problems Procs 13th Ann Mtg N Jers Mosq Fxt Ass, pp 33-54
- 467a RUDOLFS W (1930) Effects of chemicals upon the behaviour of mosquitoes Bull N Jers Agric Exp Sta, No 496

Screening

- 468 BARBER M A and KING C H (1929) Some notes on the limitations of screens in the prevention of malaria U S A Pub Hlth Rpts XLIV 10 pp 527-528
- 469 BOYD, J F M (1928) Anti malarial works and improvements Jl R A M C L, 4 pp 287-288
- 470 BOYD M F (1926) The influence of obstacles unconsciously erected against anophelines (houses and screening) upon the incidence of malaria Amer Jl Trop Med VI 9 pp 157-160

RELEVING

471 COONLE C I (1927)

CRAM C F (1909)
472 CRAIG C F (1914)

DARLING D T (1910)
DEADERICK W H (1900)
DEERY W F (1910)

473 EARLE W C (1910)

474 EIDSON R H VON (1914)

475 FINE C H (1910)

476 GAGE F H (1910)

477 GILES G M (1904)

478 HARAPIN J B (1910)

HARDENBERG W F (1910)

HIGH E (1911)

HENRI (1917)

HENSON G F (1913)

479 HILL L (1910)

480 HOPKIN J W (1913)

JACKMAN A (1911)

481 McARTHUR W I (1913)

MAN OLE F (1910)

482 McVILLIE HAYSON (1914)

483 MURKIN A C (1917)

Interim report of screening studies
Lefflore County Missouri pp 1-5
Tab Hk Rpts XLII 16 pp 110-112

See No 11 Pl 354-356
The prophylaxis of malaria with special
reference to the military service L 8
Army War Dept Off of Surg-Genl
Bull No 6 Summ Ind Med Gaz
LI 1916 p 232
See No 334

See No 14 pp 32-346
Aspects of malaria control J Trop Med
Hyg XLIX 13 pp 182-194
Malaria in Porto Rico Amer J Trop
Med X 3 pp 20-30

Prevention of malaria suggestions on
how to screen the home & keep out
effectively the mosquitoes which spread
the disease L 8 4 Tab Hk Rpts
XLIX 9 pp 503-504

The air inside a mosquito net Hist
Med JI 11 of 1910 p 1190

Observations on the passage of
phelino mosquitoes through screens of
various sized mesh L 8 4 Tab
Hk Rpts LVI No 150 pp 44-45

The wire gauze protectant in buildings
Ind J Med Hyg LVI 10 p 337

Further studies in mosquito proofing
barracks J P A M L 2
1912-130

See No 18 Chapter X
See No 11 pp 96-97

See No 11 p 304

See No 21 p 147

The air in a mosquito net Hist Med
JI 11 of 1910 p 1190

Ann Rept Mal Dept Board of Med
1910-1911

See No 19 pp 503-504

Mosquito netting J P A M L 2
1912-130

See No 30 p 310

Mosquito screening of ships Ind J Med
Hyg LVI 13 p 171-183

An experiment in mosquito proofing
barracks of Indian troops JI 11
of 1910 p 1190

See No 4 pp 219-223

- 484 ORENSTEIN, A J (1914) The technique of mosquito screening
Arch Schiffs Trop Hyg, XVIII, 1
pp 16-21
- PRINCE, J A LE, and ORENSTEIN, A J (1916) See No 34 pp 198-199
- REITLER R (1930) See No 247
- ROSS P (1911) See No 37, p 261
- 486 RUTHERFORD, N J C (1928) Mosquito proofing of barracks of British
troops in India *Jl R A M C*
L, 1, pp 76-78, and 4, pp 316-318
- 487 SBACCHI (1906) Prophylaxis by wire gauze *Atti della*
Soc per gli Studi della Malaria
See No 373
- 488 SCHUCKMANN, W. VON (1926) On the control of flies and mosquitoes
Zeitsch Ang Ent, XII 2 pp 332-339
RAEB, XV, pp 12-13
- 489 SIMPSON, W J (1917) The vizer anti mosquito and fly proof head
gear Invented by Mrs W J Simpson
Trans Foy Soc Trop Med & Hyg
X, 7, p 140
- 490 SNIDOW, H W (1919) Screening of residences *U S A Pub*
Health Bull No 104, p 19
- VISBECK and LACAZE, H (1924) See No 43, pp 215-243
- WATSON, M (1915) See No 44, p 179
- WATSON, M (1921) See No 45, pp 257-263, 364
- 491 WHYTE, G D (1910) The effect of a mosquito net on the
air within it (based on 80 analyses)
Brit Med Jl, II of 1910, pp 972-973
- 492 WURTZ, R (1907) Sûreté des tulles employés pour protéger
contre la piqure des moustiques *Rev*
Med Hyg Trop, IV, 4 p 181 In
ROSS, 1911 (see No 37)
- 493 ZIEMANN, H (1906) In MENDEL, *Hand der Tropenkrankh*, III
p. 445

Sprays.

- 494 ABBATUCCI, S, and ROUBAUD, E (1926) Expériences sur un liquide insecticide
commercial à base de pétrole et de
poudre de pyrèthre *Bull Soc Path*
Exot, XIX, 10, pp 991-993
(Fly tox)
See No 218a
(Gierson's spray)
- 495 BARRETO, J DE B. (1929) Destruction of insects by spraying with
insecticides in work against malaria
Sci Med, VII, 11, pp 527-532
RAEB, XVIII, 4, p 72

- 496 BARRETO J DE B, and PFRYASSU A G (1929) The spraying of insecticides in the prophylaxis of yellow fever *Brasil Medico* XLIII 13 pp 320-329 *PAB* XVII p. 203
- 497 BRUG S L and SLOOTEN J VAN (1927) Report of some experiments on the efficiency of insecticides of 'Flit', 'Lils' and of some other preparations *Nederl Volk Vet Ind* XVI 3 pp 514-530
- 498 FUGLINO M (1931) Malaria control Observations and researches in the Albanian malaria region *Revue Arch Schiffs Trop Hyg.* XXV 1 p 63
(Formalin lysol cresol)
- 499 FRAGA C (1928) The yellow fever epidemic at Rio de Janeiro *U S A P b Hlth Repts* XLIII 47 pp 3079-3085
(kerosene with carbon tetrachloride and methyl salicylate operated by electric motors using compressed air)
- 500 GIEMSA G (1911) Beitrag zur Frage der Stechmückenbekämpfung *Arch Schiffs Trop Hyg* XV p 533
- 501 GIEMSA G (1912) Über die Vernichtung der Stechmücken mit Hilfe des Sprayverfahrens *Ibid* XVI p 565
- 502 GIEMSA G (1913) Spraying as a means of preventing malaria and other diseases carried by biting flies *Ibid* XVII 6 pp 181-190 *PAB* I p 73
- 503 GIEMSA G (1914) Improvements in spraying methods against mosquitoes *Ibid* XVIII 1, pp 26-29 *PAB* II p 80
- 504 LEFROY H M (1919) Fly sprays *Trans Roy Soc Trop Med and Hyg.* XIII 1 pp 1-9
- 505 MALARIA COMMISSION, HOLLAND (1925) Report for 1924 *PAB* XV p 20
- 506 MANVELL P A. (1929) Some answers to the cantonment anti malarial problem *Jl R A M C* LIII 3 pp 196-201
- 507 MANVELL, P A (1930) Note on experiments towards the production of a cheap insecticide for anti mosquito work *Ibid*. LIV 2 pp 123-127
- 508 RICHMOND A E and MENDIS J C (1930) See No 239
- 509 STEINDEL (1911) *Arch Schiffs Trop Hyg* XV, p 121
(Called attention to neglect of spraying in malaria prevention)
- SWELLENGREBEL, N H et al. See No 393
(1929)

- 484 ORENSTEIN, A J (1914) . The technique of mosquito screening
Arch *Schiffs Trop Hyg*, XVIII 1,
pp 16-21
PRINCE J A LE, and See No 34 pp 198-199
ORENSTEIN, A J (1916)
REITLER R (1930) See No 247
ROSS P (1911) See No 37, p 261
- 486 RUTHERFORD N J C (1928) Mosquito proofing of barracks of British
troops in India *Jl P A M C*
L 1, pp 76-78 and 4 pp 316-318
- 487 SBACCHI (1906) Prophylaxis by wire gauze *Atti della*
Soc per gli Studi della Malaria
See No 373
- 488 SCHUCKMAN, W VON (1926) On the control of flies and mosquitoes
Zeitsch Ang Ent, XII, 2, pp 332-339
RAEB, XV, pp 12-13
- 489 SIMPSON W J (1917) The vixorant mosquito and fly proof head
gear Invented by Mrs W J Simpson
Trans Roy Soc Trop Med & Hyg
X, 7, p 140
- 490 SNIDOW, H W (1919) Screening of residences *U S A Pub*
Health Bull No 104, p 19
VISBECQ and LACAZE, H See No 43, pp 215-243
(1924)
WATSON, M (1915) See No 44, p 179
WATSON, M (1921) See No 45, pp 257-263, 364
- 491 WHYTE, G D (1910) The effect of a mosquito net on the
air within it (based on 80 analyses)
Brit Med Jl, II of 1910, pp 972-973
- 492 WURTZ R (1907) Sibiçtage des tulles employes pour protéger
contre la piqure des moustiques *Reu*
Med Hyg Trop, IV, 4, p 181 In
ROSS, 1911 (see No 37)
- 493 ZIEMANN, H (1906) In MENDEL, *Hand der Tropenkrank*, III
p. 445

Sprays.

- 494 ABBATUCCI, S, and Expériences sur un liquide insecticide
POURBAUD, E (1926) commercial à base de pétrole et de
poudre de pyrèthre *Bull Soc Path*
Exot, XIX, 10 pp 991-993
(Fly tox)
BALFOUR, A (1913) See No 218a
(Giemsa's spray)
- 495 BARRETO, J DE B (1929) Destruction of insects by spraying with
insecticides in work against malaria
Sci Med, VII, 11, pp 527-532
RAEB, XVIII, 4, p 72

- 496 BARRERO J DE B and PERYASSU A G (1909) The spraying of insecticides in the prophylaxis of yellow fever *Basil Med o* XLIII 13 pp 300-309 *PAEB* XVII p 253
- 497 BRUGS L and SLOOTEN J VAN (1917) Report of some experiments on the efficiency as insecticides of Flit R 14 and of some other preparations *Meded Volk Ved Ind* XVI 3 pp 54-530
- 498 FUGLING M (1911) Malaria control Observations and researches in the Albanian malaria region *Beihfte Arch Schffe Trop Hyg* XXV 1 p 63
(Formalin lysol cresol)
- 499 FRAGA C (1909) The yellow fever epidemic at Rio de Janeiro *U S A P b Hlth Repts* XLIII 47 pp 309-308;
(Kerosene with carbon tetrachloride and methyl salicylate operated by electric motors using compressed air)
- 500 GIEMSA G (1911) Beitrage zur Frae der Stechmuckenbekampfung *Arch Schffe Trop Hyg* XV p 533
- 501 GIEMSA G (1912) Ueber die Vernichtung der Stechmucken mit Hilfe des Sprayverfahrens *Ibid* XVI 1 p 565
- 502 GIEMSA G (1913) Spraying as a means of preventing malaria and other diseases carried by biting flies *Ibid* XVII 6 pp 181-190
PAEB I p 70
- 503 GIEMSA G (1914) Improvements in spraying methods against mosquitoes *Ibid* XVIII 1 pp 28-29 *R4FB* II p 80
- 504 IMPROY H M (1910) Fly sprays *Trans Roy Soc Trop Med and Hyg* XIII 1 pp 1-9
- 505 MALARA COMMISSION HOLLAND (1903) Report for 1904 *P4EB* XV p 21
- 506 MANSFELL P A (1909) Some answers to the cantonment anti-malarial problem *Jl R A M C* LIII 3 pp 196-201
- 507 MANSFELL P A (1930) Note on experiments towards the production of a cheap insecticide for anti-mosquito work *Ibid* LIV 2 pp 123-127
R1 20ND A L and MENDEL J C (1930) *Ses* No 238
- 508 STENDEL (1911) *Arch Schffe Trop Hyg* XV p 121
(Called attention to neglect of spraying in malaria prevention)
- SWELLENGREBEL N H *et al* *Ses* No 393
(1909)

- 509 THOMPSON, T O (1928) .. The cantonment anti malarial problem
Jl R A M C. LI, 3, pp 179-185

Traps

- BALFOUR, A (1913) See No 218a
 (Box trap)
- 509a BATH, C H (1914) Insect trap *Canal Record, Ancon*, VII,
 25 pp 239-240 *RAEB*, II, p 161
- 509b BATH, C H (1931) . The practical and research value of mos-
 quito traps *Amer Jl Trop Med*,
 XI, 2, pp 147-150
 (Traps on screened ventilators)
- 510 BLIN, G (1908) Destruction des moustiques par le procede
 des trous pièges *Bull Soc Path Exot*,
 I, pp 100-103
- 511 FLETCHER T B (1912) Traps for mosquitoes *Procs 4th Entom.*
Mtg, Pusa, pp 219-221
- 512 HEADLEE, T J (1929) The development of mechanical appa-
 ratus for detecting the presence of mos-
 quitoes in various localities *Procs 15th*
Ann Mtg N Jers Mosq Ext Ass,
 pp 160-168
 (Illuminated traps)
- JAMES, S P (1914) See No 273
- 513 LEBROY H M (1907) A preliminary account of the biting flies of
 India, pp 12-15 Govt Press, Calcutta
- LEGBOUX, R (1917) See No 230
 (Sergeant's trap)
- 514 LOFTIN, U C (1920) Mosquitoes found about Gainesville, Fla.
 Part II Traps for mosquitoes *Florida*
Buglist Gainesville, III, 4, pp 53-59,
 and 67-71 *RAEB*, VIII, p 158
- 515 MCHARDY, J W (1930) Report of the Government Entomologist
Ann Rept Med Lab, Dar es Salaam
 for 1928, pp 41-52
 (Mosquito trap house, with Le Prince
 and Orenstein's screen traps)
- 516 MEYER W (1920) On the possibilities of using mosquito traps
 in anti malaria work *U S J Pub*
Health Rpts, XXXV, 34 pp 1974-1977
- 517 NUTTALL G H F. and SHIPLEY A G (1903) Studies in relation to malaria Part II
 The structure and biology of Anopheles
Jl Hyg, II, 1, p 73
- 518 OWEN W O (1919) Illuminated trap for night flying insects
New York Med Jl, CIV, 14 p 590
- PRINCE J A LE, and ORENSTEIN, A J (1916) See No 34
 (Traps on screens)
- RICHTON, A E, and MENDIS I C (1930) See No 238
 (Crinoline trap, box traps, wire traps)

- 519 STRICKLAND, C and CHOUHURA, K L On trapping adult mosquitoes *Ind Jl Med Res*, XVII, 4 pp 1009-1014 (1930)
- 520 WILSON, C J (1930) Annual Report of the Malaria Advisory Board, F M S for the year 1929 Suppl to F M S Govt Gaz., Aug 29th
- 521 ZETEK, J (1916) Reducing malaria by reducing the number of Anopheles within buildings *Ann Ent Soc America*, IX, 3, pp 275-283 *R4EE*, V, pp 1-2

Zoophylaxis

- 521a BARBER M A and HAYNE T B (1924) Some notes on the relation of domestic animals to Anopheles *U S A Pub Hlth Rpts*, XXXIX 4 pp 143-144
- 522 BOREL E (1926) Résultats d'une enquête épidémiologique et entomologique à Yaback (Annam) *Bull Soc Path Exot*, XIX, 9, pp 845-852
- 523 BOREL E (1926) La constitution du sol et le paludisme en Cochinchine *Ibid*, 10, pp 930-942
- 524 BOREL E (1927) Résultats d'une enquête malarologique à Dalat (Cochinchine) *Ibid*, XX, 6, pp 427-434
- 525 BUCK A DE (1926) Variations in *A maculipennis* in connection with Anophelism without malaria *Dissertation, Univ Amsterdam*
- 526 BUCK, A DE SWELLEN ORRELL N H, and SCHONTE, N H (1927) Studies on anophelism without malaria *Bull Ent Res* XVII, 4 pp 351-371
- BULL, C G and KIRBY W V (1931) See No 544
- 527 BULL C G LOOT I M and FETKOLDS, B D (1933-34) Preferential feeding experiments with Anopheline mosquitoes Parts I and II *Amer Jl Hyg*, III 5 pp 514-520, and IV, 2 pp 109-118
- 528 CRAIG F W (1933) The zoophilism of *Anopheles* in relation to the epidemiology of malaria in India *Ind Jl Med Res*, X, 4, pp 962-964
- 529 DARLING S T (1935) Entomological research in malaria *South Med Jl*, XVIII 6 pp 446-449 and 452-453
- 530 DAVIS N C and SHANNON, R C (1933) The blood feeding habits of *A pseudo punctipennis* in Northern Argentine *Amer Jl Trop Med*, VIII 5 pp 443-447

- 531 DOORENOS, W B (1915) . Species sanitation and diversion to animals as measures to control malaria in the tropics *M D Thesis, Univ Leyden*
- 532 FACE, L LA, and SELLA, M (1937) Zooprophylaxis in malaria control according to the concept of Prof B Grassi *Riv Mal*, VI, 6, pp 935-943 Eng Summ, pp 1011-1012
- 533 FALLERONI, D (1926) The Italian Anopheline fauna and its habitat Methods of work against malaria *Ibid*, V, 5-6, pp 553-593
- 534 FALLERONI, D (1927) The solution of the malaria problem in Italy *Ibid*, VI, 2, pp 344-409 Eng Summ, p 493
- 535 FALLERONI, D (1927) A discussion on zooprophylaxis and on the biology of Italian Anopheles *Ibid*, 4-5, pp 751-756 Eng Summ p 879
- 536 FALLERONI, D (1929) A discussion on zooprophylaxis and other measures against malaria (4th Note) *Ibid*, VIII, 3, pp 315-326
- 537 FALLERONI, D (1929) Discussion on zooprophylaxis Principles for constructing rural buildings in malarial regions *Ibid*, 5, pp 590-606 Eng Summ, p 634
- 538 FERMI C (1928) Anti malarial zooprophylaxis *Malaria logia, Ser III*, pp 6-9
- 539 FRENKEL, H S (1926) *Ned Tijds Hyg, Microb & Serol*, I (Early precipitation tests)
- 540 FRY, A B (1922) The role of cattle in the epidemiology of malaria *Ind Med Gaz*, LVII, 1 pp 1-2
- 541 GRASSI B (1922) Domestic animals and malaria A new horizon in anti-malarial work *Ann d Igiene*, XXVII 6 pp 491-512 *BALB*, X, pp 202-203
- HACKETT, L W, and MISSIROLI, A (1931) See No 601
- 542 IVANIC (1926) *L'anophele maculipennis* est il homophile? *Belgrade Impr Drag Gregoric*, 8 pp In SWELLENGREBEL, BUCK and SCHOUTE 1929 (see No 564)
- 543 IYENGAR, M O T. (1924) The importance of animal prophylaxis in malaria (19 refs) *Proc 5th Fatom Mtg, Pusa*, 1923, pp 204-214
- 543a KING, W V, and BULL, C G (1923) The identification of the blood meal of mosquitoes by means of the precipitation test *Amer. J Hyg*, III, 5, pp 491-496

- 544 KING W V., and BULL, C G (1923) The blood feeding habits of malaria carrying mosquitoes *Amer Jt Hyg.* III 5, pp 497-513
- 545 KILGLEY I J., and LIEBMAN E (1938) Studies on malaria in an uncontrolled hyperendemic area Part II *Jt Prev Med.* II 5, pp 433-440
- 546 LEGENDRE, J (1929) Role du bétail et de la basse cour dans la défense contre la malaria *C R Hebd Acad Sci.* CLXX, 12 pp 766-769 *RAEB.* VIII, p 98
- 547 LEGENDRE J (1929) La concurrence entre moustiques zoophiles et anthropophiles *C R Acad Sci Fr* CLXXXVIII, 1 pp 95-97 *RAEB.* VIII p 80
- 548 MARCHOUY E. (1927) Le paludisme dans les Dombes et en Camargue *Full Acad Med* p 67 *RAEB* XV, p 137
- 549 MARTINI, E (1923) A criticism of the theory of misanthropism or zoophilous Anophelines *Arch Schiffs Trop Hyg.* XXVI, 9 pp 257-265 *RAEB* V, pp 223-229
- 550 MISSIROLI, A and HACKETT I W (1927) The spontaneous retrogression of malaria in some regions of Italy *Riv Mal.* VI 2 pp 193-243
- 551 MISSIROLI A., and HACKETT L W (1929) The precipitin test as a means of determining the source of Anopheline blood meal (MS) *League of Nations Hlth Organ.* CH Mal 131
- 552 PRADO A (1929) Zootropismo des Anopheles *Sci Med* VII 8 pp 379-381 *RAEB* XVII 12, p 236
- 553 RIZZI M (1919) A new form of anti malaria prophylaxis in *d'Igiene* XXIX, 11 pp 743-751 *RAEB* IX p 131
- 554 ROUBAUD E (1919) Antagonisme du bétail et du l'homme dans la nutrition sanguine de l'*Anopheles maculipennis* Le rôle anti paludique du bétail domestique *Bull Agric Alger Tun Maroc.* XXV, 11, pp 272-273 *RAEB.* VIII p 33
- 555 ROUBAUD E (1920) Les conditions de nutrition des Anophèles en France, et le rôle du bétail dans la prophylaxie du paludisme *Ann Inst Past.* XXXIV 4, pp 181-228.
- 556 ROUBAUD E (1921) La différenciation des races zootropiques d'anophèles et la régression spontanée du paludisme *Bull Soc Path Exot.* XIV, 9 pp. 577-595

- 557 ROUBAUD, E (1922) A propos des races zoophiles d'Anophèles *Bull Soc Path Exot*, XI, 1 pp 36-39
- 558 ROUBAUD, E (1925) Peuplements humains et peuplements animaux comme facteurs économiques comparés de la déviation anophéenne antipalustre *Ibid*, XVIII 9, pp 73-730
- 559 ROUBAUD, E (1928) Nouvelles recherches sur l'évolution zoophile des faunes d'Anophèles en Europe, d'après les données de l'armement maxillaire *Ann Inst Past*, XLII 5, pp 553-618 (38 refs)
- 560 ROUBAUD, E (1930) Quelques remarques à propos de l'interprétation théorique des index maxillaires *Bull Soc Path Exot*, XXIII 1, pp 47-53
- 561 ROUKHADZE, N (1927, 1928) Stabbing of animals as a means of malaria reduction *Ibid*, XIX, 2, pp 100-105, XX, 2 pp 153-170
- 561a ROUKHADZE, N (1929) Rôle of domestic animals in the prophylaxis of malaria *Arch Trop Med* II 2, pp 84-88 *RAEB*, XVII, p 107
- 562 SWELLENGREBEL, N H (1925) Considérations sur la prophylaxie du paludisme par la déviation des Anophèles *Ibid* XVIII 9, pp 715-727
- 563 SWELLENGREBEL, N H (1929) The dissociation of the sexual and nutritive functions—Gonotrophic Dissociation—in *A. maculipennis* as the cause of malaria in the Low Countries and its connection with house infection *Ann Inst Past*, XLIII 10, pp 1370-1389 *RAEB*, XVIII, pp 53-54
- 564 SWELLENGREBEL, N H, BUCK, A DE and SCHOUTE E (1927-29) On Anophelism without malaria around Amsterdam *Proc Roy Acad Sci, Amsterdam*, XXX, 1, pp 61-68, XXXI, 4-5, pp 531-539, XXXII 6, pp 772-779
- 565 SYMES, C B (1930) Anophelines in Kenya *Kenya and E Afric Med J*, VII, 1, pp 2-11
- 566 TREVSZ, F (1930) Recherches sur l'index maxillaire d'*Anopheles maculipennis* *Bull Soc Path Exot*, XXIII, 1, pp 43-47
- 567 TREVSZ, F (1930) L'index maxillaire d'*Anopheles maculipennis* et la théorie du zootropisme anophélien *Arch Inst Past d'Algérie*, VIII 1, pp 1-70 (75 refs)

- 567a. WALCH, E. W. (1928) Some remarks on malaria in the Dutch East Indies *South Med J.*, XVIII, 6, pp. 434-438
- 568 WALCH, E. W. and SARDJITO, M. (1928) Identification of the blood meal of Netherlands Indian *Anopheles* by means of the blood test *Nederl Volk Ned Ind.*, XVII, 2, pp. 234-250
- 569 WEIDANZ (1909) *Cent f Bakt.* XLII, Beihefte, p 108 (Early work with precipitation test for blood)
- 570 WESSENBRO-LUND, C. Contributions to the biology of Danish Culicidæ. *Mem Acad Sci & Letters Copenhagen*, VII, 1, pp 1-210
- 570a WESSENBRO-LUND, C. Reasons for the change in feeding habits of *A. maculipennis* *C R Soc Biol.*, LXXXV, 28, pp. 383-386 *RAEB.*, IX, p 153

ADDITIONS TO BIBLIOGRAPHY.*

(1931-1940.)

A. GENERAL.

- 571 AHMED, N (1939) Five years of anti malaria work at Barwadih Railway Settlement. *Ind Med Gaz*, LXXIV, 8, pp 472-476
TDB, XXVII, p 364.
- 572 BALFOUR, M C (1936) Some features of malaria in Greece and experience in its control *Riv. Mal.*, XV, 2, pp 114-131 *RAEB*, XXIV, p 163
- 573 BANNER, M A (1932) Malaria control work in West Africa. *South Med J.*, XXV, 6, pp 649-651
- 574 BECKMAN, H (1940) The prophylaxis of malaria *Ibid.*, XXXIII, 5, pp. 616-622
- 575 BOYD, I E M (1932) Malaria in India *Jl R A. M C*, LVIII, 2-4, pp 81-91, 177-194, 248-261
RAEB, XX, p 122
- 576 BURCA, B DE (1939) Note on anti-malaria measures in Quetta Cantonment during 1938 *J Mal Inst Ind*, II, 2, pp 121-130
- 577 CHAMBERLAIN, W P (1933) The medical activities of the Panama Canal *Med Surg*, LXXII, 2, pp 108-124
- 578 CHOWDHURY, K L (1936) Three years' mosquito control work in Calcutta *Rec Mal Surv Ind* VI, 3, pp 467-481
- 579 CLARKSON, L M (1940) Engineering phases of malaria control *South Med J.*, XXXIII, 8, pp 900-904
- 580 COVELL, G (1940) Anti-mosquito measures, with special reference to India *Govt India Hlth Bull*, No 11, Pd 5 58 pp
581. COVELL, G, and AFRIDI, M K (1939) Anti malaria operations in Delhi: Part II. *J Mal Inst Ind*, II, 4, pp 315-340
- 582 CRAIG, W J F. (1936) Malaria prevention in Java *Jl R. A. M C*, LXVII, 4 pp 233-245 *RAEB*, XXV, p. 31

* *RAEB* = *Review of Applied Entomology*, Series B
TDB = *Tropical Diseases Bulletin*

- 583 EARLE W C, *et al* (1936) Methods used to control malaria in Puerto Rico *Puerto Rico J Pub Hlth*, XI, 3 pp 434-456 *RAEB*, XXIV, p 141
- 584 GIES, R W (1936) Mosquito control engineering III Control technique and organization *Engng News Rec* Aug 13, pp 225-238
- 585 HACKETT, L W (1931) Recent developments in the control of malaria in Italy *South Med J*, XXIV, 5, pp 426-430
- 586 HARRISON, C C, and RAMSAY G C (1933) Malaria survey of tea estates in the Eastern Duars Bengal, and malaria control recommendations *J Trop Med Hyg*, XXVI 3 pp 33-41 *RAEB* XXI p 76
- 586½ HERNS W B, and GRAY H F (1940) Mosquito control 317 pp New York
- 587 HOWARD L O (1936) Mosquito control engineering I The growth and importance of anti mosquito work *Engng News Rec* July 23, pp 118-122
- 588 HOWARD L O and BISHOPP F C (1937) Mosquito remedies and preventives *U S A Dept Agric Farmers' Bull* No 1570 11 pp *RAEB*, XXI, p 33
- 589 HOWARD S C, *et al* (1939) Anti malarial measures on estates Control of urban and rural malaria in Malaya *Bull Inst Med Res F M S* No 2 23 pp
- 590 HUSSAMUDDIN (1931) The anti malaria campaign in Turkey *C R 2nd Cong Int Pul Alger*, 1930 II pp 359-401 *RAEB* XV, p 74
- 591 MACKAY R (1939) Second (final) report of the malaria unit, Dar es salaam Govt Press Dar es salaam 61 pp
- 592 MALARIA ADVISORY BOARD F M S (1931-40) Annual Reports Govt Press, Kuala Lumpur
- 593 MANSON D, and RAMSAY G C (1933) Further observations on a malaria survey in Jorhat District Assam *Rec Med Surv Ind*, III, 3 pp 479-494
- 594 MATHESON R (1937) Medical entomology, pp 274-298 London
- 595 MISHNAEVSII M N (1937) The malaria mosquito and measures for its control (in Russian) 214 pp *RAEB*, XXV, p 206
- 596 MISSIROLI A (1931) Lectures on the epidemiology and prevention of malaria 531 pp Rome

- 597 MULLIGAN, H W, and The prevention of malaria incidental to
AFRIDI M K (1938) engineering construction *Govt Ind
Hlth Bull*, No 25 *PAEB*, XXVII
p 184
- 598 NEZITCH E (1934) Malaria in Dalmatia *Bull Off Int Hyg
Pub*, XXVI, 3, pp 463-473 *R4EB*,
XXII, p 117
- 599 PRINCE, J A LE (1931) Cost of malaria control *Amer J Pub
Hlth*, XXI 4, pp 378-391 *R4EB*,
XX, p 2
- 600 RANKOV, M (1934) Malaria in Southern Serbia *Bull Off
Int Hyg Pub*, XXVI 3 pp 474-499
R4EB, XXII, p 118
- 601 RAO B A. (1939) Malaria control in Malaya Java Ceylon
and S India 66 pp Govt Press
Bangalore
- 602 RICHARDS, A G (1938) Mosquito control on Long Island, New
York *Bull N Y State Museum*,
No 316, pp 85-180 *RAEB*, XXVIII,
p 60
- 603 ROSAI, G (1935) How Maccarese was fully reclaimed in
spite of malaria *Ann R Inst Sup
Agrar Portica* (3) VII, pp 163-379
R4EB, XXV, p 68
- 604 RUSSELL P F (1933) Malaria in the Philippine Islands *Amer
J Trop Med* XIII 2 1p 167-178
R4EB XXI, p 148
- 605 SCHARFF J W (1932) A note on Public Health Administration
in the Northern Settlement 35 pp
Singapore
- 606 SCHARFF, J W (1935) Anti mosquito measures in the Northern
Settlement of Malaya *Trans 9th Cong
F F A T M*, II, pp 201-212 *RAEB*,
XXVI, p 127.
- 607 SENIOR WHITE, R A, and Studies in malaria as it affects Indian
NEWMAN, C D (1932) Railways Part II *Tech pap Rly
Board India* No 258, 71 pp Govt
Press, Calcutta
- 608 SERGIEV P G, and KOV Organization of anti malaria work in
TUN A S (1937) the U S S R *Med Paras*, VI,
6, pp 723-765 *R4EB*, XXVI,
p 237
- 609 SHAPIRO, J M (1933) Rural malaria control in Palestine *J
Egypt Med Assoc*, XVI, 8 pp 850-857
RAEB, XXII, p 238
- 610 SILER, J F (1931-1934) *Ann Rpts Hlth Dept Panama Canal*
- 11 STROTHER SMITH F F (1938) Anti-mosquito measures in Cantonments.
Ind Med Gaz, LXXIII, 10 pp 606-
608 *RAEB*, XXII, p 81

- 612 SWELLENOREBEL N H and BUCK A DF (1933) Malaria in the Netherlands 267 pp Amsterdam and London *RAEB* XXVII p 27
- 613 VICKERS W J (1934) Recent local advances in the economics of practical malaria control. *Malay Med J* IX 2 pp 40-43 *RAEB* XXII p 191
- 614 WALCZ E W and SOESILO R (1935) Malaria control in the Netherlands Indies. *Med Volk Med Ind* XXIV 3 pp 88-94 *PAEB* XXIV p 16
- 614a WILLIAMS L L (1931) Rural malaria control. *South Med J* XXIV 5 pp 440-442
- 615 WILLIAMS L L (1936) Mosquito control engineering VI Effective malaria control. *Engng News Pac* Sept 3 pp 341-343
- 616 WILSON D B (1938) Report of the malaria unit Moshu Govt Press Dar es Salaam 34 pp

B SPECIAL SUBJECTS

Biological Control

- 617 ALVARADO C A (1939) Anti malaria measures in Argentina. *Bol San t Dep Ac H g Argent* III 1 pp 891-915 *RAEB* XXVII p 247 (Art fic al shad ng)
- 618 AMBLAET R (1935) An attempt to destroy *Anopheles* larvae by Williamson's method (herbage cover). *Arch Inst Past Alg r* XVI 2 pp 161-165 *RAEB* XXVII p 33
- 619 ANANYAN S A (1939) Interrupted irrigation of rice fields in Armenia. *Trop Med Let* VII 1 pp 47-55 VIII 8-9 pp 39-47 *RAEB* XXVII p 117 XIX p 154
- 620 ANTOINE M (1936) Prevention of rural malaria by intermittent irrigation of rice fields. *Bull Soc Med Ch r Indoch ne* XIV 6 pp 612-615 *TDB* XXIV p 208
- 621 BAYERJEE A C (1939) The Guntur Weir Lucknow. *J Mal Inst Ind* II 3 pp 257-259 (Shading)
- 622 BARBER M A and RICE J B (1935) The relation of housing to malaria in certain villages of East Macedonia. *Amer J Hyg* XXII 3 pt 512-538 *RAEB* XXIV p 35
- 623 BARROWMAN B (1939) Replantation and malaria. *J Malay Br E t Med Ass* III 2 pp 170-174 *RAEB* XXVIII p 63 (Shading jungle-clearing)

- 623a BEATTIE M V F (1932) Physico chemical factors in relation to mosquito breeding *Bull Ent Res.* XXIII, 4, pp 477-496
- 624 BINYUKOV, V I (1938) Dynamic action of water as a means of combating Anopheles *Med Paras* VII 5 pp 701-715 *R4FB* XXVII p 173
- 625 BISHOP E L (1936) Malaria control activities of the Tennessee Valley Authority *U S A Pub Hlth Rpts*, LI 29 pp 970-975 *TDB* XXXIV, p 142
(Fluctuating water level)
- 626 BLACKLOCK D B (1936) Cultivation of dense shade plants as an anti malaria measure *Ann Trop Med Paras*, XXV, 2, pp 181-186 *RAEB*, XXIV, p 277
- 627 BLACKLOCK D B (1939) Notes on siphon action, with special reference to anti mosquito work *Ibid* XXXIII 2 pp 141-160 *RAEB* XXVIII, p 16
- 628 BOER H S DE (1936) Malaria control in swamps *Quart Bull Hlth Org L of N*, V, 1 pp 139 139 *RAEB*, XXIV p 153
(Planting of trees to dry up land)
- 629 BRUNELLI G (1933) The surrounding channel and hydro biological measures against malaria *Att. R Acad Na Lincei* XVII 8 pp 671-674 *R4FB*, XXI p 219
(Method of dealing with fish ponds)
- 630 BUXTON, F A (1934) Chemical factors affecting the breeding of Anopheles in Trinidad *Bull Ent Res.* XXV, 4, pp 491-494 *RAEB* XXIII p 79
(Ammonia nitrogen content, hydrogen ion concentration carbon dioxide dissolved oxygen nitrites, nitrates phosphates chlorides, total solids)
- 631 CARE H P (1938) *A. albimanus* breeding in relation to degree of shade *South Med J*, XXXI 7 pp 804 805 *RAEB* XXVII p 30
- 632 CHRISTOPHERS, S R, and MISSIROLI A (1933) Report on housing, and malaria *Quart B H Hlth Org, I of N*, II pp 355-482 *RAEB*, XXII p 147
- 633 COLLIGNON, E (1938) A great anti larval measure The draining of Lake Halloula *Arch Inst Past Alger*, XVI, 2 pp 166-175 *RAEB* XXVII p 34

- 634 COVELL G (1940) Malaria problems in the Wynaad *Planters Chronicle* XXX 15 pp 309-310
(Herbage packing)
- 635 COVELL G and HARBHAG WAT (1939) Malaria in the Wynaad South India *J Mal Inst Ind* II 4 pp 341-376
(Herbage packing)
- 636 CROSS D (1933) Observations on malarial prevention in estates *Malay Med J* VIII 1 pp 261-274 *RAEB* XXII p 66
(Breeding places sprayed with cow dung solution)
- 637 CUNY T A (1939) Flood flushing schemes in Bengal *Proc Nat Inst Sci Ind* IV 2 pp 215-218
RAEB XXVII p 7
- 638 DRYENSKI K (1936) Influence of intermittent irrigation of rice-fields on malaria and the rice crops *Mitt Bulg Ent Ges* IX pp 11-14
PAEB XVI p 103
- 639 EBERSTEIN A (1935) Biological control of anopheline vectors in the Philippines *J Phil Isl Med Ass* XV 4 pp 177-191
- 640 EBERSTEIN A (1936) Present orientation in the control of malaria in tropical countries *Ibid* XVI 2 pp 61-71
- 641 EBERSTEIN A (1938) Biological control of *Anopheles minimus* sub-group breeding in the Philippines *Ibid* XVIII 7 pp 415-430 *RAEB* XXVII p 253
(Training of streams stagnating by dams)
- 642 FAKOLOPOV S K (1931) Interrupted irrigation of rice fields *Trop Med Vet* IX 2 pp 58-63 *RAEB* XVI p 114
- 643 FAYO W B R (1937) Successful malaria control at Banjoewangi *Med Volk Ned Ind* XXI 2 pp 41-60
RAEB XX p 218
(Biological control of fish ponds)
- 644 FERMI C (1938) Shading as an anti-malarial measure for anopheline areas *Malariaologia* Jan 31 pp 60-68
- 645 FERMI C (1933) The biological clearing of aquatic vegetation *R Mal* XII 3 pp 523-531
RAEB XVI p 260
- 646 GRIFFIN F C (1931) Anti-malarial engineering *Sonar Bangla* VI 10 pp 49-51 11 pp 53-56
(Flood flushing)

- 647 GRIFFIN, F C (1933) .. Note on the evils of land reclamation by marginal embankments in tidal areas 8 pp Govt Press Calcutta
(Flood flushing)
- 648 HACKETT, L W (1936) Biological methods in anti-larval work. *Riv Mal*, XIV, 6 Suppt., pp 58-70
RAEB, XXIV, p 114
- 649 HACKETT L W (1936) Biological factors in malaria control. *Amer J Trop Med*, XVI 3 pp 341-352
RAEB, XXIV, p 222
- 650 HACKETT, L W (1937) Malaria in Europe 336 pp London
RAEB, XXV, p 114
- 651 HACKETT, L W, and MISSIROLI A (1931) The natural disappearance of malaria in certain regions of Europe. *Amer J Hyg*, XIII 1, pp 57-78
RAEB, XIV, p 107
- 652 HACKETT, L W, *et al* (1938) The present use of naturalistic measures in the control of malaria. *Bull Hlth Org L of N*, VII, 6 pp 1016-1064
RAEB, XXVII, p 211
- 653 HARGREAVES, H (1933) Ann Rept Govt Entomologist Ann Med San Rept Uganda 1932 pp 86-90
RAEB, XXII, p 83
(Pollution of breeding places by cut elephant grass)
- 654 HILL R B (1937) The malaria problem in Portugal. *South Med J*, XXX 9, pp 953-956
RAEB, XXVI p 217
(Intermittent drying of rice fields)
- 655 HINMAN E H (1938) Biological effects of fluctuation of water level on anopheline breeding. *Amer J Trop Med*, XVIII, 5 pp 483-495
TDB, XXVI p 334
- 656 HINMAN, E H, and HURT BUT, H S (1940) The relation of shade to *A. quadrimaculatus* breeding. *J Parasit* XXVI, 2, pp 145-156
- 657 HODGKIN, E P (1938) Naturalistic methods of malaria control. *J Malay Br Brit Med Ass*, II, 1, pp 24-29
RAEB, XXVI, p 230
- 658 JESUS, P I DE (1933) Hydrodynamics of mosquito breeding places. *Amer J Hyg*, XVII, 2, pp 502-514
RAEB, XXI, p 147
- 659 KIEER, C C, and STROM QUIST W Q (1939) Importance of reservoir preparation in mosquito control of impounded water. *Proc 21st Mtg Nat Mal Comm* 1938, pp 60-71 (Typescript)

- 660 BIRK, J B (1937) The relation of shade to the natural breeding places of *A. costalis* in Mauritius *Ann Trop Med Paras.* XXXI, 1, pp 1-6 *RAEB*, XXV, p 180
- 661 KONSULOFF, S (1922) The control of malaria mosquitoes in rice-fields *Z Angew Ent*, VIII, 2, pp 283-294 *RAEB*, X, p 193 (Intermittent drying)
- 662 KONSULOFF, S (1933) Irrigation and malaria *Ibid*, XX, 3, pp 461-463 *RAEB*, XXII, p 52 (Intermittent drying of rice-fields)
- 663 LAZUK, A. D and UTENKOV, I V (1939) Drying out of mill ponds as an anti larval measure *Med Paras*, VII, 6, pp 847-859 *RAEB*, XXVII, p 180
- 664 LEGWEN, W. A., and KIRBY, A J (1939) The practical application of pH on larvicidal operations *Proc 21st Mtg Nat Mal Comm* 1938, pp 72-96 (Typescript)
- 665 MACDONALD, G (1939) A design of flushing siphon for control of anopheline breeding *J Mal Inst Ind*, II, 1, pp 63-69
- 666 MAXSON D (1936) Velocity, silt and larval drift *Ind Med Gaz*, LXVI, 5, pp 270-275 *RAEB*, XXIV, p 204
- 666a MANSON, D, and RAMSAY G C (1933) Malaria survey in Jorhat, Assam *Rec. Mal Sure Ind*, III 3, pp 479-494 (Shading)
- 667 MARTINI E (1934) Salinity of water collections and malaria *Ent Brish Berl*, I, pp 28-44 *RAEB*, XXII, p 202
- 668 MAURICE, A (1931) The employment of a rodent in anti malarial work *Asiatr Med*, July August Abstr in *Piv Mal*, XI, 4, pp 533-538 (Chilean beaver feeds on aquatic plants)
- 669 MEASHAM J E (1939) The transmission and prevention of malaria on the tea estates of the Madras Presidency 56 pp Valparai (Sluicing, shading)
- 670 MONDAL, P S (1940) The species control of a hill stream by training *J Mal Inst Ind*, III, 4, pp 463-474
- 671 MORIN, H G S (1934) First results of agricultural measures in the malaria control campaign in Indo-China *Bull Soc Mal Chir Indochine*, XII, 2, pp 230-244 *RAEB*, XXIII, p 26

- 672 MORIN H G S (1934) Destruction of larvae in certain streams
Bull Soc Méd Chir Indochine XII,
8 pp 743-746 *RAEB* XXIII p 93
(Flushing)
- 672a MORIN, H G S and BADER H (1933) Physico chemical factors in the ecology of
Anopheles larvae *Ann Inst Past*
LI 5 pp 656-668
- 673 MULDER J G A (1936) The campaign against malaria at Tjaling
on the west coast of Atjeh *Genesal*
Tijd Ned Ind LXXVI 30 pp 1864-
1889 *TDB* XXXIV p 149
(Lagoons and swamps connected with
the sea)
- 674 MURPHY R A (1934) Anti malaria work on a group of tea
estates in South Sylhet *Ind Med*
Gaz LXIX 8 pp 437-439 *RAEB*
XXII p 290
(Siltng shading)
- 675 NICHOLAS W A (1937) Anti malaria measures in Java *J Malay*
Br Brit Med Ass I 1 pp 38-50
RAEB XXV, p 261
(Control of fish ponds, intermittent
drying shading)
- 676 NICHOLAS W A (1939) A four day automatic sluice *Ibid*
III 1 pp 44-46 *TDB* XXXVIII
p 293
- 677 OVERBEEK, J G, and STOKER W J (1937) Malaria in the Netherlands Indies and
its control 60 pp Batavia
(Control of fish ponds intermittent
drying shading)
- 678 OVERBEEK J G, and STOKER W J (1938) Malaria in the Netherlands Indies and
its control *J Malay Br Brit Med*
Ass I 4 pp 281-304 *RAEB*
XXVI, p 161
(Control of fish ponds intermittent
drying shading)
- 679 OVTCHINIKOV, K M et al (1937) Effect of different rates of flow in canals
and of small waterfalls on larvae
of *A. maculipennis* *Med Paras*
VI 1 pp 37-54 *TDB*, XXXIV
p 639
- 680 PAVLOVA P I (1932) Artificial alteration of the active reaction
of water reservoirs for the control of
mosquito larvae *Ibid*, I, 5-6 pp 265-
266 *RAEB* XXII p 21
- 681 PODGURSKII, P F (1936) Growing of trees along the margins of
breeding places as an anti mosquito
measure *Ibid* V, 1, pp 47-51
TDB, XXIV, p 227

- 682 PROSOLODOV, K. G. (1931) Observations on rice-fields *Trop Med Vet*, IX, 7-8, pp 357-358 *RAEB*, XX, p 116
(Intermittent drying)
- 683 RAM RAJA (1935) Control of anopheline larvae by flushing, *Ind Engng*, XCVIII, 4, pp 122-127
PAEB, XXIV, p 48
- 684 RAMSAY, G. C., and ANDERSON, I. P. (1940) Use of automatic siphon sluices on a group of tea estates in Northern Bengal *J Mal Inst Ind*, III, 1, pp 93-93
- 685 RAMSAY, G. C., and MACDONALD, G. (1936) The species control of anophelines in India *Ind Med Gaz*, LXXI, 12, pp 699-710 *RAEB*, XXV, p 95
(Shading)
- 686 RICE, F. M. (1935) Observations on malaria in Assam with special reference to cold weather and pre monsoon anti larval control *Rec Mal Sure Ind*, V, 4, pp 371-388
- 687 ROBERTSON, R. C., and CHANG, T. L. (1937) Rice cultivation in relation to malaria *China J*, XXVI, 6, pp 347-360
PAEB, XXVI, p 124
(Effects of manuring)
- 688 RUDOLFS, W. (1930) What are the prospects of eliminating food of mosquito larvae from breeding pools? *Procs 17th Ann Mtg N J Mosq Ext Ass*, pp 113-123 *RAEB*, XIX, p 129
- 689 RUSSELL, P. F. (1942) Control of *A. minimus* larvae in the Philippines by stranding and flushing *Phil J Sci*, XLVII, 4, pp 439-445
RAEB, XX, p 122
- 690 RUSSELL, P. F., and JACOB, V. P. (1939) Naturalistic control of Anopheles breeding in Casuarina pits *J Mal Inst Ind*, II, 3, pp 293-313
(Pollution of water agitation of water, shading changing of flora)
- 691 RUSSELL, P. F., and PAO, H. R. (1940) The Anopheles of rice-fields in South eastern Madras *Ibid*, III, 4, pp 427-446
(Intermittent drying)
- 692 SCHAEFF, J. W. (1938) Some methods of malaria control *J Malay Br Brit Med Ass*, II, 3, pp 165-168 *RAEB*, XXVII, p 126
(Flooding, silting, sluicing, fluctuating water levels, agitation of water surface)
- 693 SENIOR WHITE, R. A. (1936) Williamson's herbage cover method of larval control *Rec Mal Sure Ind*, VI, 3, pp 439-445

- 694 SINHA, J D (1940) . The theory and practice of rural malaria control in Bengal *Beng Pub Hlth J*, II, 1, pp 1-12
(Results of flood flush schemes)
- 695 SMALT, F H (1937) The periodic draining of rice fields to control malaria *Med Volk Ned Ind*, XXVI, 4, pp 285-299 *RAEB*, XXVI, p 105
- 696 STRICKLAND, C (1938) Holland and Bengal Reclamation and malaria in the two countries *Trans R Soc Trop Med Hyg*, XXXII 2, pp 277-286 *RAEB*, XXVII, p 28
- 697 STRICKLAND, C (1938) Malaria in relation to the coastal lowlands of Bengal *Ind Med Gaz*, LXXIII 7, pp 399-402 *RAEB*, XXVII, p 9
- 698 STRICKLAND, C (1939) . A study in physiography in relation to malaria in the Orissa coastal tracts *Riv Mal*, XVIII, 1, pp 38-41 *RAEB*, XXVII, p 248
- 699 STRICKLAND, C, and GIBSON, D (1934) *Backdoor drainage*, an anti malarial measure designed to meet a particular physiologic situation in Sylhet District, Assam *Ind Med Gaz*, LXIX, 8 pp 432-437 *RAEB*, XXII, p 219
- 700 STRICKLAND, C, and MURPHY R A (1932) River sand sifting and other anti malarial measures in South Sylhet, Assam *Rec Mal Surv Ind*, III 1, pp 125-142
- 701 STROMQUIST W G (1935) Malaria control in the Tennessee Valley *Civ Engng*, V, 12, pp 771-774 *RAEB*, XXIV, p 130
(Fluctuating water level)
- 702 SWEET, W C (1938) Irrigation and malaria *Procs Nat Inst Sci Ind*, IV, 2, pp 185-189 *RAEB*, XXVII, p 7
(Advocates regulation of water supply)
- 703 THOMSON R C M (1940) Studies on the behaviour of *A minimus* Paris I-III. *J Mal Inst Ind*, III 2 3, pp 265-348
(Shading, water movement, etc)
- 704 UBBINO, C M (1938) Malaria control and agricultural settlements *Mon Bull Bur Hlth Philipp*, XVIII, 7, pp 301-323 *RAEB*, XXVII, p 164
(Clearing vegetation, muddying, etc)
- 705 WALCH, E W (1933) Malaria control at Banjoewangi *Med Volk Ned Ind*, XXII, 1, pp. 16-19 *RAEB*, XXI, p 191
(Control of fish ponds)

- 706 WALCH, E. W., *et al* (1930) Sanitation of the salt water fish ponds of Batavia *Med Volk Ned Ind.*, XIX, 3, pp 400-430 *RAEB*, XIX, p 170
- 707 WASHBURN, R. E. (1913) An epidemic of malaria at Falmouth, Jamaica *Amer J Hyg*, XVII, 3, pp 656-665 *RAEB*, XXI, p 181 (Changing salinity of water)
- 708 WATSON, M. (1930) Anti malarial work *Bull Rubber Growers Assoc*, XII, 11, pp 603-606 *TDB*, XXVIII, p 609 (Pollution of water)
- 709 WATSON, L. B. and BISHOP E. I. (1940) Control of *A. quadrimaculatus* in the Tennessee Valley *Procs 27th Ann Mtg N J Mosq Ext Ass*, pp 145-153 (Fluctuating water level)
- 709a WIGGLESWORTH, V. B. (1933) Adaptation of mosquito larvae to salt water *J Exp Biol*, X, I, pp 1-37 *RAEB*, XXI, p 74
- 710 WILLIAMSON, K. B. (1933) *Ann Rpt Mal Advisory Board F M S for 1932*, p 9 (Iron pyrites, herbage packing, paddled clay)
- 711 WILLIAMSON, K. B. (1933-34) The control of rural malaria *Malay Agr Hort Sci J*, III, 3 t, pp 145-150, 201-206, IV 1 2 pp 224-228, 281-291 *RAEB*, XXIII, p 130
- 712 WILLIAMSON, K. B. (1933-36) The control of rural malaria by natural methods and references and notes supplementary to above L of N Pbn., 119 pp Singapore *RAEB*, XXV, p 263
- 713 WILLIAMSON, K. B. (1935) Need for action in relation to the biochemical investigation of anopheline breeding places *Trans 9th Cong F E A T M*, II, pp 83-87 *RAEB*, XXVI, p 126
- 714 WILLIAMSON, K. B. (1936) The soil in relation to malaria L of N Pbn., 7 pp. Singapore *RAEB*, XXV, p 69
- 715 WILLIAMSON, K. B., and SCHARFF, I. W. (1936) Anti larval flushing *Malay Med J*, XI, 3 pp 123-150 *RAEB*, XXV, p 30
- 716 WORTH, H. N. (1937) The control of anopheline breeding in river beds *Trans Foy Soc Trop Med Hyg*, XXV, 5, pp 521-530 *RAEB*, XXV, p 168 (Training of streams)
- 717 WORTH, H. N. and SUBRAMANYAN, K. (1940) Anti larval flushing of rivers and streams in Ceylon *J Mal Inst Ind*, III, 1, pp 81-92

- 718 YAKUSHEV M R (1937) Anti larval methods in mill ponds *Med Paras* VI 2 pp 31-39 *RAEB* XVI p 46
(Intermittent drying)
- 719 ZON B B (1939) Salt water fish ponds and malaria *Geneesk Tijd Ned Ind* LXXIX 9 pp 529-540 *RAEB* XXVII p 159

Deleterious Plants and Trees

- 720 BARRER M A and FORBES I R (1933) Malaria in the irrigated regions of New Mexico *U S A Pub Hlth Rpts* XLVIII 22 pp 610-613 *RAEB* XXI p 196
(Lucerne clover *Chara*)
- 721 BEKLEMISHEV V N and POLOVODOVA V (1933) Influence of plants on the larvae of *A. maculipennis* *Med Paras* II 6 pp 341-363 *RAEB* XVII p 5
(*Eleocharis* etc)
- 722 BRADLEY G H (1931) Feeding tests of anopheline mosquitoes with leguminous plants *J Econ Ent* XXIV 6 pp 1229-1233 *RAEB* XX p 54
(Clover)
- 723 DUTT N B (1931) Influence of leguminous crops on malaria *Ind Med Rec* LI 11 pp 343-6
- 724 HAMLYN HARRIS P (1932) Some further observations on *Chara fragilis* in relation to mosquito breeding in Queensland *Ann Trop Med Paras* XXVI 4 pp 519-524 *RAEB* XVI p 52
- 725 DUFFRELL F (1932) Clover and malaria *Amer J Hyg* XVI 2 pp 609-617 *RAEB* XX p 250
- 726 MATHESON R (1930) Aquatic plants as factors in mosquito control *Procs 17th Ann Mtg N J Mosq Ext Ass* pp 81-104
- 727 MATHESON P (1930) Utilization of aquatic plants as aids in mosquito control *Smithson Rept* 1931 pp 413-430 *RAEB* XVI p 176
- 728 MATHESON, R and HINMAN E H (1931) Further work on *Chara* spp and other biological notes on Culicidae *Amer J Hyg* XIV 1 pp 99-108 *RAEB* XIX p 209
(*Chara* *Hydra*)
- 729 SOESILO R (1930) Further note on malaria and plants containing coumarin *Geneesk Tijd Ned Ind* LXXII 6 pp 329-380 *RAEB* XX p 121
(Clover etc)

- STRATMAN THOMAS, W K (1931) Supposed antagonism between alfa alfa and malaria *Amer J Hyg.* XIV, 2, pp 394-410 *RAFB*, XIX, p 249
- TWINN, C R (1931) See No 904
- ZIEMANN, H (1931) Possible relation between plants containing coumarin to the epidemiology of malaria *Arch Schiffs Trop Hyg.* XXXV, 7, pp 410-420 *RAEB*, XIX, p 193 (Clover, etc.)

Drainage

- ADAMS, C M (1936) Mosquito control engineering VII Ending malaria in New Mexico *Engng News Rec*, Sept 10 pp 372-374
- BARROWMAN B (1932) An example of anti malarial subsoil drainage under rural conditions *Malay Med J*, VII 4 pp 116-118 *TDB*, XXX p 493
- BARROWMAN B (1939) An expedient in anti malarial subsoil drainage *J Malay Br Brit Med Ass* III 2 pp 175-176 *RAEB*, XXVIII, p 65
- BLACKLOCK, D B (1936) Simple forms of subsurface drainage for anti mosquito work *Ann Trop Med Paras*, XX 4 pp 501-510 *RAEB*, XXV p 103
- CLARKE, J L (1936) Mosquito control engineering, IV Inland control methods *Engng News Rec*, Aug 20 pp 266-269
- CLARKE, A G (1931) Proper drainage as a means of mosquito abatement *Municipal Sanitation*, II, 5 pp 223-226 Also *Abst VI, Pub Hlth Engng RAFB* XIX, p 261
- COLLIGNON, F (1934) Inadequacy of flat bottomed drains to prevent mosquito breeding *Arch Inst Past Alger*, VII, 1 pp 26-29 *TDB*, XXVI p 712
- CORKEAN, W S (1936) Mosquito control engineering VII Mosquito abatement in Delaware *Engng News Rec* Sept 10, pp 374-376
- CORNER, J D (1932) The employment of the portable pump in anti malaria schemes *J R I W C*, LIX, 2 pp 134-141 *PAEL*, XX, p 215
- CORY F N., and CROSTH-WAIT S L (1937) A large scale demonstration in mosquito control (Maryland) *J Econ Ent.* XXV, 5 pp 711-712 *PAEB*, XXVI p 54

- 742 CRAIG, W J F (1937) Anti malaria drainage work in the new Chanai Cantonment (Malaya) *J R A M C*, LXVIII, 1-2 pp 15-6 73 85 *RAFB*, XXV p 194
- 743 DEBWERKER, R J van (1936) Mosquito control engineering V The salt marsh problem *Engng News Rec* Aug 27, pp 304-307
- 744 ELMENDORF J E et al (1939) Concrete invert and tile manufacture *Procs 21st Mtg Nat Mal Comm* 1938 pp 46-59 (Typescript)
- 745 GRAY H F (1936) Mosquito control engineering VIII California's campaign *Engng News Rec* Sept 17, pp 405-406
- 746 CRIFFIN F C (1938) Surface and subsoil drainage *Procs Nat Inst Sc Ind*, IV, 2 pp 219-222 *RAEB* XXVII p 8
- 747 HARKNESS J W P, and SAMANA, M Y (1937) The use of dynamite in anti malarial drainage *J Roy San Inst*, LVIII 5 pp 337-345 *TDB* XXXV, p 96
- 748 HERMS W B (1934) Mosquito control in California under the CWA *J Econ Ent* XXVII 5 pp 1014-1029 *RAEB* XXIII p 23
- 749 LAL R B and SHAH N S (1933) Malaria and water logging *Rec Mal Surv Ind*, III 4 pp 583-634 (Effect of reclamation by drainage for agricultural purposes)
- 750 MAIL, G A (1933) Mosquito control in Montana *Montana Agric Exp Sta Circ*, No 143 13 pp *RAEB* XXII p 181
- 751 ONEJLL J H (1938) Concrete lined ditches for mosquito control *American City* LIII 8 p 38 *Summ Pub Hlth Engng Abst*, XIX, 1939 *TDB*, XXXVII p 364
- 752 PRINCE J A LE (1936) Mosquito control engineering. IX Malaria and the Mississippi Valley *Engng News Rec*, Sept 17, pp 404-405
- 753 RANDLE, T A (1940) New developments in underground drainage for malaria control *Amer J Pub Hlth*, XXX, 7 pp 736-742
- 754 RECTOR N H (1938) Different phases of permanent drainage for malaria control in Mississippi *South Med J* XXXI 7 pp 814-815 *RAFB*, XXVII, p 30
- 755 SAMUELS W F (1931) Anti malarial work at Tanjong Rambutan Malaya *J Mental Sci*, LXXVII 318 pp 555-561 *TDB*, XXIX, p 336 (Subsoiling with milk tins)

- 756 SCHAEFF, J W (1933) Method of planting and growing grass in connection with anti malarial drainage *Malay Med J*, X, 4, p 153 *RAEB*, XXIV, p 248
- 757 SCHARFF J W (1935)
 † Anti malarial drainage from the point of view of the Health Officer 104 pp Penang *RAEB*, XXIII, p 236 Also in *Malay Med J*, X, 4 pp 119-137 XI, 1-2, pp 40-52, 67-70 *RAEB*, XXIV p 248
- 758 SCOTT D F (1936) Malarial control by vertical drainage *Proc Ann Mtg Assn Br Brit Med Ass* pp 22-23 *TDB*, XXXIV, p 638
- 759 SCOTT G W (1930) Report on malaria prevention work carried out at Chenderoh *Malay Med J*, V 3, pp 86-89 *RAEB*, XIX p 27
- 760 SCOTT G W (1934) Fascine drainage in mosquito control *Ibid* IX 1, pp 65-66 *RAEB* XXII, p 149
- 761 SEXTON WHITE R A (1936) Malaria control by subsoil drainage at Waltair *Rec Mal Surg Ind*, VI 1 pp 13-18
- 762 STRUTHERS F A and NINHA, S N (1934) The use of bamboo in subsoil drainage *Malay Med J* IX 4, pp 197-199 *RAEB*, XXIII, p 92
- 763 TWEEDIE D R (1940) Fascine drainage as an anti-malaria measure *J Malay Pr Prot Med Ass*, IV 2 pp 167-171
- 764 VERTEUIL E J DE and SPENCE T (1937) Malaria in Trinidad Low tide level culvert system in coastal drainage *Trans Roy Soc Trop Med Hyg*, XXX 4 pp 449-460 *RAEB*, XXV p 106
- 765 WOOD T F M (1941) Two simple anti mosquito measures *J I A M C* LIX 6, pp 450-452 *RAEB* XXI p 47
Treatment of borrowpits)

Fish

- 766 ARNAUD J (1943) Malaria control at In Salah (Tidikelt) Trial of *Gambusia* in a Sahara oasis *Arch Inst Past Algér*, XIII, 3 pp 369-376 *RAEB* XXIV, p 11
- 767 BELENNIKHEV A N and LAZUSKI G F (1935) Use of *Gambusia* *Med Paras*, IV, 4, pp 327-328 *RAEB*, XXIV, p 32
- 768 BORRERO C P (1933) On the identity of the species of topminnow introduced into Italy for malaria control *Zool Anz*, CV, 1-2 pp 9-14 *RAEB*, XXII p 20

- 768 BOGDANOVICH, I N. (1935) Acclimatization of *Gambusia* in Turkmenistan *Med Paras*, IV, 5, pp 413-416 *RAEB*, XXIV, p 73
- 769 BOGOJAWLENSKI, N A (1936) Failure of *Gambusia* to destroy *Anopheles* larvae in nature (Russian Azerbaijan) *Arch Schiffs Trop Hyg*, XL 5, pp. 201-203 *RAEB*, XXIV, p 140
- 770 COULON, G, and SAUTET, J (1931) *Gambusia* and malaria in Corsica *Ann Paras Hum Comp*, IX, 6, pp 530-545 *RAEB*, XX, p 87
- 771 DANILOVA, M I, and IAPPIV, G I (1936) Transport of *Gambusia* *Med Paras*, V, 4, pp 579-583 *RAEB* XXV, p 141
- 772 ENIKOLOPOV, S K (1935) Biology of *Gambusia* acclimatized in Dagestan *Ibid*, IV, 5, pp 408-413 *RAEB*, XXIV, p 73
- 773 GUPPY, P L (1934) Observations on Trinidad larvicidal fishes *Trop Agric*, XI, 5, pp 117-122 *RAEB*, XXII, p 128
- 774 HOLLAND, E A (1933) Control of malaria in New Ireland by *Gambusia* *Trans Roy Soc Trop Med Hyg*, XXVI, 6, pp 529-538 *RAEB*, XXI, p 156
- 775 HORA, S L, and NAIR K K (1938) Observations on the nutrition of *Panchax panchax* *Procs Nat Inst Sci Ind*, IV, 2, pp 245-251 *RAEB*, XXVII, p 8
- 776 JOHN, C C (1940) Utility of *Aplocheilichthys lineatus* for mosquito control *J. Nat Inst Ind*, III, 1 pp 67-80
- 777 KAIANDADZE, J, and MCHELIDZE, I (1934) History of introduction of *Gambusia* into Europe and its establishment in Russia *Med Paras*, III, 4, pp 336-339 *RAEB*, XXIII, p 76
- 778 KULAGIN, S M and MARTZINOVSKIY V I (1936) Attempt to acclimatize *Gambusia* in the cooling tanks of a power station *Ibid*, V, 1, pp 62-61 *RAEB*, XXIV, p 227
- 779 LEMASSON, J (1937) Use of *Gambusia affinis* and *Gambusia guppyi* in malaria control *Bull Econ Indochine*, XL, 2 pp 328-330 *RAEB*, XXVI, p 21
- 780 LINDBERG, G U (1934) *Gambusia* *Mag Paras*, IV, pp 351-367 *RAEB*, XXIV, p 54
- 781 MORIN, H G S, et al (1936) Use of fish against mosquitoes *Arch Inst Past Indochine*, VI, 24, pp 413-505 *RAEB*, XXVI, p 20
- 782 MULLIGAN, H W, and MAJID, S A (1936) Care, transportation and use of *Gambusia* under Indian conditions *Rec. Mal Surv Ind*, VI, 4, pp 537-547

- 783 ORYER (1931) *Gambusia* on the Cote d'Ivoire Abst in *Bull Mens Off Int Hyg Pub*, XXIII, 6, pp 1099-1100 *RAEB*, XIX, p 168
- 784 PANAOLA, A (1932) Experiments and observations on work against anopheline larvae *Rev Med*, XI, 4, pp 453-467 *RAEB*, XX, p 264
- 785 PETRISHCHEVA, P A (1936) Role of *Gambusia* and other fish in the control of malaria in Turkmenistan. In *PAYLOVSKII, E., Animaux Pathog.*, pp 111-138 *RAEB*, XXVI p 9
- 786 PRASAD, B, and HORA S L (1936) A general review of the larvivorous fishes of India *Rec Med Surv Ind*, VI, 4, pp 631-648
- 787 PRENDEL A P et al (1933) Acclimatization of *Gambusia* in the U S & R *Med Paras*, I, 5-6 pp 261-264. *RAEB*, XXII, p 20
- 788 REBRIN, M (1938) Acclimatization of a new larvicidal fish in Uzbekistan *Soviet Sci Tech*, VI 7, pp 106-107 *RAEB*, XXVII, p 76 (*Lebistes reticulatus*)
- 789 RIMBAUT G, and MATHIS M (1935) Use of 'millions' against larvae at Dakar *Bull Soc Path Exot*, XXVIII, 7, pp 575-577 *RAEB*, XXIII, p 255 (*Lebistes reticulatus*)
- 790 FOULE, S (1934) Role of larvivorous fish in prevention of malaria 61 pp Paris *RAEB*, XVII, p 258
- 791 ROY D N (1938) Control of malaria mosquitoes in Bengal by fish *J Mal Inst Ind*, I, 4, pp 405-416 (*Panchax panchax* *Barbus phatunio*)
- 792 RUIBINSKII S V, and LEVIT, M S (1934) Fish in malaria control in the Ukraine *Pre Microbiol*, XIII, 2, pp 151-159 *RAEB*, XXIII p 86
- 793 RUIBINSKII S V and OVCHINNIKOV, M M (1935) Rearing *Gambusia* in ponds near Kiev *Rev Med*, XIV 4 pp 369-378 *RAEB*, XXIV p 11
- 794 ROCKADZE N P (1934) *Gambusia* in Abkhazia *Med Paras*, III, 1 pp 60-69 *RAEB*, XXII p 193
- 795 PERRYELL, P F and JACOB V P (1939) Use of fish to control *Anopheles* breeding in Casuarina pits *J Mal Inst Ind*, II 3 pp 273-291 (*Gambusia affinis* *Panchax* *Mochis*, *Aplocheilichthys melanocephala*)
- 796 SFRONCT, ET (1932) Attempt to stock an Algerian river with *Gambusia* *Arch Inst Past Alg*, X, 3 pp 348-355 *RAEB*, XXI, p 20

- 797 SERGENT, ET (1939) Cannibalism of *Gambusia*, and a method of combating it *Arch Inst Past Alger*, XVII, 1, pp 139-142 *RAEB*, XXVII, p 233
- 798 SERGENT, ET, *et al* (1932) Control of Anopheles with *Gambusia* in an Algerian reservoir *Ibid*, X, 2, pp 157-156 *RAEB*, XXI, p 20
- 799 SICAULT, G (1934) . Adaptation of *Gambusia* to saline waters *Bull Soc Path Exot*, XXVII 5 pp 485-488 *RAEB*, XVII p 158
- 800 SICAULT, G and ROULE, S (1935) Biology of *Gambusia* in Morocco *Ibid* XXVIII, 2, pp 134-141 *RAEB* XXIII, p 147
- 801 SOKOLOV N P (1936) Acclimatization of *Gambusia* in Central Asia *Riv Mal*, XV, 5, pp 325-344 *RAEB*, XXV, p 32
- 802 TOUMANOFF C (1933) On a first attempt to acclimatize *Girardinus guppyi* at Tonkin *Bull Soc Path Exot*, XXVI, 4, pp 632-638 *RAEB*, XVI, p 168
- 803 TRAUSMILLER, O (1932) Limits of malaria control by *Gambusia* *Arch Schiffe Trop Hyg*, XXXVI 10 pp 530-539 *RAEB*, XV, p 265
- 804 WATSON, M (1936) A larvivorous fish in the Belgian Congo *Ann Soc Belge Med Trop*, XVI, 1 pp 153-158 *RAEB*, XXIV, p 187 (*Tilapia heudeloti*)
- 805 ZAVATTARI, E (1934) The acclimatization of *Gambusia* in Ferran *Riv Mal*, XIII, 5, pp 617-623 *RAEB*, XXIII, p 59

Fumigants

- 806 GRIFFITHS, T H D, *et al* (1935) Destruction of mosquitoes in aircraft *Bull Off Int Hyg Pub* XXXII 3 pp 550-560 *RAEB*, XXIII p 154 (HCN carbonyl)
- 807 GUTZEVICH, A V, and PODOLYAN, V. Y (1935) Pyrethrum smoke candles to control mosquitoes and sandflies *Rev Microbiol*, XIV, 1, pp 87-98 *RAEB* XXIII, p 291
- 808 JAMES, J F (1935) Fumigation and trapping of mosquitoes *Jl R A M C*, LXV, 4, pp 267-269 *RAEB*, XXIII, p 281 (Burning pyrethrum coils)
- 809 SELIVANOV, K P (1937) .. Use of traps in the fumigation of mosquitoes *Med Paras*, VI, 2, pp 243-249 *RAEB*, XXVI p 48

- 810 SYMES C B (1935) Insects in aeroplanes *Rec Med Res Lab, Kenya* No 6 16 pp *RAEB* XXIV p 65
(HCN gas)
- 811 THOMPSON, T O (1936) Mosquito destruction by James method *J R A M C*, LXVII 5, pp 331-335
RAEB, XXV, p 73
- 812 TRUILLARD, M (1934) The destruction of adult Anopheles as an anti-malaria measure in S Indo-China *Bull Soc Path Exot*, XXVII 10 pp 937-939 *RAEB*, XXIII, p 81
- 813 WILLIAMS C L and DREXSEN, W C (1935) The destruction of mosquitoes in airplanes *U S A Pub Hlth Rpts* L, 20, pp 663-671 Also in *Bull Off Int Hyg Pub*, XXVII, 7 pp 1350-1359 *RAEB* XXIII p 206
(Carbocide)

Miscellaneous Measures against Adult Mosquitoes

- 814 RADLETZ N A, and KIZMINA, L A (1935) The dispensary method of controlling hibernating anophelines *Parasites transmetteurs anm acmeux* pp 264-268 *RAEB* XXIV p 180
(Destruction of hibernating adult mosquitoes)
- 815 KREMER B I and KUVICHINSKII B S (1936) A new method for destroying adult mosquitoes *Med Paras* V, 6 pp 955-958 *RAEB* XXV p 200
(Imagocade-anabasine sulphate mixed with slaked lime)
- 816 KREMER B I and KUVICHINSKII B S (1937) Anabadust for killing adult mosquitoes *Med Paras* VI 4, pp 550-555 *RAEB* XXVI p 170
(Anabasine sulphate mixed with slaked lime)
- 817 PIVOVAROV V M, and GUTERMAN I M (1937) Use of anabadust against hibernating mosquitoes. *Ibid*, 3 pp 424-432 *IHP* XXVI p 169
- 818 SMOLNIKOV, V M and KONDRATYEV, P V (1937) The Sh apparatus *Ibid* 3 pp 433-435 *RAEB*, XXVI p 169
(For insecticidal dusts)
- 819 THOMSON I (1932) A possible means of cutting down the mosquito population *Science*, LXXX 1133 pp 78-79 *RAEB*, XX, p 209
(Sound apparatus to attract males)

Miscellaneous Measures against Larvæ of Mosquitoes.

- 320 AFANASEY, S I (1938) Use of residue from conversion of petroleum to coke as a larvicide *Med. Paras*, VII, 5, pp 691-700 *RAEB*, XXVII, p 173
- 321 ALFSSANDRINI, G (1934) Malaria in the pontine marshes in 1933, *Conquista D Terra* 1934, No 2, reprint, pp 17 *RIFB*, XXII, p 133 (Carburel)
- 322 ANON (1934) Destruction of mosquitoes by leaves of the prickly pear (*Opuntia vulgaris*) *Rev Hort Agric Afr Nord*, XXVIII, 9, p 250 *RAEB*, XXII, p 233
- 323 BALI, B (1938) Action of extracts of tobacco on mosquito larvæ *Riv Mal*, XVII, 1, pp 122-125 *RAEB*, XXVI, p 231
- 324 BANA, F D (1936) A practical way of dealing with *Aedes ægypti* breeding in country craft *Ind Med Gaz*, LXXI, 2, pp 79-80 *RAEB*, XXIV, p 115 (Systematic emptying of water containers)
- 325 BANA, F D (1938) Control of *Aedes ægypti* mosquitoes in Indian country craft by a mosquito proof metal cap (Bennett pattern) for drinking water receptacles *Ibid* LXXIII, 6, pp 344-346 *RAEB*, XXVI, p 207
- 326 BAROODY, B J (1932) Effect of various chemicals on larvæ and pupæ at various temperatures *J. Elish Mitchell Sci Soc*, XLVIII, 1, pp 123-129 *RAEB*, XXI, p 10 (Iodine, formaldehyde, sodium hydroxide; mixture of copper sulphate, mercury bichloride, iodine and formaldehyde)
- 327 BOTSFORD R C, and TURNER, N (1933) Mosquito control in Connecticut, 1932 *Bull Conn Agric Exp Sta*, No 349, pp 439-444 *RAEP*, XVI, p 210 (Rotenone fuel oil, fuel oil, naphthalene fuel oil; pyrethrum powder)
- 328 BRINK, C J H, and CHOWDHURY, D K D. (1939) Ammonium sulphate as a combined fertilizer and mosquito larvicide (Abstract). *J Mal Inst Ind*, II, 1, pp 111-112
- 329 BUCHMANN W (1934) Size of particles of the active material in aqueous suspensions of pyrethrum insecticides *Z Angew Ent*, XX, 1, pp 136-149 *RAEB*, XVI, p 171

- 830 BUTT, N M (1938) An inexpensive apparatus for mixing (liquid) larvicides *J Mal Inst Ind.* I 2 pp 147-148
- 831 CAMPBELL T L, *et al* (1934) Insecticidal tests of synthetic organic compounds, chiefly sulphur compounds, against culicine mosquito larvae *J Econ Ent* XXVII 6 pp 1176-1185 *RAEB* XXIII p 118
- 832 CAWSTON F G (1936) *Opuntia* as a larvicide *J Trop Med Hyg.* XXIX 12 pp 137-138 *RAEB* XXIV p 217
- 833 CHOPRA R N ROY D N and GHOSH S M (1940) Insecticidal and larvicidal action of *Tephrosia vogelii* *J Mal Inst Ind.* III I pp 183-189
- 834 CHOPRA R N ROY D N and GHOSH S M (1940) Action of pyrethrum on mosquito larvae *Ibid* 4 pp 457-463
- 835 CHOPRA R N ROY D N, and GHOSH S M (1940) *Blumea densiflora* and *Aztecium vulgare* Their insecticidal and larvicidal properties *Ibid* pp 494-498
- 836 DANILOVA M I and BUDIMKO F A (1939) Acidol as a larvicide *Med Paras.* VIII 3 pp 177-178 *RAEB* XXVIII p 8
(Mixture of naphthalene acids obtained from distillation of petroleum oils)
- 837 DUREN A N (1938) Creolin as a larvicide *Ann Soc Belge Med Trop* VIII 1 pp 23-25 *TDB.* XXVII p 581
- 838 FROBISHER M and SHANNON R C (1931) Effects of certain poisons on mosquito larvae *Amer J Hyg* XIII 2 pp 614-633 *RAEB* XIX p 140
(Iodine bromine potassium cyanide potassium hydroxide HCN gas calcium hypochlorite)
- 839 GIBSON A (1931) Mosquito pyrethrum in Canada in 1930 *Proc 18th Ann Mtg V J Mosq Ext Ass* pp 23-39 *RAEB* XX p 52
(Dusts made by mixing pyrethrum extracts with kaolin powdered gypsum or slaked lime)
- 840 GINSBURG J M (1930) Studies of pyrethrum as a mosquito larvicide *Ibid* 17th Mtg pp 57-62 *RAEB* XIX p 128
- 841 GINSBURG J M (1934) Soap as a larvicide *Science* LXVIII, 2014 p 210 *RAEB* XXII p 87
- 842 GINSBURG J M (1931) Mosquito larvicides *Proc 21st Ann Mtg V J Mosq Ext Ass* pp 191-197 *RAEB* XXIII p 151
(Pyrethrum kerosene-emulsion)

- 842 GINSBURG J M (1935) Larvicides and a method for temporary protection from adult mosquitoes in limited areas *Procs 22nd Ann Mtg N J Mosq Ext Ass* pp 147-151 *RAEB XXIII*, p 205
(Improved pyrethrum emulsion)
- 843 GINSBURG J M (1937) Studies in reducing volume of oil necessary to kill mosquito larvae by incorporating pyrethrum *J Econ Ent*, **XXX** 9 pp 328-332 *RAEB XXV* p 201
- 844 HANLYN HARRIS P (1931) Mosquitoes breeding in tree cavities in Queensland *Bull Ent Res* **XXII** 1 pp 51-59 *RAEB XIX* p 119
(Paradichlorobenzene)
- 845 HARGREAVES H (1936) Ann Rpt Govt Entomologist *Rept Med Dept Uganda* 1935 pp 69-73 *RAEB XXIV* p 273
(Cotton seed tar *Opuntia*)
- 846 HENDERSON, J M and HOWARD R S (1939) A comparative evaluation of paris green and pyrethrum emulsion larvicide for malaria control in Georgia *Procs 21st Mtg Nat Val Comm* 1938 pp 97-111 (Typescript)
- 847 HOSKINS W M (1934) Toxicity of acid and basic solutions of sodium arsenite to mosquito pupae *J Econ Ent* **XXV**, 6 pp 1219-1224 *RAEB XVI* p 34
- 848 ILLINGWORTH J F (1931) Use of arsenic in mosquito control *Proc Hawaii Ent Soc*, **VII** 3 p 414 *RAEB XIX* p 169
(Paris green lead arsenate)
- 849 JORDAN J (1937) Mosquito larvicidal measures *Chinese Med J* **LI** 6 pp 927-936 *TDH XXIV*, p 944
(Pyrethrum emulsion larvicide)
- 850 KALANDADZ L P and LEMER M (1936) Chloropicrin as a larvicide *Med Paras V* 4 pp 568-578 *RAEB XXV* p 144
- 851 KEMPER H (1933) Effect of pyrethrum powder on different animals especially aquatic species *Z Gesundheits U Stadtehyg* **XXV** 3 pp 149-164 *RAEB XVI* p 170
- 852 KING W V (1938) Experiments with phenothiazine and mosquito larvae *J Econ Ent* **XXXI** 5 pp 610-611 *RAEB XXVII* p 53
- 853 KING W V and McNEEL T E (1938) Experiments with paris green (as a spray) and calcium arsenite as larvicides for culicine mosquitoes *Ibid* 1 pp 85-86 *RAEB XXVI* p 144

- 854 KUCHER S S (1934) Anthracene a new mosquito larvicide
Med J aas III 2 pp 141-143
RAEB XVII p 19
 (By product of the coke benzene industry)
- 855 LEVENSON F D (1931) Observations on the effect of various chemical substances on the development of the larvae of *Anopheles maculipennis*
Trop Med Vet IX 10 pp 497-501
RAEB XX p 119
 (Various acids and alkalis used in pest b g ts)
- 856 MCDWELL F I (1939) Larvicides and contact sprays used in mosquito control
Quart Bull Mal Agric Exp Sta XVII 1 pp 3-34
RAEB XXVIII p 43
 (Popular account)
- 857 MA G A (1936) Thiodiphenolamine (phenothiazine) a new repellent of mosquito larvae
J Econ Ent XXIX 6 pp 1144-1146
RAEB XXV p 83
- 858 MANSON D (1939) The action of certain Assamese plants as larvicides
J Mal Inst Ind II 1 pp 85-93
 (*Duranta Polygonum Garden Zanthoxylum Tephrosia*)
- 859 MAY I M (1938) Mosquito control by hexachlorethane
Rev Pathog XVIII 1 pp 5-19
FAFB XXIV p 18
 (Applied as dust with talc as diluent)
- 860 MUELLER, P P and GUYER RAY F M (1937) Control of mosquito breeding in fish ponds
Med Paras VI 2 pp 239-24
RAEB XXVI p 47
 (Arsenical dusts)
- 861 NABOKOV V A et al (1933) Arsenical (copper arsenite) as a larvicide
Ibid II 12 pp 59-63
RAFB XII p 37
- 862 O'HANE W C WELTGATE W A and GLOVER, L C (1934) Studies of contact insecticides
Tech Bull N H Agr Exp Sta No 58 3 pp
RAEB XVIII p 55
 (Nicotinic heptyl acetate caproacetate)
- 863 PERCOTT and ESCALER G (1934) Anti-malaria campaign round Rome in 1933
R Mal XIII 5 pp 673-688
RAEB XXIII p 59
 (Carbureol)
- 864 POLITOV V K (1932) Chloroform and carbon bisulphide in the control of mosquito larvae and pupae
Med Paras I 1 pp 32-3
RAFB XI p 257

- 865 ROMAN, E. and NETIEN, G (1939) Physiological action of *Derris elliptica* Benth on mosquito larvae *Bull Soc Linn Lyon*, VIII, 3, pp 74-79 *RAEB*, XXVII, p 232
- 866 ROSS, G A P (1932) Bird lime for killing mosquitoes *J Trop Med Hyg*, XXV, 16, p 256 *TDB* XXX, p 45
- 867 ROUBAUD, F (1931) On the advantages and methods of action of larvicidal dusts *Proc 2nd Int Cong Mal Algér*, II, pp 39-44 *RAFB*, XX, p 73
- 868 RUSSELL P F, and WEST, A F (1932) Larvicidal effects of arsenicals other than paris green *Phil J Sci*, XLIX 1 pp 97-103 *RAEB*, XX p 237 (Arsenic trioxide, zinc arsenite, etc)
- 868a RUSSELL P F, and WEST, A F (1932) Various toxic substances partially adsorbed on charcoal as larvicides *Ibid*, 2 pp 211-215
- 869 SATTA, E (1938) Anti larval potency of hexachlorethane *Rend Ist San Publ*, I, 3, pp 885-892 *TDB*, XXXVII p 136 (Unfavourable report)
- 870 SCHAFFER, P S, et al (1937) Toxicity of phenothiazine derivatives to culicine mosquito larvae *J Econ Ent*, XX, 2, pp 361-363 *RAEB*, XX p 214
- 871 SCHARFF, J W (1936) Filling tree holes, an anti mosquito measure *Malay Med J*, XI, 1, p 68 *TDB*, XXXIV, p 71
- 872 SCHWARDT, H H (1939) Biologies of Arkansas rice field mosquitoes *Bull Arkansas Agric Exp Sta*, No 377, 22 pp *RAEB*, XXVII, p 257 (Pyrethrum emulsion)
- 873 SHANNON, R C, and FROBISHER, M (1931) Effect of various substances on larvae of *Aedes aegypti* Lmer *J Hyg*, XIV, 2, pp 426-432 *RAFB*, XIV, p 250 (Creolin a proprietary substance of the crude carbolic type, probably containing phenol the cresols and related substances, phenol, alum borax)
- 874 SHEPARD, H H (1931) Relative toxicity of rotenone and nicotine to mosquito larvae *J Econ Ent*, XXIV, 3 pp 725-731 *RAFB*, XIV, p 198
- 875 SHIKHOV, N I, and SHCHERBAKOVA, O A (1937) Effect of paris green calcium arsenite and oleogumbrin on rice crops *Med Paraz*, VI, 5, pp 683-695 *RAEB*, XXVI p 171

- 86 SHUPITZINA N K (1938) Action of larvicide powders on larvae of Anopheles *Med Pa ss VII 3* pp 346-371 *RAEB XXVII* p 68 (Paris green arsenical acid)
- 87 SHVETZ S I and LAFIN G I (1936) Larvicide action of residual products of crude petroleum *Ib id V 9* pp 171-173 *RAEB XXIV* p 93
- 88 SMIT J W (193) One year's experience with pyrethrum larvae *Proc 19th Ann Mtg N J Mosq Ext 4 :* pp 128-131 *RAEB XX* p 241
- 89 SOESILO R and HOOT O H VAN DER (193) The connection between malaria and the disposal of manures in relation to channels *Centrnl Trop Med Ind LXVII 13* pp 84-89 *PAEB XX* p 193
- 90 STEWART A I and MOORTHY V N (1933) Lethal properties of aqueous extract of young bamboo shoots *Ind Med Gaz LXVIII 6* pp 320-323
- 91 STRONG L A (193) Rpt Bureau Entom and Plant Quarantine *U S A Dept Agric 98* pp Washington *IAEB XXVI* p 110 (Cal. marconite)
- 92 STRONG L A (1938) Rpt Bureau Entom and Plant Quarantine *Ibid 84* pp Washington *RAEB XXVII* p 156 (Pyrethrum based oil mixture & hew nut shell oil)
- 93 TILLI P (1933) Control of anopheline larvae by calcium cyanamide *Riv Mal XII 4* pp 2-730 *RAEB XXI* p 275 (Also experiments with superphosphates potassium sulphate sodium nitrate and ammonium sulphate)
- 94 TILLI P (1935) Control of anopheline larvae near Rome with calcium cyanamide *Ib id XIV 9* pp 19-90 *RAEB XXIII* p 94
- 95 TILLI P (1937) Infestation with calcium cyanide *Ib id XVI 1* pp 54-59 *RAEB XXV* p 114
- 96 TILLI P (1938) Modern agricultural methods and the use of calcium cyanamide *Ib id XVII 1* pp 55-930 *RAEB XXVI* p 75
- 97 TURNER N (1933) Traps for larvae *Proc 19th Ann Mtg N J Mosq Ext 4 :* pp 100-101 (Najolia no pyrethrum larvae)

- 888 ULYANDZHEV, V. I., and RIFLING, E. A. (1936) Olearsenite for control of mosquito larvae. *Summary Sci. Res. World Inst. Plant Prot.*, 1935, pp. 473-475. *RAEB*, XXV, p. 68.
(A 1 in 4 mixture of calcium arsenite in oleogumbrin, a waste product of petroleum.)
- 889 VANDI, V. (1933) A further advance in the control of mosquito larvae. *Riv. Mal.*, XII, 3, pp. 532-534. *RAEB*, XXI, p. 260.
(Dust from residue of distillation of bituminous schists.)
- 890 VANNOTE, R. L., and GINSBURG, J. M. (1931) Practical application of pyrethrum mosquito larvicide. *Procs. 18th Ann. Mtg. N. J. Mosq. Ext. Ass.*, pp. 111-120. *RAEB*, XV, p. 52.
- 891 VOLZHENSKII, L. E. (1935) Laboratory experiments with schists decomposition products. *Med. Paras.*, IV, 4, pp. 316-320. *RAEB*, XXIV, p. 31.
- 892 WASFY, O. (1936) A new larvicide. *J. Egypt. Med. Ass.*, XIX, 10, pp. 592-608. *RAEB*, XXIV, p. 294.
(Acid oil sludge, a waste product of oil refining.)
- 893 WASSILIEFF, A. (1931) A new arsenical larvicide. *Arch. Inst. Past. Tunis*, XXIII, 4, pp. 449-454. *RAEB*, XVIII, p. 187.
(CuHAsO_4 adsorbed on chalk.)
- 894 WATTS, R. C., and BHARUCHA, H. H. (1938) Larvicides for anti-mosquito work, with special reference to cashew nut shell oil. (Abstract.) *J. Mal. Inst. Ind.*, I, 2, pp. 217-219.
- 895 WEST, A. P., and RUSSELL, P. F. (1932) Experiments with various substances partially adsorbed on charcoal as larvicides. *Phil. J. Sci.*, XLIX, 2, pp. 211-217. *RAEB*, XXI, p. 9.
(Barium chloride, pyrethrum powder, nicotine, tannic acid, rotenone.)
- 896 WORSLEY, R. R. DE G. (1934) The insecticidal properties of some East African plants. I. *Ann. App. Biol.*, XXI, 4, pp. 649-669. *RAEB*, XXIII, p. 27.
(*Tephrosia vogelii*.)
- 897 YATZENKO, F. I. (1934) Chloropicrin for control of mosquito larvae. *Med. Paras.*, III, 1, pp. 91-93. *RAEB*, XXII, p. 19.
- 898 YURCHAN, I. F., and HENKO, V. P. (1939) Aqueous suspensions of larvicides. *Ibid.*, VIII, 2, pp. 170-176. *RAEB*, XXVIII, p. 7.
(Calcium arsenite, Paris green.)

Natural Enemies *

- 879 BISHOP, S. C. and HART R. C. (1931) Natural enemies of the mosquito in Colorado *J. N. Y. Ent. Soc.* XXXIX 2 pp 1st 157 *P4EB*, XIX p 18.
(Frogs, toads, salamanders, *Dolicho-*
poda Lyons, *diplopoda* snails)
- 900 GAGANOV A. P. (1938) Destruction of the eggs and larvae of mosquitoes by *Limnoria ovata* *Med. Pribl.* VII 4 p 617 *P4EB*, XXVII p 78
(Gastropod)
- 901 HICMAN E. H. (1934) Predators of the Culex (larvae, pupae and adults) *J. Trop. Med. Hyg.* XXXVII 9 pp 19-174 10, pp 14-150 *R4FB* XXII p 1-0
(Comp. chrysomelids, etc. 123 refs.)
- 902 SHAPIROV L. A. (1938) Value of aquatic birds as destroyers of anopheline larvae *Med. Pribl.* VII 5 p 778 *R4EB* XXVII p 125
- 903 TURBETT C. I. (1978) The giant toad *Agrotis J. F.* IX, 3 p 9 *P4FB* XXVII p 80
(*Bufo marinus* does not eat larvae)
- 904 TWINE C. P. (1931) Aquatic animal and plant enemies of mosquitoes *Can. Ent.* LXIII 3 pp 51-61 *P4FB* XIX p 148
(*Culex*, *Chalcidius*, *Encorethra*,
Dytiscus, *Nitidulidae*, *Hemiptera*, *Chironomidae*,
Flutellidae, *Utricularia*, *Lemna*, *Wolffia*,
Sparganium)
- 905 WLADIMIROV M. and SMIRNOV F. (1932) Aquatic insects that feed on Culex larvae *Zh. Anz.* XLIX 7-8 1p 192-200
R4EB XX p 731

Oil

- 906 ALLEN R. P. (1938) A summary of the literature on the repellent effect of oil on mosquito larvae *Proc. Oil Ann. C. of Mosq. Abolition Officers California* pp 21-35.
(Typecript)
- 907 BARROWMAN B. (1931) Anti-natural oils and their application. *Utah J.* VI, 1 pp 7-11
R4FB XIX p 145

*References to Deterrent Plants and Trees (Nos 723-731) and Fish (Nos 763-803) are listed under separate headings.

- 907 BLACKLOCK, D B (1938) A device for applying oil or other liquids to flushing cisterns *Ann Trop Med Paras*, XXVII, 2, pp 109-114 *RAEB*, XXVII, p 14
- 908 BOSE K (1931) Application of larvicides *Ind Med Gaz*, LXVI, 8, pp 436-440 *RAEB*, XI, p 225
- 909 CAUCHI J, *et al* (1936) A method of testing oils and other chemical agents for killing mosquito larvae *Bull Ent Res*, XXVII, 4, pp 649-652 *RAEB*, XXV, p 94
- 910 CLARK D (1931) Methods and approximate costs of malarial preventive measures on High Rock Lake in North Carolina *South Med J*, XXIV, 5, pp 442-449 (Figure of oil water spraying outfit)
- 911 CORBETT, G H, and HODGKIN E P (1931) Laboratory experiments on the larvicidal properties of mineral oils *Bull Inst Med Res, F M S*, No 5 20 pp *RAEB*, XV, p 199
- 912 GINSBURG, J M (1932) Airplane oiling to control mosquitoes *Science*, LXXV, 1951, p 542 *RAEB*, XX, p 176
- 913 GINSBURG, J M (1933) Specifications for mosquito oils and larvicides *J Agric Exp Sta Circ* No 291 4 pp *RAEB*, XXIII, p 25
- 914 GRAY, H F, and BENT F A (1938) The spreading properties of oils on water surfaces *Procs 9th Ann Conf Mosq Abatement Officials, California*, pp 2-16 (Typescript)
- 915 KALANDADSE, L (1933) Studies of the respiration processes in mosquito larvae, and of the effects of kerosene, shade and pollution of water *Arch Schiffs Trop Hyg*, XXXVII, 2 pp 88-103 *RAEB*, XVI, p 93 (How oil acts on larvae)
- 916 LIKER, C C (1936) Observation on mosquito control on Lake Wilson, Tennessee River *South Med J* XXIX, 8 pp 867-873 (Application of oil by power sprayers from boats)
- 917 KURO, M, and OHITO, E (1939) A new mosquito larvicide oil *J Orient Med*, XXVI, 3, pp 407-409 *TDB* XXVII, p 365 (Mixture of oil factory products)
- 918 MANSON, D. (1938) A method of anti larval oiling for rivers *Ind Med Gaz*, LXXIII, 5, pp 284-285 *RAEB* XXVI, p 181

- 919 MICHELSON F G (1938) An apparatus for continuous oiling of streams *Ind Med Gaz*, LXIII, 10, p 612 *RAEB XXVII* p 81
- 920 MILLER F W (1930) Mosquito control in New Jersey *60th Ann Ppt Ent Soc Ontario* 1929 pp 89-92 *RAEB XIV*, p 23 (Apparatus mounted on trucks)
- 921 MIYAK, O J (1933) Mosquito control in Haiti *C S A Agr Med Bull XXVI* 3 pp 323-334 *RAEB XXII* p 34 (Panama wheel for open ditches)
- 922 MURRAY, D R P (1936) Mineral oils as larvicides *Bull Ent Res XXVII* 2 pp 289-306 *RAEB XXIV* p 241
- 923 MURRAY, D R P (1938) Problems concerning the efficiency of oils as mosquito larvicides Part I The stability of oil films on the surface of the water *Ibid XXIX* 1 pp 11-35 *RAEB XXVI* p 135
- 924 MURRAY, D R P (1939) Problems concerning the efficiency of oils as mosquito larvicides Part II The spreading power of oils and methods of increasing it *Ibid XXX* 2 pp 211-236 *RAEB XXVII* p 259
- 925 POKROVSKI, N N, and DZHANGIROV, K G (1939) Application of oil from a mechanical sprayer *Mal Paras VIII* 1 pp 145-146 *RAEB XXVII* p 222
- 926 POWERS, G E, and HEADLEE T J (1939) How petroleum oil kills culicine mosquito eggs *J Econ Ent XXVII* 2 pp 219-222 *RAEB XXVII* p 216
- 927 QUAIFE W T (1931) The brushing method of oiling *Malay Med J VI* 3 pp 62-63 *RAEB XX* p 121
- 928 QUAIFE, W T (1934) Further remarks on the brushing method of oiling *Ibid IX* 1 p 33-43 *RAEB XXII* p 148
- 929 RAMSAY, G C, and CARPENTER J A (1932) An investigation of petrolum oils for malaria control purposes *Rec Mal Surv Ind III* 2 pp 203-218
- 930 RICE F M (1935) Anti-larval oil application by a one man brushing method *Ibid V* 4 pp 499-500
- 931 SAUTT F T (1929) Oils for mosquito control. *Chem Abst.* XXIII, 2 p 466
- 932 SICHAULT G, and MESSERLIN A (1936) Destruction of larvae by mineral oils and the activation of such oils *Bull Soc Path Exot XXIX* 9 pp 1023-1032 *RAEB XXV*, p 86

- 933 SINTON, J A, and MAJID, S A (1936) The dispersion of anopheline larvae by the flow of streams and the effect of larvicides in preventing this *Rec Mal Surv Ind*, V, 1, pp 3-17
- 934 STRONQUIST, F T (1929) Oil supplies for anti mosquito campaigns *U S A Pub Hlth Bull*, No 156, pp 123-124
- 935 SYMES, C B, and ROBERTS, J I (1934) Medical entomology *Rpt Med Res Lab Kenya*, 1932, pp 21-27 *RAEB*, XXII, p 169
- 936 SYMES, C B, and ROBERTS, J I (1935) Section of entomology *Ibid*, 1933, pp 16-18 *RAEB*, XXIII p 233
- 937 VAINSHTEIN, N V (1939) Behaviour of *Anopheles* larvae after oiling water containing vegetation *Med Paras*, VIII, 1, pp 118-122 *RAEB*, XXIII, p 201
- 938 VANNOTE, R L (1932) Use of the aeroplane for the distribution of oils and larvicides *Procs 19th Ann Mtg A J Mosq Ext Ass*, pp 97-101 *RAEB*, XX, p 240
- 939 VICKERS, W J, et al (1934) Economy in large scale anti malarial control in Kuala Lumpur, F M S *Trans Roy Soc Trop Med Hyg*, XXVIII, 1, pp 85-99 *RAEB*, XXII, p 174
(Brushing method of oiling)
- 940 WALLACE R B (1932) Cessation of oiling in a malarious area during a certain season of the year *Malay Med J*, VII, 3, pp 73-81 *RAEB*, XXI, p 8
- 941 WATSON, G I (1937) Observation on mosquito larvae dying in anti malaria oils and other substances *Ann Trop Med Paras*, XXXI, 3, pp 417-426 *RAEB*, XXVI, p 38
- WOOD, T F M (1932) See No 764
(Modification of Wetmore's apparatus)

Paris Green.

- 942 AZIZ, M (1939) The water paris green mixture in anopheline control in the Tiliwa area, Cyprus *Med Dept Papers Govt Cyprus*, No 5, 16 pp
(Barber's method)
943. BARBER M A, RICE, J B, and MANDELOS, A (1936) A dustless method of diluting and spreading paris green *Amer J. Hyg*, XXII, 1, pp 41-44 *RAEB*, XXIV, p 289

941 BELON Y M (1931)

942 BENEDETTI A DE (1933)

BENEDETTI A DE (1934)

BENEDETTI A DE (1934)

BENEDETTI A DE (1934)

949 BENEDETTI A DE (1935)

950 BENEDETTI A DE (1935)

BOSE K (1931)
BOSE R (1933)

BRANNING CRACKEN B (1938)

BUDUILIN V (1931)
VODREKVA V V (1931)

CAMPBELL M S (1940)

CLARK D (1931)
COOK N S (1931)

Method of applying paris green in air
and safety of its application on a large
scale *Tr P M 11* IX 1 pp 13 18
RIFB VI p 114

A method of using paris green
while the last arrivals of paris green
Pr Med VII 1 11 *RIFB*
VII p 14

A method of using paris green
in the last arrivals of paris green
Pr Med VII 1 11 *RIFB*
VII p 14

For the use of paris green
in the last arrivals of paris green
RIFB VII 1 11 *RIFB*
VII p 14

Method of using paris green
in the last arrivals of paris green
Pr Med VII 1 11 *RIFB*
VII p 14

Application of paris green
in the last arrivals of paris green
Pr Med VII 1 11 *RIFB*
VII p 14

Application of paris green
in the last arrivals of paris green
Pr Med VII 1 11 *RIFB*
VII p 14

Application of paris green
in the last arrivals of paris green
Pr Med VII 1 11 *RIFB*
VII p 14

Application of paris green
in the last arrivals of paris green
Pr Med VII 1 11 *RIFB*
VII p 14

Application of paris green
in the last arrivals of paris green
Pr Med VII 1 11 *RIFB*
VII p 14

Application of paris green
in the last arrivals of paris green
Pr Med VII 1 11 *RIFB*
VII p 14

Application of paris green
in the last arrivals of paris green
Pr Med VII 1 11 *RIFB*
VII p 14

Application of paris green
in the last arrivals of paris green
Pr Med VII 1 11 *RIFB*
VII p 14

- 956 COVELL, G (1935) The effect of paris green dusting on rice crops *Rec Mal Surs Ind*, V, 2, pp 153-157
- 957 COVELL, G and AFRIDI M K (1937) Experimental application of paris green from aircraft *Ibid*, VII, 1, pp 93-103
- 958 COVELL, G BAILY J D and PRASAD V (1935) An experiment with paris green in a hyperendemic village in Sind *Ibid*, V, 2, pp 131-152
- 959 DANILCHENKO (1937) Aeroplane dusting in Kiev in 1936 *Med Paras*, VI, 3, p 446 *RAEB* XVI p 169
- 960 DUNDALE J N (1936) Malaria in Portugal *J Trop Med Hyg* XXIX, 13, pp 149-150 *RAFB* XXIV, p 229
(Said to injure growing rice)
- 961 FIGUEIRA L, and LANDEIRO F (1939) Report of the first year's anti malarial work at Benavente *Arg Inst Bui Cam Pest*, VI, 3, pp 191-243 *RAEB*, XVI, p 59
(Dusting from aircraft)
- 962 GRANT H G, NEWMAN, B M, and WOOD P D. (1932) Action of colloidal paris green on the larvae of *Culex appicalis* *U S A Pub Hlth Rpts*, XLVII, 23, pp 1239-1247 *RAEB* IX, p 175
- 963 HAKKI I (1931) Mosquitoes of Turkey and control of anopheline breeding *Arch Schiffs Trop Hyg*, XXXV, 5 pp 303-309 *RAEB*, XI, p 180
- 964 HALE C (1932) Anopheles mosquito control 20th Ann Rpt United Fruit Co, 1931, pp 98-107 *RAEB* XII p 233
- HENDERSON J M and HOWARD R S (1939) See No 846
- 965 KEIRIS, L M, and KLOKOV, N A (1939) Use of aqueous suspensions of paris green *Med Paras* VIII, 3 pp 303-306 *RAEB*, XXVIII p 37
(Barber's method)
- KIKER, C C (1936) See No 916
- 966 KIKER C C, et al (1938) Further observations on airplane dusting for Anopheles larvae control *South Med J*, XXXI, 7, pp 808-811 *RAFB*, XXVII, p 30
- KING, W V, and MCKEEL, T F (1938) See No 853
- 967 LANDEIRO, F (1933) A sifter for paris green dust *Rev Mal*, VII, 6, pp 1165-1168 *RAFB* XII p 97

- 958 LEZSOV, H S (1931) Anopheline mosquitoes in Southern Rhodesia *Lond Sch Tr p Med Memoir* No 4
(Cyanide dust as diluent)
- 959 FROEYDREK, F (1934) Application of larvicidal powders from aircraft in Madagascar *Bull Soc Path Exot* XXVII 6 pp 603-608 *RAEB*, XXII p 190
- 970 MACKAY, R. *et al* (1934) Arsenical poisoning associated with larvicidal treatment of water with paris green *Trans Roy Soc Trop Med Hyg* XXVII 5 pp 517-523 *RAEB*, XXII p 84
- 971 MARTINI, E (1931) Dusting especially with paris green in malaria control *Z Desinfekt* XXIII 4 pp 151-166 *RAEB* XIX p 107
- 972 MONVIER, E, and MARTIN, P (1934) Chemical estimation of paris green *Bull Soc Méd Chir Indochine* VII 2 pp 253-267 *RAEB* XXII p 27
- 973 MORIN, H G S. *et al* (1933) The mode of action of paris green *Bull Soc Path Exot* XXVI 10 pp 1267-1273 *RAEB* XXII p 31
- 974 MORIN, H G S. and MARTIN, P (1933) Value of paris green as a larvicide in Indo China *Arch Inst Past Indochine*, XVII pp 103-111 *RAEB* XXII, p 91
- 975 NABOKOV, V A (1931) The use of aeroplanes in the control of mosquito larvae *Trop Med Vet* IX 7-8, pp 395-397 *PIFR* XX p 132
- 976 NABOKOV, V A (1932) Control of larvae of malaria mosquitoes *Mater 1st All Union Auto Chem Conf Moscow* 1930 pp 55-83 *RAEB* XX, p 152
- 977 NABOKOV, V A (1934) Results of dusting paris green from aircraft in 1931 *Med Paras* III 1 pp 63-73 *RAEB* XXII, p 194
- 978 NABOKOV, V. A (1939) Results of spraying paris green from aircraft in the U S S R *Ibid* VIII 2 pp 165-169 *RAEB* XXVIII p. 7
- 979 NABOKOV, V A *et al* (1931) Use of aeroplane in larval control *Trop Med Vet* IX 6 pp 270-288 *PIFR*, XX p 116
- 980 PALADINO BLANDINI, A (1933) The organization and first results of general anti mosquito work in Calabria *Bull Mal* VII 1, pp 119-195 *RAEB* XXI, p 249
(Compulsory application by 11 000 l/m²)

- PANAGIA, A (1932) See No 784.
(Paris green said to attract by its colour)
- 981 PAUL, W L C, and GOMFY W A (1938) The paris green treatment of jaddy fields in anti malarial work *Trop Agric*, XC, 2, pp 88-90 *RAEB*, XXVI, p 141
- 982 POPOV V M, and FERRI I V (1936) Action of paris green on larvae of *A maculipennis* at low temperatures *Med Paras*, V, 4, pp 506-509 *RAEB*, XXV, p 141
- 983 RAO B A (1938) Control of Anopheles breeding in irrigation channels by paris green *Proc Nat Inst Sci Ind*, IV, 2 pp 203-209 *RAEB* XXVII, p 7
- 984 RAO, B A, and SWART W C (1937) Paris green and jaddy *Rec Mal Surv Ind*, VII, 2 J, pp 185-189
- 985 RICE F M, and ROY B K D (1938) A rapid method for the estimation of the arsenious oxide content of paris green *J Mal Inst Ind*, I, 2 pp 123-128
- 986 RUKAVISHNIKOV B I (1936) Scorching of crops by larvicidal dusts applied from aircraft *Med Paras*, V, 3, pp 426-437 *RAEB*, XXV, p 113
(Attributable to faulty technique)
- 987 RUSSELL, P F (1933) Automatic distribution of paris green for malaria control *J Parasit*, XIX, 3 pp 215-224 *RAEB*, XXI, p 117
- 988 RUSSELL, P F (1935) The automatic distribution of paris green for control of Anopheles larvae *Trans 9th Cong F F A T M*, II, pp 229-230 *RAEB*, XXVI, p 126
- 989 RUSSELL, P F and BATON L S (1934) An automatic distributing machine for paris green mixtures *Phil J Sci*, LIII, 4, pp 497-503 *RAEB*, XXVII, p 223
- 990 RUSSELL P F and JACOB V P (1939) Treating *Casuarina* pits with paris green *J Mal Inst Ind*, II, 3, pp 261-271
- 991 RUSSELL, P F KNIFE P W, and RAO T R (1940) Water instead of dust for diluting paris green *Ind Med Gaz*, LXXV, 12 pp 740-742
- 992 RUSSELL, P F, and WEST A P (1932) Charcoal as a diluent for paris green *Phil J Sci*, XLVIII, 3, pp 291-296 *RAEB*, XX, p 215
- 993 RUSSELL, P F and WEST A P (1932) The effect on *Culex* larvae of paris green diluted with charcoal *Ibid*, XLIX, 4, pp 651-675 *RAEB*, XXI, p 86

- PANAGIA, A (1932) See No 784.
(Paris green used to attract by its color)
- 981 PAUL, W R C, and COMPTON, W A (1938) The paris green treatment of paddy fields in anti malarial work *Trop Agric*, XC, 2, pp 88-90 *RAEB*, XXVI p 141
- 982 POROV, V M, and FRUM, I V (1936) Action of paris green on larvae of *A. maculipennis* at low temperatures *Med Paras*, V, 4 pp 506-509 *RIEB*, XXV, p 141
- 983 RAO, B A (1938) Control of Anopheles breeding in irrigation channels by paris green *Procs Nat Inst Sci Ind*, IV, 2, pp 203-209 *RIEB*, XXVII, p 7
- 984 RAO, B A, and SWEET, W C (1937) Paris green and paddy *Rec Mal Surv Ind*, VII, 2-3 pp 185-189
- 985 RICE, E M, and ROY, B K D (1938) A rapid method for the estimation of the arsenious oxide content of paris green *J Mal Inst Ind* I, 2, pp 123-128
- 986 RUKAVISHNIKOV, B I (1936) Scorching of crops by larvicidal dusts applied from aircraft *Med Paras*, V, 3 pp 426-437 *RAEB*, XXV, p 113
(Attributable to faulty technique)
- 987 RUSSELL, P F (1933) Automatic distribution of paris green for malaria control *J Parasit*, XIX, 3 pp 215-224 *RAEB* XXI, p 117
- 988 RUSSELL, P F (1935) The automatic distribution of paris green for control of Anopheles larvae *Trans 9th Cong F F A T U*, II, pp 223-232 *RAEB*, XXVI, p 126
- 989 RUSSELL, P F and FAYON, L S (1934) An automatic distributing machine for paris green mixtures *Phil J Sci*, LIII, 4, pp 497-503 *RAEB*, XXII p 223
- 990 RUSSELL, P F and JACOB, V P (1939) Treating *Casuarina* pits with paris green *J Mal Inst Ind*, II, 3, pp 261-271
- 991 RUSSELL, P F KNIFE, F W and RAO, T P (1940) Water instead of dust for diluting paris green *Ind Med Gaz*, LXXV, 12 pp 740-742
- 992 RUSSELL, P F, and WEST, A P (1932) Charcoal as a diluent for paris green *Phil J Sci*, XLVIII, 3, pp 291-296 *RAFR*, XX, p 215
- 993 RUSSELL, P F and WEST, A P (1932) The effect on *Culex* larvae of paris green diluted with charcoal *Ibid*, XLIX 4 pp 651-675 *RAEB*, XXI, p 86

- 994 SAVAGE J de la M (1933) Malaria sect on *Rpt Annua Med Pes*
oc 1933 7 pp PAEL VII
p 141
(Recebran a bluent)
- 995 SAZONOV I V and
STREGLINA F S (193) Control of mosquito larvae in Armenia
Mater Ist Ill Un on Ar o Chem C of
Moscow 193 pp 183-210 R4EB
XX p 10
(Aircraft dust n)
- 996 SEN P (1933) an paris green k l hsh? *Fengal Pub*
Ill h J 1 1 pp 9 1^o TDB
XXXXVII p 303
- 997 SERGEYEV I T (1932) Use of paris green as a larvicide in Algeria
Arch Inst Lat Algé X 3 p1 353-
38 R4EB XXI p 30
- 999 SERGIEV I G et al (1939) A mechan al sprayer for larvichidial
d ts Berna) *Med Ia at* VIII
1 pp 99 3 R4FB XXVIII
p 36
Mounted on bo t or tru k and worked
by a motor
- SHIKI V N I and
SICHENNAKOVA O A (1937) No 8
Effect on rice crop
- SHIPITZINA N K (1939) No 8 6
- 999 SICACLY G (1933) Appl at n f par green from a r salt
n Morocco *Bull Inst Hyg Marre*
- pp 93 1 r R4EB XXIV
1 14
- 1000 SICACLY G (193) Manufa ture tr a l use f j m
green larvicide mixtures n Morocco
Bull Soc Pa k Exot XXX " pp 84-
58 TDB XXXIV 1 310
- SILVER J F (193) No 610 (R pt for 1931)
(Aircraft dusting)
- 1001 SWEET W C and IIA
B A (1933) Notes on malaria in Mysore State IV
Experimental control with paris green
and plasmodium *Lee Mal Surv Ind*
III 4 pp 689 718
- 1002 SWEET W C and IIA
B A (1931) Notes on malaria n Mysore State V
The control of anopheline breed n in
Bangalore City and its cost in Mysore
State *R Mal Surv Ind* IV 2,
1 p 90-110
- 1003 SYDDIQ M M (1939) Observat ns of pract al importance for
the malaria logist *Ind Med Gaz*
LXXXIII II pp 6 6-670 R4FP
XXXXII p 173
(Effects on rice crops.)

- 1004 SYMES, C B (1931 32) Observations on anophelines and malaria in Kitale, with notes on experimental control with paris green *Kenya and E Afr Med J* VIII 9-10, pp 256-267 280 283 *RAEB*, IX, p 76
- 1005 VARIOUS AUTHORS (1932) First conference in the U S S R on use of aircraft in malaria control *Med Paras*, I 2, pp 65-116 *RAEB*, XII, p 87
- 1006 WALLACE, R B (1933) Paris green as a larvicide on an inland hilly estate in Malaya *Trans Roy Soc Trop Med Hyg* XXVII, 2 pp 131-146 *RAEB*, XXI, p 229
- 1007 WATSON, R B (1936) Airplane dusting for Anopheles larvae control *South Med J*, XXIX 8 pp 862 867 *RAEB*, XXV, p 117
- WATSON R B and BISHOP, E L (1940) *See* No 709 (Aircraft dusting)
- 1008 WATSON R B KILMER, C C, and JOHNSON, H A (1938) The role of airplane dusting in the control of Anopheles breeding associated with impounded waters *U S A Pub Hlth Rpts* LIII, 7 pp 261-263 *RAEB* XXVI, p 132
- 1009 WEST A P and RUSSELL, P F (1937) Paris green partially adsorbed on charcoal as a larvicide *Phil J Sci* XLVIII, 4, pp 545-561 *RAEB*, XX, p 236
- 1010 YAO Y T, and WU C C (1935) Anti larval measures by the use of paris green *Trans 9th Cong F E A T M* II, pp 213-221 *RAEB*, XXVI, p 126
- YURCHAK, F F, and BOZHENKO, V P (1939) *See* No 898 (Barber's method)

Repellents

- GINSBURG J M (1935) .. *See* No 842 (Pyrethrum emulsion)
- 1011 GINSBURG J M (1935) Protection from mosquito bites in outdoor gatherings *Science* LXXXII, 2134, pp 490-491 *RAEB* XXIV, p 34 (Pyrethrum emulsion)
- 1012 GINSBURG, J M (1936) Protection of outdoor gatherings from the mosquito pest *Proc 23rd Ann Mtg N J Mosq Ext Ass* pp 166-173 *RAEB* XXIV, p 270 (Pyrethrum emulsion)
- 1013 GINSBURG, J M (1937) Protection of outdoor meetings from the mosquito pest *Ibid* 24th Mtg pp 5-11 *RAEB* XXV, p 263 (Pyrethrum emulsion)

- 1014 GRANETT, P (1933) Comparison of mosquito repellency tests under laboratory and field conditions *Procs 25th Ann. Mtg N J Mosq Ext Ass.* pp 61-57 *RAEB*, XXVII, p 29
- 1015 GRANETT, P (1940) The development of a practical mosquito repellent *Ibid*, 27th Mtg, pp 36-43
(Stay away Insect Repellent Lotion*, mixture of diethylene glycol monobutyl ether acetate, diethylene glycol monoethyl ether, ethyl alcohol, corn oil and perfume)
- 1016 KING W V *et al* (1937) Experiments in Florida in repelling mosquitoes by outdoor spraying *Ibid*, 24th Mtg, pp 163-172 *RAEB*, XXV, p 265
(Pyrethrum emulsion)
1017. MACKEY C G (1938) An effective repellent for biting insects *Canad Entom*, LXX, 8, pp 175-176 *RAEB*, XXVII, p 14.
(Oil of thyme, $\frac{1}{2}$ oz, Extract of pyrethrum 1 oz Castor oil, 23 oz)
- 1018 MACKAY C G (1939) Studies on repellents for biting flies *Ibid*, LXXI, 2, pp. 38-41
- 1019 MAIL, G V (1934) The mosquitoes of Montana *Bull Montana Agric Exp Sta.* No 288 72 pp *RAEB*, XXII, p 182
(Derivatives of salicylic acid)
- MEILLON B DE (1934) See No 1039
(Citronella oil)
- SYDDIQ M M (1934) See No 1003
(Beeswax, 2 dr spermaceti 4 dr, coconut oil 4 dr fuller's earth, 4 dr lq paraffin, 1 oz, oil of citronella, 1 oz carbolic acid, 10 min)
- 1020 VANNOTE, R L. (1937) Methods of applying the larvicide as a repellent *Procs 24th Ann Mtg N J Mosq Ext Ass.* pp 11-13 *RAEB*, XXV p 263

Screening.

- 1021 BEANLEF C (1934) On anti mosquito screens *Ric Mal.*, XVII, 1, pp 51-61 *RAEB*, XXVI, p. 159
(Steel, aluminium, monel metal, bronze, galvanized iron; brass; copper)

- 1004 SYMES C B (1931-32) Observations on anophelines and malaria in Kisale, with notes on experimental control with paris green *Kenya and E Afr Med J*, VIII 9-10, pp 256-267 280 283 *RAEB*, XV p 76
- 1005 VARIOUS AUTHORS (1932) First conference in the U S S R on use of aircraft in malaria control *Med Paras*, I 2, pp 65-116 *RAEB*, XXII p 87
- 1006 WALLACE, R B (1933) Paris green as a larvicide on an inland hilly estate in Malaya *Trans Roy Soc Trop Med Hyg*, XXVII 2 pp 131-146 *RAEB*, XXI, p 229
- 1007 WATSON, R B (1936) Airplane dusting for *Anopheles* larvae control *South Med J*, XXIX 8 pp 862 867 *RAEB*, XXV, p 117
- WATSON, R B and BISHOP, E L (1940) *See No 709*
(Aircraft dusting)
- 1008 WATSON R B KIKER C C, and JOHNSON H A (1938) The role of airplane dusting in the control of *Anopheles* breeding associated with impounded waters *U S A Pub Hlth Rpts*, LIII 7 pp 251-263 *RAEB*, XXVI, p 132
- 1009 WEST A P and RUSSELL, P F (1932) Paris green partially adsorbed on charcoal as a larvicide *Phil J Sci* XLVIII 4, pp 545-561 *RAEB*, XX, p 236
- 1010 YAO Y T, and WU, C C (1935) Antilarval measures by the use of paris green *Trans 9th Cong F E A T M*, II pp 213 221 *RAEB* XXVI, p 126
- YURCHAK, F F, and BOZHENKO V P (1939) *See No 898*
(Barber's method)

Repellents

- GINSBURG J M (1935) .. *See No 842*
(Pyrethrum emulsion)
- 1011 GINSBURG J M (1935) Protection from mosquito bites in outdoor gatherings *Science* LXXII, 2134, pp 490-491 *RAEB*, XXIV, p 34
(Pyrethrum emulsion)
- 1012 GINSBURG, J M (1936) Protection of outdoor gatherings from the mosquito pest *Proc 23rd Ann Mtg N J Mosq Ext Ass*, pp 166-173 *RAEB*, XXIV, p 270
(Pyrethrum emulsion)
- 1013 GINSBURG J M (1937) Protection of outdoor meetings from the mosquito pest *Ibid* 24th Mtg pp 5-11 *RAEB*, XXV p 265
(Pyrethrum emulsion)

- 1014 GRAYETT P (1939) Comparison of mosquito repellency tests under laboratory and field conditions *Procs 25th Ann Mtg N J Mosq Ext Ass*, pp 51-57 *RAEB*, XXVII p 29
- 1015 GRAYETT P (1940) The development of a practical mosquito repellent *Ibid* 27th Mtg, pp 36-43
(Stay away Insect Repellent Lotion' mixture of diethylene glycol monobutyl ether acetate diethylene glycol monoethyl ether, ethyl alcohol, corn oil and perfume)
- 1016 KING W V *et al* (1937) Experiments in Florida in repelling mosquitoes by outdoor spraying *Ibid* 24th Mtg pp 163-172 *RAEB* XXV, p 263
(Pyrethrum emulsion)
- 1017 MACNAY C G (1938) An effective repellent for biting insects *Canad Entom LXX* 8 pp 175-176 *RAEB* XXVII p 14
(Oil of thyme, $\frac{1}{2}$ oz Extract of pyrethrum 1 oz Castor oil, ~ 3 oz)
- 1018 MACNAY C G (1939) Studies on repellents for biting flies *Ibid*, LXXI 2 pp 78-81
- 1019 NATH C A (1934) The mosquitoes of Montana *Bull Montana Agric Exp Sta* No 283 72 pl *RAEB* XXII p 182
(Derivatives of salicylic acid)
- MILLON B E (1934) See No 1059
(Citronella oil)
- SYDDIQ M M (1934) See No 1003
(Beeswax 2 dr spermaceti 4 dr, coconut oil 4 dr fuller's earth, 4 dr 1 lb paraffin, 1 oz, oil of citronella 1 oz carbolic acid 10 min)
- 1020 VANNOTE R I (1937) Methods of applying the larvicide as a repellent *Procs 24th Ann Mtg N J Mosq Ext Ass* pp 11-13 *RAEB*, XXV p 283

Screening

- 1021 BREXLER C (1934) On anti mosquito screens *Pir Mol*, XVII 1 pp 51-61 *RAEB*, XXVI, p 159
(Steel aluminum monel metal bronze galvanized iron brass; copper)

- 1004 SYMES, C B (1931-32) Observations on anophelines and malaria in Kitale, with notes on experimental control with paris green *Kenya and E Afr Med J*, VIII, 9-10, pp 256-267, 280-283 *RAEB*, XX, p 76
- 1005 VARIOUS AUTHORS (1932) First conference in the U S S R on use of aircraft in malaria control *Med Paras*, I, 2, pp 65-116 *RAEB*, XXII, p 87
(RUSSIAN)
- 1006 WALLACE, R B (1933) Paris green as a larvicide on an inland hilly estate in Malaya *Trans Roy Soc Trop Med Hyg*, XXVII, 2, pp 131-146 *RAEB*, XXI, p 229
- 1007 WATSON, R B (1936) Airplane dusting for *Anopheles* larvae control *South Med J*, XXIX, 8, pp 862-867 *RAEB*, XXV, p 117
- WATSON, R B, and *See No 709*
BISHOP, E L (1940) (Aircraft dusting)
- 1008 WATSON, R B, KIKER, C C, and JOHNSON, H A (1938) The role of airplane dusting in the control of *Anopheles* breeding associated with impounded waters *U S A Pub Hlth Rpts*, LIII, 7, pp 251-263 *RAEB*, XXVI, p 132
- 1009 WEST, A P, and RUSSELL, P F (1932) Paris green partially adsorbed on charcoal as a larvicide *Phil J Sci*, XLVIII, 4, pp 545-561. *RAEB* XX, p 276
- 1010 YAO Y T, and WU, C C (1935) Anti larval measures by the use of paris green *Trans 9th Cong F E A T M*, II, pp 213-221 *RAEB*, XXVI, p 126
- YURCHAK, F. F, and *See No 898*
BOZHENKO, V P (1939) (Barber's method)

Repellents

- GINSBURG J M (1935) .. *See No 842*
(Pyrethrum emulsion)
- 1011 GINSBURG J M (1935) Protection from mosquito bites in outdoor gatherings *Science*, LXXII, 2134, pp 490-491 *RAEB*, XXIV, p 34
(Pyrethrum emulsion)
- 1012 GINSBURG, J M (1936) Protection of outdoor gatherings from the mosquito pest *Proc 23rd Ann Mtg N J Mosq Fct Ass*, pp 166-173 *RAEB*, XXIV, p 270
(Pyrethrum emulsion)
- 1013 GINSBURG, J M (1937) Protection of outdoor meetings from the mosquito pest *Ibid*, 24th Mtg pp 5-11 *RAEB*, XXV, p 265
(Pyrethrum emulsion)

1014 GRANETT, P (1934)

Comparison of mosquito repellency tests under laboratory and field conditions
Procs 25th Ann Mtg N J Mosq Ext Ass, pp 51-57 *PAEB*, XXVII, p 29
 The development of a practical mosquito repellent *Ibid*, 27th Mtg, pp 36-43
 (b) a) way Insect Repellent Lotion', mixture of diethylene glycol monobutyl ether acetate, diethylene glycol monobutyl ether, ethyl alcohol, corn oil and perfume)

1015 GRANETT P (1940)

Experiments in Florida in repelling mosquitoes by outdoor spraying *Ibid*, 24th Mtg, pp 163-172 *RAEB*, XXV, p 263
 (Pyrethrum emulsion)
 An effective repellent for biting insects
Canad Entom LXX, 8, pp 175-176.
IAFB, XXVII p 14
 (Oil of thyme, 1 oz. Extract of pyrethrum 1 oz. Castor oil, 23 oz)

1016 KING W V et al (1937)

Studies on repellents for biting flies
Ibid, LXXI, 2, pp 39-41
 The mosquitoes of Montana *Bull Montana Agric Exp Sta*, No 283
 "2 pp *RAEB*, XXII, p 182
 (Derivatives of salicylic acid)

17. MACNAY C G. (1934)

See No 1059
 (Citronella oil)

1018 MACNAY C G (1937)

See No 1003
 (Beeswax, 2 dr spermaceti 4 dr. coconut oil 4 dr fuller's earth, 4 dr liq paraffn, 1 oz., oil of citronella, 1 oz carbolic acid, 10 min)

1019 MAH G A (1934)

Methods of applying the larvicide as a repellent *Procs 24th Ann Mtg N J Mosq Ext Ass*, pp 11-13 *PAEB*, XXI, p 263

MEILLON R DE (1934)

STODOL V M (1934)

1020 VANKOTZ, R L. (1937)

Screening

On anti mosquito screens *Ric Mal*, XVII, 1 pp 51-61 *RAEB*, XXVI, p 159
 (Steel aluminium monel metal bronze, galvanized iron brass, copper)

21. DESLAFER C (1934)

- 1022 BLACKLOCK, D B (1935) .. Screencloth for houses in the tropics
Ann Trop Med Paras, XXIX, 2
pp 261-263 *RAEB*, XXIII, p 229
(Recommends Barronia metal)
- 1023 BLACKLOCK, D B (1937) Screencloth for houses in the tropics
Ibid, XXXI, 3, p 447. *RAEB*,
XXVI, p 39
- 1024 BOOKER, C G (1935) *Ann Rpt S Afr Rlways and Harbours*
Hlth Organization Rpt Dept Pub
Hlth S Afr, 1934-35, pp 79-88
RAEB, XXIV, p 77
(Monel metal screening, 16 mesh, 30
ISWG)
- 1025 BROWN J Y (1934) Safe mosquito nets for use in Nigeria
N Afr Med J, VII, 4, pp 147-148
RAEB, XXII, p 213
(Mesh 25/26)
- 1026 DAVEY, T H, and GORDON, R M (1938) The size of aperture necessary in screen
cloth *Ann Trop Med Paras*, XXXII,
4, pp 413-424 *RAEB*, XXVII, p 130
(Must not exceed 0.017 in., i.e., 16
mesh, 28 ISWG)
- 1027 FARLE W C (1932) Anti mosquito screening and screening
materials *Puerto Rico J Pub Hlth*
Trop Med, VIII, 2, pp 227-234
RAEB, XXI, p 89
(Bronze, monel metal; galvanized iron
recommends 12 mesh screen of
0.015 in wire)
- 1028 GATER B A R (1930) Mosquito nets *Malay Med J*, V, 1,
pp 29-32 *RAEB*, XIX, p 5
(Recommends 22/23 mesh and 60/100
cotton)
- 1029 GIGLIOTTI, G (1933) Aluminium gauze for mosquito screening
Riv Mal, XII, 1, pp 196-197 *RAEB*,
XXI, p 181.
- HARREAVFS, H (1936) See No 845
(Recommends 14 mesh, 30 ISWG wire
gauze, or 25/26 mesh nets woven of
30 or 40/60 counts cotton)
- 1030 IREDELL A W (1933) Mosquito proofing carried out by the
Royal Air Force in India *Jl F A*
M. C, IX, 1, pp 33-37 Also in
Procs Roy Soc Med, XXVI, 1, pp 1-3
RAEB, XVI, p 65
- 1031 MARNITTE, H (1935) The mosquito net in malaria prevention
in Indo China *Bull Soc Med Chir*
Indochine, VIII, 9, pp 1265-1297
RAEB, XXIV, p 126

ACCEPTING SPRAYS

- 1037 MARTINOVSKII F I the n f r a mosq to net Bull Soc
Pth Expt XXVI 8 pp 1018-1020
(1933-1934) Also in U I Inras III 4 Pl 313
314 R I F B XXII p 15
- 1038 MELENEY H E and Mosq to pro fine a l treatment n the
Griffin R B (1931) con trol of malar a on a plantation.
South Med J XXIV 5 pp 418-44
- 1039 MOORE J I (1931) Possibilit a of mosq to-prooing all rural
homes n a county Id J pp 431-439
- 1040 MULLIGAN H W and 18 a y f w re gauzo screening of
Majid S A (1931) different apertures n ex lusion, ano-
pheline n n eq wa n In A R c Vol
Surv Int III 1 pp 163-169
(Aperture of 0.015 to 0.030 is necessary
for absolute ex lusion but 0.030 is
con trol adequate for general use)
Man fac re and lfo of screen w re
South Med J XXIV 5 pp 449-451
P I F B XX p 1
(Recom mends electro galvan ed steel
w re)
- 1041 POTTALL I F and Mosq to net for use n the Pl l pp ne
A M (1934) 1 lanl Phil J Agr Liff "
pp 107-140 P I F B XXII p 146
(Snamay ne ing made from fibres of
Musa text ile)
- 1042 SHMELEVA Y D and Val e of a ree ng ho es and using
S LENOVA M F (1937) n mosq to n te Med P os VI "
pp 25-54 R I F B XXVI p 48
- 1043 ZAVATTARI E (1936) A amplified a th mosq o net for tropical
con trols R c M l XIV 3
pp 29-36 P I F B XXIII p 287

Ssprays

- Dept Ent mology and Zoology Ann
Rpt Tech e Dept 1936 Hs 1
1936 29 R II No 17
(1 e brom-carbon tetrachloride-petrol
spray)
- 1044 AZOV (1938) Destruct m of mosq to n in Aeroplanes.
A test Eight June 1 CCXXIV
p 1414 R I F B XXIII p 31
(Aq co a bane extract f pyrethrum)
Insect ul l sprays for the lstruction of
mosq itors and flies JI R A M C.
LXII 6 pp 411-418 R I F B XXII
p 19
(Pyrethrum kerosene ctrols o l
and petrol)

- 1043 BOER H DE (1937) *Anti mosquito measures in aerodrome*
Bull Off Int Hyg Pub XXIX 6
 pp 1157-1158 *RAEB* XXI p 53
 BOOKER C G (1935) *See No 1094*
 (Pyrethrum spray)
- 1044 BOOKER C G (1936) *Ann Rpt S Afr Rivers and Harbours*
Hlth Organ Rpt Dept P b Hlth
S Afr 1935-36 pp 93-101 *RAEB*
 XXI p 137
 (Pyrethrum spray)
- 1045 CALDWELL A F (1938) *A note on the chemistry and preparation*
of insect sprays containing pyrethrum
J Maln Br B t Med Ass I 4
 pp 336-341 *RAEB* XXVI p 159
- 1046 CHOPRA B L (1938) *Anti malarial measures in the railway*
area at Delhi Ind Med Gaz LXXIII
 3 pp 100-101 *RIFB* XXVI
 p 148
 (Pyrethrum spray)
- 1047 COVELL G (1940) *The cultivation of pyrethrum in*
India Planters Chronicle XXXV 18
 pp 370-371
- 1048 COVELL G MULLIGAN
 H W and AFRID M K
 (1938) *An attempt to control malaria by the*
destruction of adult mosquitoes with
insecticidal sprays J Mal Inst Ind
I 1 pp 105-113
 (Pyrethrum spray)
- 1049 GERTLER S I and
 HALLER H L (1939) *The pyrethrin content of home made*
fly sprays Soap XVI pp 93-94
RIFB XXVII p 155
- 1050 GNADINGER C B (1936) *Pyrethrum flowers Minneapolis Minn*
Ed (with Suppl)
- 1051 GRAY C G (1931) *Sensitivity to insecticides J Amer Med*
Ass XCVI 3 p 913
 (Dermatitis following use of Flit)
- GRIFFITHS T D H *et al* *See No 806*
 (1935) (Pyrethrum spray)
- 1052 HICKS E P and CHAND
 D (1936) *Transport and control of Aedes aegypti*
in aeroplanes Rec Mal Surv Ind
VI 1 pp 73-90
 (Pyrethrum spray)
- 1053 HOLT F L (1931) *A new mosquito spray Mal Surg on*
IXIX 6 pp 69-77 *RAEB* XX
 p 100
 (Pyrethrum with carbon disulphide
 acetone and chloroform)
- 1054 HOLT R L and KINTNER
 J H (1932) *Anti mosquito sprays Phil J Sc*
XLVII 4 pp 437-438 *RAEB* XX
 p 100

1055. HOYER, D G (1936) Effect of metals on fly sprays (Pyrethrum liquid insecticides) *Soap*, XII, Preprint No 1, 2 pp *R4EB*, XXIV, p 151.
1056. JITTA, N M J (1935) Destruction of mosquitoes in aircraft *Bull Off Int Hyg Pub*, XXVII, 7, pp 1360-1361 *R4EB*, XXIII, p 247 (Pyrethrum spray)
1057. LOWMAN, M S, and SULLIVAN, W N (1938) Pyrethrum evaluation. Pelition of pyrethrum content of pyrethrum flowers to their toxicity to mosquito larvae *Soap* XIV 11 pp 89-91, 93, 119 *R4EB* XXVIII, p 64
- McDANIEL F I (1939) See No 836 (Popular account)
1058. MACKIE, F P and CRABTREE, H S (1938) The destruction of mosquitoes in aircraft *Lancet*, CCXXXI pp 447-450 *R4EB*, XXVIII, p 118 (Aqueous base pyrethrum spray)
1059. MEILLON, B DE (1934) Observations on *Anopheles funestus* and *Anopheles gambia* in the Transvaal S Afr Inst for Med Res Pbn 249 pp (Pyrethrum spray)
1060. MEILLON B DE (1936) Control of malaria in S Africa by measures against adult mosquitoes in habitations. *Quart Bull Hlth Organ*, I of N, V, 1, pp 134-137 *R4EB*, XXIV, p 157 (Pyrethrum spray)
1061. MORRIS P (1939) Value of insecticidal sprays in malaria control *Rev Med Franc Extr Orient* XVI, 3, pp 264-274 *R4EB*, XXVII p 138 (Aqueous base pyrethrum spray)
1062. NIJKAMP, J A and SWELLENGREBEL, A H (1934) Destruction of *Anopheles* with Pyrethrum preparations *Ned Tijdschr Geneesk*, LXXXIII, 21, pp. 2327-2333 *R4EB*, XXII p 197
1063. PANNEWITZ, F (1931) On the toxicity of the combinations of mineral oils with essential oils in insecticides with a petroleum base. *J Desinfekt* XXIII, 8, pp 339-342 *R4EB*, XIV, p 251
1064. RICHARDS, H (1928) Report on the disinfection of aircraft at Khartoum *Bull Off Int Hyg Pub*, XX, 3, pp 563-567 *R4EB*, XXVI, p 157 (Pyrethrum extract, 58, citronella oil, 2 carbon tetrachloride, 49, kerosene oil, 432)

- 1065 RICHARDSON, H H (1932) Insecticidal studies of mid continent distillates as bases for pyrethrum extracts *Indust Engng Chem*, XXIV, 12, pp 1394-1397 *RAEB*, XXII, p 10
- 1066 ROSS, G A P (1936) Insecticide as a major method of control of malaria *Quart Bull Hlth Organ*, L of N, V, 1, pp 114-131 *RAEB*, XXIV, p 153
(Pyrethrum spray)
- 1067 ROSS G A P (1938) Automatic destruction of mosquitoes in aircraft *Bull Off Int Hyg Pub* XXV, 3, pp 2002-2031 *RAEB*, XXVII, p 48
(Aqueous base pyrethrum spray)
- 1068 RUSSELL, P F, and KNIFE, I W (1939) Malaria control by spray killing adult mosquitoes *J Mal Inst Ind*, II, 3, pp 229-237
(Pyrethrum spray)
- 1069 RUSSELL, P F, and KNIFE, I W (1940) Malaria control by spray killing adult mosquitoes Second season's results *Ibid*, III, 4, pp 531-541
(Pyrethrum spray)
- 1070 SCHOFFNER, W A P, and SWELLENGREBEL, N H (1938) Report for 1936 and 1937 of the Malaria Commission of the Health Council of Holland *Versl Meded Volk*, Reprint 26 pp *RAEB*, XXVII p 51
(Pyrethrum spray)
- 1071 SINTON, J A, and WAYS R C (1935) The efficacy of various insecticidal sprays in the destruction of adult mosquitoes *Rec Mal Surv Ind*, V, 3, pp 275-306
- 1072 SWELLENGREBEL, N H (1934) Seven years' experience in Anopheles control at Medemblik *Ned Tijdschr Geneesk*, LXXVIII, 3, pp 345-353 *RAEB*, XXII, p 116
(Kerosene with 0.5 per cent pyrethrum extract, 0.5 per cent oil of saesafra, and 2 per cent methyl salicylate)
1073. SWELLENGREBEL, N H, et al (1936) Investigations on the transmission of malaria in some villages north of Amsterdam *Quart Bull Hlth Org L of N*, V, 2, pp 295-332 *RAEB*, XXIV, p 283
(Spraying confined to houses with several children)

- 1074 SYMES, C E (1937) Insects in aircraft *Bull Off Int Hyg Pub*, XXIX, 6, pp 1150-1157 *RAEB*, XXV, p 253
(Pyrethrum kerosene-carbon tetrachloride)
- 1075 THORNTON, E V (1933-36) Malaria *Fpts Dept Pub Hlth S Afr*, 1932-35 *RAEB* XXII, p 53
XXIII p 8 XXIV, p 76, XXV, p 136
(Pyrethrum spray)
- 1076 VIKTOROV S V (1938) Destruction of adult mosquitoes *Med Paras*, VII 1, pp 61-63 *RAEB*, XXVI p 219
(Watery solution of soft natron soap and methylated spirit)
- 1077 WATS, R C, and BHARUCHA, A H (1940) The choice of mechanical sprayers for mosquitocides *J Mal Inst Ind*, III, 1, pp 123-136
- 1078 WATS, R. C, and SIMON, J (1937) Mosquitocidal value of indigenous *Derris* and other drugs *Rec Mal Surr Ind*, VII 1, pp 109-147
(Sprays containing 60 different substances)
- 1079 WELCH, L V (1939) Insects found in aircraft at Miami, Fla, in 1938 *U S A Pub Hlth Rpts*, LIV, 14, pp 561-566 *RAEB*, XXVII p 215
(Pyrethrum spray)
- 1080 WHITFIELD, F G S (1939) Air traffic insects and disease *Bull Ent Res*, XX 3, pp 365-442
- 1081 WILLIAMS, C L (1940) Disinsectionization of aircraft *U S A Pub Hlth Fpts* LV 23, pp 1003-1010
(Description of special spray gun)
- WILLIAMS, C. L., and DRESSER, W C (1935) See No 813
(Pyrethrum spray)
- 1082 WILLIAMS, C L., and DRESSER, W C (1935) A non inflammable pyrethrum spray for use in airplanes *U S A Pub Hlth Rpts*, L, 41 pp 1401-1404 *RAEB*, XXIV, p 33
(kerosene pyrethrum-carbon tetrachloride.)
- 1083 WILLIAMS L L (1932) Mosquito control activities of the United States Public Health Service *Procs 19th Ann Mtg N J. Mosq Ext. Ass.*, pp. 61-67 *RAFI*, XX, p 239
(Pyrethrum oil extract sprayed over marsh to kill adult mosquitoes, range 60 ft.)

Traps

- 1084 BRADLEY, G. H. and McNEEL, T. E. (1935) Mosquito collections in Florida with the New Jersey light trap *J. Econ. Ent.*, XXVIII, 5, pp 780-786 *RAEB*, XXIV, p 6
- 1085 BRIGHENTI, D. (1930) Research on the attraction effected by colours in *Anopheles maculipennis* *Rev. Mal.*, IX, 3 pp 224-231 *RAEB*, XIX, p 67
(Preference for red and violet, blue and green attract much less, grey neutral)
- 1086 CARMANAN, C. T. (1939) A two year record of adult mosquito trapping in Dade County, Florida *U. S. A. Pub. Hlth. Rpts.*, LIV, 15 pp 608-611 *RAEB*, XXVII, p 215
(Suction light traps)
- 1087 CHATTERJI, A. C., et al (1939) Report on tests made with the Entoray apparatus 25 pp Govt Press Calcutta
- 1088 DEERWERKER, R. J. VAN (1935) Mosquito traps *Procs. 22nd Ann. Mtg. N. J. Mosq. Ext. Ass.*, pp 164-170 *RAEB*, XXIII, p 205
(Suction light traps)
- 1089 GATER, B. A. R. (1933) Notes on Malayan mosquitoes, II *Malay. Med. J.* VIII, 1 pp 42-45 *RAEB*, XXI, p 147
(Bed trap catches)
- 1090 GORE, R. N. (1933) A feather duster mosquito trap 10 pp Bombay *RAEB*, XXII, p 17
- 1091 GORE, R. N. (1936) A village mosquito trap *Ind. Med. Gaz.*, LXXI, 8, pp 460-461 *RAEB*, XXV, p 5
- 1092 GORE, R. N. (1937) A modified village mosquito trap *Ibid.* LXXII, 11 pp 674-675 *RAEB*, XXVI, p 50
- 1093 GORE, R. N. (1937) An improved feather duster mosquito trap *Rec. Mal. Surv. Ind.*, VII, 2-3 pp 209-211
- 1094 GOURDON, G. (1930) The capture and destruction of insects by ultra violet rays *C. R. Cong. Int. Appar. Util. Lutte contre les ennemis des cultures* Lyon, 1929, pp 127-133, 209-213 *RAEB*, XXII, p 59
- 1095 HEADLEE, T. J. (1932) Development of mechanical equipment for sampling the mosquito fauna and some results of its use *Procs. 19th Ann. Mtg. N. J. Mosq. Ext. Ass.*, pp 106-126 *RAEB*, XX, p 241
(Suction light traps)

- 1096 HEADLEE, T J (1936) Mosquito control engineering. II Mosquitoes species and habits *Fungus News Rec* Aug 6 pp 199-201 (Suction light traps)
- JAMES J F (1935) See No 808 (Trapping and fumigation combined)
- 1097 JOHNSON H A (1937) Attractivity of light for Anopheles mosquitoes *J Tenn Acad Sc* VII 1 pp 104-106 *RAFB* XXVI p 105 (Best results with 100 watt frosted Mazda lamp)
- 1098 MACCREARY D (1939) Comparative results obtained by the use of several mosquito traps in a limited area *J Econ Ent* XXXII " pp. 216-219 *RAFB* XXVIII p 216 (Suction light traps)
- 1099 MAGDOFF T H (1935) A portable stable trap for capturing mosquitoes *RII Ent Res* XXVI 3 pp 363-371 *RAEB* XXIII p 300
- 1100 MILLER F W (1930) Light traps *Rpt. \ J Agric Exp Sta* 1929-30 pp 125-140 *RAFB* XIX p 57
- 1101 MURKIN T D (1934) A new development in mosquito traps *Procs 21st Ann Mtg \ J Mosq Ext Ass* pp 137-140 *RAEB* XXIII p 151 (Suction light traps.)
- 1102 PERAGALLO I (1938) A transparent trap for mosquitoes *Proc Mal* XVII 3 pp 231-234 *TDR* XXXVI p 16 (Constructed of acetyl-cellulose)
- 1103 RUSSELL P F and SANTIAGO D (1934) An earthen trap for anopheline mosquitoes *Procs Ent Soc Wash* XXXVI 1 pp. 1-21 *RAFB* XXVII p 65
- SELIVANOV K P (1937) See No 809 (Combined trapping and fumigation)
- 1104 SENIOR WHITE I A (1936) Experiments with an automatic mosquito catching machine *Rec Mal. Surv Ind* VI 4 pp 595-609 (Entomograph machine)
- 1105 STACK H H (1939) Use of the New Jersey mosquito trap in the Pacific North west *Procs 9th Ann. Conf Mosq Abatement Officials Cal* pp 63-64 (Typeset pt) (Suction light trap)

- 1106 SYDDIQ, M. M. (1938) .. Some observation with the Entoray machine (Abstract) *J Mal Inst Ind*, I, 4, p 473
- THOMPSON, T. O (1936) See No 811
(Combined trapping and fumigation)
- 1107 WATS, R C, and Experiments with the Entoray machine as an anti mosquito measure *Rec Mal Sur Ind*, VI, 1, pp 549-555
- BILDERBECK, C. L (1936)
- 1108 WELLS R W (1931) Electrified screens and traps *J Econ Ent*, XXIV, 6, pp 1242 1247 *RAEB*, XX, p 55

Zooprophylaxis.

- 1109 DAVIS, G E. and PHILIP, Identification of the blood meal in W African mosquitoes *Amer J Hyg*, C B (1931) XIV, 1 pp 130-141 *RAEB*, XIX, p 209
- 1110 FSCALAR, G (1933) Experimental application of zooprophylaxis at Arden (Agro Romano) *Ris Mal*, XII, 2, pp 373-380 *TDB*, XXX, p 866
(Pig sties)
- 1111 FGOAZZA, E (1933) Observations on anti malarial zooprophylaxis at Milan in 1932 *Ibid*, 3, pp 535-539 *RAEB*, XXI, p 261
- 1112 GRUIC, I (1932) The zoophilism of *Anopheles* as a factor in the disappearance of malaria *Glasn Trent Khig Zavoda*, VII, 4-6, pp 166-173 *RAEB*, XXI, p 59
- 1113 HACKETT, L W (1934) The present state of our knowledge of the subspecies of *A maculipennis* *Trans Roy Soc Trop Med Hyg*, XXVIII, 2, pp 109-140 *RAEB*, XXII, p 200
- 1114 HACKETT, L W, and Housing as a factor in malaria control MISSIROLI, A (1932) *Ibid*, XXVI, 1, pp 66-72 *RAEB*, XX, p 191
- 1115 HACKETT, L W, and The varieties of *A maculipennis* and their relation to the distribution of malaria in Europe *Rev Mal*, XIV, 1, MISSIROLI, A (1935) pp 45-109 *RAEB*, XXIII, p 152
- 1116 HU, S M K, and HU, H Blood preferences of *A. hyrcanus* v (1936 1937) *sinensis* in Shanghai region *Chinese Med J*, L, Suppt 1, pp 379-386, LI, 5, pp 639-642 *RAEB*, XXI, p 191, XXV, p 234
- 1117 KLIGLER, I J, MER, G, Seasonal variations in the food preferences of *Anopheles elutus* *Trans Roy Soc Trop Med Hyg*, XXVI, 3, pp 233-237 and OLITZKY, L (1932) *RAEB*, XXI, p 45

- 1118 MARTINI, F., MISSIROLI A. and HACKETT L W (1931) The races of *A. maculipennis* Arch. Sch. Hyg. Trop. Hyg. XXV, 11 pp 622-643 R4FB XX, p 57
- 1119 MISSIROLI, A. HACKETT L. W. and MARTINI E (1933) The races of *A. maculipennis* and their importance in the distribution of malaria. *Riv. Mat.*, XI, 1, pp 1-56 R4FB XXI p 177
- 1120 MORIN, H G S (1933) Maxillary index of Anopheles of Indo China. *Bull. Soc. Path. Exot.* XVI 2, pp 293-300 R4EB, XXI p 140
- 1121 MORISHITA K. and KATAOKA T (1937) Examination of blood in stomachs of Anopheles in Formosa. *Dokuts. Zasshi* XLV pp 90-92 R4EP XXI, p 158
- 1122 PECORI, G. and ESCALAR G (1973) Anti malaria work round home in 1934. *Riv. Mat.*, XIV 6 pp 469-519 R4EB, XXIV, p 163 (Fig. 11)
- 1123 PICK J B. and BARBER M A (1935) A modification of the Uhlenhuth Weitzel precipitin test for determining the source of blood meals in mosquitoes and other insects. *J. Lab. Clin. Med.* XX, 8 pp 876-883
- 1124 ROUBAUD F (1931) Anopheline zootropism. *Bull. Soc. Path. Exot.* XXV, 3, pp 209-216 R4EB, XIX, p 162
- 1125 ROUBAUD, F (1932) The trophic races of *A. maculipennis* disclosed by experimental breeding. *C. R. Acad. Sci.* CXIV, 19, pp 1694-1696 R4FB, XX, p 151
- 1126 ROUBAUD, F (1932) Trophic and biologic variations of *A. maculipennis*. *Bull. Soc. Path. Exot.*, XXV 7 pp 755-762 R4FB XX, p 212
- 1127 ROUBAUD, F (1933) The anopheline standpoint in anophelism and malaria. *Ibid.* XXVI 10, pp 1215-1233 R4EB, XXII, p 50
- 1128 ROUBAUD, F (1937) Principles and possibilities of zoophyllum in malaria control. *Arch. Inst. Past. Tunis* XXVI 4 pp 625-664 R4EB, XXVI, p 65
- 1129 ROUBAUD, E., et al (1933) Data furnished by maxillary index correlated with infectivity of species of Anopheles of Indo-China. *Bull. Soc. Path. Exot.*, XXVI, 2, pp 282-293. R4EB, XXI p 140.

- 1130 ROUBAUD, E. and GASCHEN, H. (1931) Zoophilism of *A. maculipennis* Bull. Soc Path Exot, XXIV, 3, pp 203-209 RAEB, XIV, p 161
- 1131 ROUBAUD, E. and GASCHEN, H (1933) Insufficiency of egg characters to distinguish trophic and biologic races of *A. maculipennis* Ibid, XXVI, 3, pp 447-451 RAEB, XXI, p 151
- 1132 ROUBAUD, E. and TOUMANOFF, C (1935) The maxillary index and trophism of anophelines of the Far East Ibid, XXVIII, 9 pp 835-838 RAEB, XXIV, p 47
- 1133 RUSSELL, P F. (1933) The value of an animal barrier in malaria control Science, IXXVIII, 2014, pp 101-102 RAEB, XXI, p 230
- 1134 RUSSELL, P F (1934) Zoophylaxis failure An experiment in the Philippines Riv Mal, XIII, 5, pp 610-616 RAEB, XXIII, p 59
- 1135 SENIOR WHITE, R A (1940) Studies on the behaviour of adult *A. culicifacies* Part III J. Mal Ind Ind, III, 2-3, pp 363-382
- 1136 SWEET, W C, and RAO, B A (1937) Races of *A. stephensi* Liston, 1901 Ind Med Gaz, LXXII, 11, pp 665-674 RAEB, XXVI, p 49
- 1137 SWELLENGREBEL, N H, and BURK, A DE (1933) Dutch races of *A. maculipennis* and their relation to those of Italy Bull Soc Path Exot, XXVI, 2, pp 273-282 RAEB, XXI, p 139
- 1138 THIEL, P H VAN (1933) Investigations on *A. maculipennis* races and their bearing on malaria in Italy Riv Mal, XII, 2, pp 281-318 RAEB, XXI, p 211
- 1139 THIEL P H VAN (1935) Behaviour of Anopheles towards man and animals Geneesk Tijds Ned Ind, LXXV, 25, pp 2101-2118 RAEB, XXIV, p 89
- 1140 THIEL, P H VAN, et al (1939) Zoophilism and anthropophilism of Anopheles biotypes and species Riv Mal, XVIII, 2, pp 95-124 RAEB, XXVIII, p 10
- 1141 THIEL, P H VAN, and BEVERE, L (1939) Experimental proof of the anthropophilism of *A. maculipennis labranchiae* and *clutus* Bull Soc Path Exot, XXXII, 1, pp 103-109 RAEB, XXVII, p 162
- 1142 THIEL, P H VAN, and SAUTET, J (1937) Existence of anthropophile biotypes of *A. maculipennis* Ibid, XXX, 2, pp 186-193 RAEB, XXV, p 170

- 1143 TOUMANOFF, C. (1934) The trophic habits of anophelines of Hong Kong *Bull Soc Path. Exot.*, XXVII, 8, pp 745-749 *RAEB.*, XXIII, p. 34
- 1144 TOUMANOFF, C (1936) Anophelines of the Far East *Soc Path. Exot. Monograph IV.* 434 pp *RAEB.*, XXV, p 138
- 1145 TOUMANOFF, C, and HU, S M K (1934) First results regarding the zoophilum of *A. hyrcanus* var *sincensis* in China *Bull Soc Path. Exot.*, XXVII, 8, pp 741-745 *RAEB.*, XXIII, p. 33.
- 1146 TOUMANOFF, C, and HU, S M K (1935) Trophism of *A. hyrcanus* var *sincensis* around Shanghai *Ibid.*, XXVIII, 9, pp 832-835 *RAEB.*, XXIV, p 46
- 1147 TRAUTSMILLER O (1935) Malaria in the Adriatic Islands, Krk, Rab and Pag *Bull Off Int Hyg Pub* XXVII, 2 pp 291-303 *RAEB.*, XXIII, p 148
(Cattle stabled between reservoir and village)
- 1148 WALCH F W (1932) On the relation between the preference for animal or human blood of Netherlands Indian Anophelids and their ability to transmit malaria under natural conditions 1st *Brev. Verh. Phynol.*, II pp 206-210 Also in *Geneesk. Tijdschr. Ned. Ind.* LXXII, 11 pp 682-709 *TDB.*, XXXI p 159

*What The Director, Malaria Institute of India, says
about our*

SOFT STONE POWDER

"The Soft Stone Powder sent by you to the Officer in charge, Ross Field Experimental Station for Malaria Karnal, has given encouraging results, and it appears that it will be a suitable diluent for Paris Green

Large stock of the above powder is held by

Calcutta Mineral Supply Co., Ltd.

31, Jackson Lane, Calcutta

*The biggest dealers in Indian Minerals and
Mineral products*

Asbestos Composition
Barytes
China Clay
French Chalk
Fire Brick and Fire Clay
Felspar
Glass Powder
Graphite Powder

Manganese Dioxide
Mica Dust
Plaster of Paris
Quartz Sand
Red and Yellow Ochre
Soap Stone Powder
Silicate of Soda
Talc Powder

For particulars and rates please write to

H. DUTT & CO.

Managing Agents

31, Jackson Lane Calcutta

Telephone Butra Bazar 1397

Telegrams Chmamat "

APPENDIX I

Anti-Mosquito Equipment.

| ARTICLE | WHERE OBTAINABLE |
|------------------------------|---|
| Paris Green Equipment | |
| Haverack | Dina Nath & Sons Gumti Bazaar Lahore |
| Mixer | Dina Nath & Sons Lahore Shaw Wallace & Co 4 Bankshall St Calcutta General Export Co 63 Ezra St Calcutta |
| Paris green | F Klein 1 Royal Exchange Place Calcutta Bishan Das Arora & Sons Bharat Buildings Nisbet Rd Lahore Dina Nath & Sons Lahore General Export Co Calcutta Howie Robson Barnett & Co Ltd Agents |
| Peerless Dust Gun | Shaw Wallace & Co Calcutta |
| Screeners | Shaw Wallace & Co Calcutta Dina Nath & Sons Lahore |
| Soapstone powder | |

Oiling Equipment.

Eclair No. 2 Model Knapsack | McLeod & Co. Post Box 74 Dalhousie St.,
Sprayer (Vermorel Eclair) | Calcutta Messrs Ltd., Main Rd., Poy-
31 cala. | suram Madras.

Anti-Mosquito Equipment—*concl'd*

| ARTICLE | WHERE OBTAINABLE. |
|--|--|
| Oiling Equipment—<i>concl'd</i>. | |
| Four Oaks Ross Knapsack Sprayer, 3½ gals | Planters' Stores and Agency Co, Calcutta |
| Fuel (diesel) oil, Malariol | Burmah Shell Oil Storage and Distributing Co. of India, Ltd., Ambala, Bombay, Calcutta, Delhi, Karachi |

Equipment for Spray-killing Adult Mosquitoes.

| | |
|--------------------|--|
| Deskito, Pye fly | Stafford Allen & Sons, London, Anglo Siam Corporation, Ltd, Ewart House, Bombay |
| Pyrocide 20 | F S Kerr & Co, Vulcan House, Ballard Estate, Bombay. |
| Pyrefume | Penick & Co, 132, Nassau St, New York, U S A; Standard Chemical & Pharmaceutical Co, Bombay |
| Pyrin, Pyrentol 20 | John Powell & Co, 114, East St, New York, U. S. A; Standard Chemical & Pharmaceutical Co, Bombay |
| Shell tox | Burmah Shell Oil Storage and Distributing Co of India, Ltd |
| Power Sprayers | Gallanders Arbuthnot & Co, Connaught Place, New Delhi, Calcutta, Bombay, Madras, Lahore |

Wire Gauze.

| | |
|---|---|
| Wire Gauze Screen (Greening Rustless Bronze Gauze Wire) | Wright & Partners, Ltd, 19, Aurangzeb Rd, New Delhi |
|---|---|

APPENDIX II

Table of Equivalents

Measures of Capacity

| | |
|-------------------------------------|--|
| 1 Cubic centimetre = 0.0352 ounce | 1 Gallon = 4.546 litres |
| 1 Ounce = 28.4123 cubic centimetres | 1 United States gallon = 0.833 Imperial gallon |
| 1 Pint = 568.2454 cubic centimetres | 1 Imperial gallon = 1.2 United States gallons |
| 1 Gallon = 160 ounces | |
| 1 Litre = 0.2199 gallon | |

Measures of Weight

| | |
|------------------------------|---|
| 1 Gramme = 15.432 grains | 1 Fluid drachm = 3.5437 grammes or 54.6873 grains |
| 1 Grain = 0.0648 gramme | 1 Fluid ounce = 437.5 grains or 28.350 grammes |
| 1 Pound = 0.4535 kilogramme | |
| 1 Kilogramme = 2.2046 pounds | |
| 1 Ton = 2,240 pounds | |

Measures of Length and Area.

| | |
|----------------------------|-----------------------------|
| 1 Centimetre = 0.3937 inch | 1 Chain = 66 feet |
| 1 Inch = 2.54 centimetres | 1 Acre = 4,840 square yards |
| 1 Metre = 1.0936 yards | 1 Square mile = 640 acres |
| 1 Yard = 0.9144 metre | 1 Acre = 0.4047 hectare |
| 1 Foot = 30.48 centimetres | 1 Hectare = 2.4711 acres |

Approximate Equivalents

| | |
|---------------------------------------|-------------------------------------|
| 1 Millimetre = $\frac{1}{8}$ inch | 1,000 kilogrammes = 1 ton |
| 1 Centimetre = $\frac{1}{2}$ inch | 1 Litre = $1\frac{1}{2}$ pints |
| 1 Metre = 39 $\frac{1}{2}$ inches | 1 Pound = 453 $\frac{1}{2}$ grammes |
| 1 Foot = 30 centimetres | 5 Kilogrammes = 11 pounds |
| 5 Miles = 8 kilometres | 1 Gramme = 15 grains |
| 1 Kilogramme = 2 $\frac{1}{2}$ pounds | 1 Hundredweight = 50 kilogrammes. |

INDEX

| A M V | I A E | Chemical | 140K |
|-----------------------------|---------|--------------------------|---------|
| Abies | 50 | Chemical Larvicides | 107-109 |
| Aeroplanes for Paris green | 61 | Chlorine gas | 27 |
| Agitation of water | 116 | Chloroform tubes | 11 |
| Alibree | 91 | Cisterns | 82 |
| Alternation of flow | 100 | Cladophora | 83 |
| Animal prophylaxis | 1 | Clayton mach n s | 19 |
| Anti malarial oil | 30 | Cleaning junct | 27 |
| Army Fly spray | 6 | Clematis trap | 33 |
| Asobis | 81 | Clover | 16 |
| Balls oil | 53 | Cold weather control | 27 |
| Bamboo | 4 | Cone of oil | 99 |
| Barrel pump | 53 | Copper sulphate | 20 |
| Bats | 24 | Cotton waste | 77 |
| Baquito extra | 50 | Coin arm | 52 |
| Bellows hand | 63 | Crank case oil | 23 |
| Biological control | 81 | Crocin (as fumigant) | 50 |
| Blowers electric | 118-119 | Crocin (as larvicide) | 22 |
| Blowers knapsack | 64 | Cromote (as larvicide) | 75 |
| Blowers rotary | 63 | Croton (as repellent) | 5 |
| Booms, oil | 63 | Crowl (as fumigant) | 21 |
| Boots, mosquito | 100 | Crowl (as larvicide) | 48 |
| Borax | 3 | Crown line trap | 13 |
| Borrow pits | 91 | Cruel oil | 49 |
| Box traps | 14 | Culicifuges | 3-5 |
| Bug and Van Slooten's spray | 21 | Cunettes | 94 |
| Brushing method of oiling | 103 | Cutting down jungle | 34 |
| Buckets fire | 93 | Cylin | 27 |
| Building water used in | 97 | Datura | 74 |
| Cactus paste | 74 | Depressions in ground | 90 |
| Calco gates | 81 | Derris | 78 |
| Camphor and carbolic acid | 22 | Deterrent aquatic plants | 81 |
| Carbolic acid | 73 | Deterrent trees | 27 |
| Cashew nut shell oil | 110 | Diesel oil | 50 |
| Castor oil | 47 | Divers and cistern | 3, 9 |
| Catching stations | 32 | Drainage | 102-104 |
| Cham | 87 | Drainage, fascine | 102 |
| Chalk traps | 17 | Drainage, Lido system | 103 |
| | | Drainage vertical | 43 |
| | | Drains contour | 33, 83 |

| | PAGE | | PAGE |
|----------------------------|-----------|----------------------------|--------|
| Drains herring bone | 37 | Holes in trees | 91 |
| Drains open | 36 | Household traps | 15 |
| Drains parallel | 37 | Housing and malaria | 101 |
| Drains, subsoil | 39-43 102 | Hycol | 77 |
| Drip cans | 54 | Hydrants | 93 |
| Drying intermittent | 93 116 | Hydrated lime (as diluent) | 59 |
| Duckweeds | 82 | | |
| <i>Duranta</i> | 36 | Illuminated traps | 18 95 |
| | | Impounded water | 87 107 |
| Electric blowers | 64 | Intermittent drying | 93 116 |
| Emprasin | 76 | Intermittent irrigation | 116 |
| <i>Eupatorium</i> | 36 | Irrigation and drainage | 90 |
| | | Irrigation channels | 89 |
| Fans hand | 3 | Irrigation intermittent | 116 |
| Fascine drainage | 102 | Isotex | 76 |
| Fauna and flora changing | 118 | Ival | 22 |
| Ferric hydroxide | 84 | | |
| Fire buckets | 93 | Jack fruit trees | 36 |
| Fish | 79-81 118 | James, J F, trap | 95 |
| Fish ponds control of | 118 | James, S P trap | 17 |
| Fletcher's trap | 13 | Jungle clearing | 27, 35 |
| Flit | 24 | | |
| Flooding | 115 | Katol | 3 |
| Fluctuating water level | 115 | Kerosene (as larvicide) | 49 |
| Flushing | 113 | Kerosene (as spray) | 24 |
| Fly bed | 24 | Key walls | 38 |
| Flyosan | 24 | Knapsack sprayers | 63 107 |
| Flytox | 24 | | |
| Formalin | 22 | Lakes | 87 |
| Formic acid | 22 | Lefroy's spray | 26 |
| Fuel oil | 49 105 | Lefroy's trap | 13 |
| Fumigation | 18-23 95 | <i>Lemna</i> | 81 92 |
| | | Leron | 51 |
| <i>Gambusia</i> | 79 118 | Lido system of drainage | 103 |
| Giemsa's spray | 24 26 | Lime (as diluent) | 59 |
| Giles cones | 20 | Lime (as larvicide) | 75 |
| Gloves | 3 | Liquid fuel | 49 |
| Guddas | 53 | Liquid paraffin | 51 |
| Gumti Weir | 113 | Lysol | 22 |
| Gutters rain water | 93 | | |
| | | M D B | 50 |
| Hand bellows | 63 | Mansell's spray | 25 27 |
| Hand catching | 11 | Malaria | 105 |
| Hand nets | 11 | Marshes | 85 |
| Haversacks for Paris green | 63 | Mercuric chloride | 23 |
| Herbage packing | 112 | 'Millions | 79 |
| Hill foot seepages | 89 | M mm's chloride | 22 |
| Holes in rocks | 91 | Mixer for Paris green | 62 |

| | PAGE | | PAGE |
|-----------------------------------|----------------|------------------------------------|-------------------|
| Monel metal | 9 | Pyrethrum extract (as spray) | 24, 26, 95-100 |
| Mosquit | 24 | Pyrethrum powder . . | 18 |
| Mosquito boots . . . | 3 | Pyrocide | 97 |
| Mosquito nets | 1-3 | | |
| Motor oil, waste . . . | 50 | Rain water gutters . . | 93 |
| Muddying | 116 | Repellents | 3-5, 94 |
| Naturalistic measures . | 111-119 | Research trap | 16 |
| Necrosene, emulsified (larvicide) | 76 | Reservoirs | 87 |
| Necrosene (spray) . . | 24 | Rice-fields | 89, 116 |
| Nets hand | 11 | Richmond and Mendis traps | 13 |
| Nets, mosquito | 1-3 | Rods | 24 |
| Oil, anti malarial . . | 50 | River beds | 88 |
| Oil balls | 53 | Rocks, holes in | 91 |
| Oil, Bamber | 4 | Rotary blowers | 63 |
| Oil booms | 106 | Sal hycol | 77 |
| Oil, crank case | 50 | Salinity, changing of . | 112 |
| Oil, crude | 49, 105 | Salt water swamps . . | 86, 112 |
| Oil, diesel | 50, 105 | Sanitas Okol | 74 |
| Oil, fuel | 49, 105 | Sawdust | 47, 54 |
| Oil, kerosene | 49 | Schweinfurt green . . | 57 |
| Oil, solar | 50, 105 | Screen trap | 17 |
| Oiling | 41-57, 104-107 | Screeners for paris green diluents | 60 |
| Oils, spread of | 46, 105 | Screening | 5-10, 94-95 |
| | | Seepages | 89 |
| Panama Canal larricide . | 74 | Sergent's trap | 15 |
| Paraformaldehyde . . | 72 | Shading | 30, 117 |
| Paris green | 57-71, 107 109 | Shell tox | 24 |
| Paris green, dustless method . | 108 | Sifting | 113 |
| Paristan | 76 | Slaked lime (as diluent) | 59 |
| Pelitory root | 78 | Sludge | 50 |
| Praterine | 50 | Sluice-valves | 93 |
| Petrol | 51, 92 | Sluicing | 113 |
| Pine oil | 51 | Smoke | 23 |
| Pollution of water . . | 111 | Soap | 75 |
| Ponding | 116 | Soapstone | 59 |
| Ponds | 87 | Soft stone | 59 |
| Pools | 91 | Solar oil | 60, 105 |
| Iron Block traps . . . | 15 | Spray, Brug and Van Slooten's | 24, 26 |
| Iyagra | 97 | Spray, Giesman's . . . | 24, 26 |
| Pre-fly | 97 | Spray, Lefroy's | 26 |
| Pyrentol | 97 | Spray, Mansell's . . . | 25 27 |
| Pyrethrum emulsion (as larricide) | 110 | Sprayers, knapsack . . | 63, 107 |
| Pyrethrum emulsion (as repellent) | 94 | Sprayers, oil | 51 |
| | | Sprays | 23-27, 95-100 |
| | | Stagnating | 116 |

| | PAGE | | PAGE |
|----------------------------|-------------|-----------------------------|------------|
| Standp pes | 93 | Traps Strickland and | |
| Stone packing | 103 | Chowdhury s | 17 |
| Storal | 71 | Traps Wilson s | 33 |
| Stramonium | 23 | Traps wire | 15 |
| Streama | 83 | Trees (as screens) | 23 |
| Strickland and Chowdhury s | | Trees deterrent | 27 |
| trap | 17 | Trees holes in | 91 |
| Sub aqueous saws | 88 | Trioxymethylene | 71 |
| Subso l drains | 39 107 | Tubes chloroform | 11 |
| Sulpl ate of copper | 77 | | |
| Sulphur dio x de | 19 | <i>Utricularia</i> | 82 |
| Swamps | 85 | | |
| Swatters | 11 | Vaseline liquid | 51 |
| | | Vegetable larvic des | 78 109 |
| Tanks | 87 | Vegetable o ls | 47 |
| Tobacco | 23 78 | Veils | 3 |
| Trap holes | 18 | Vertical drainage | 43 |
| Trap house McHardy s | 18 | | |
| Traps | 17 18 92-96 | Waste cotton | 51 |
| Traps box | 14 | Waste motor o l | 50 |
| Traps chatta | 17 | Waste tobacco | 23 |
| Traps Clemesha s | 16 | Water hyac nth | 87 |
| Traps crinol ne | 13 | Water impounded | 107 |
| Traps Fletcher s | 13 | Water logging | 90 |
| Traps household | 15 | Watering pots for oil | 52 |
| Traps illuminated | 18 90 | Weep holes | 38 |
| Traps J F James | 95 | Wells | 91 |
| Traps S P James | 17 | Wetmore s apparatus for o l | 54 |
| Traps Lefroy s | 13 | Whiz | 24 |
| Traps Proudlock s | 15 | Wilson s trap | 33 |
| Traps research | 16 | Wire traps | 15 |
| Traps Richmond and Mend s | 13 | <i>Wolffia</i> | 85 92 |
| Traps screen | 17 | | |
| Traps Sergeant s | 15 | Zooprophylaxis | 29 100-101 |

"VERMOREL-ECLAIR"


SPRAYING EQUIPMENT

"SEM"

PNEUMATIC KNAPSACK SPRAYERS

HIGHLY
EFFICIENT
PRESSURE
RETAINING
MODEL

ALL LIQUID
CONTENTS
ARE SPRAYED



SUITABLE
FOR ALL
LIQUIDS
EXCEPT
ACIDS

BOTH HANDS
FREE DURING
SPRAYING

Also available—

The "Eclair" Model No. 2 Knapsack Sprayer
The "Eclair" Knapsack Limewashing Machine
The "Noe" Pneumatic Knapsack Sprayer
The "Vermorel" Pneumatic Hand Sprayer
The "Hydra" Self-Supplying Syringe
The "Salva" Bucket Sprayer, etc., etc.

MCLEOD & COMPANY, LIMITED, CALCUTTA

RUSTLESS BRONZE GAUZE MOSQUITO PROOF SCREENING

Large stocks held in India

As standardized by

ARMY HEADQUARTERS, INDIA

AND SUPPLIED TO

THE INDIAN STORES DEPARTMENT

Manufactured by

THE B. GREENING WIRE CO., LTD.

HAMILTON, ONTARIO, CANADA

Sole Agents for India, Burmah and Ceylon :

R. WRIGHT & PARTNERS, LTD.

19, AURANGZEB ROAD, NEW DELHI

Everything Required for

MALARIA SURVEY WORK

and all sorts of apparatus and appliances for the study of Malaria Mosquitoes, such as Entomological Pins, Cork Sheets, Insect Store Boxes, Larvæ Collection Pipettes, Portable Breeding Cages, Lenses and Magnifiers, Cover Slips, Slides, Slide Boxes, Microscopes, etc., are available from us. We can also supply Health Charts, Lanterns and Lantern Slides required for Health Lectures

The largest and most varied stock of scientific apparatus and chemicals in India

SCIENTIFIC SUPPLIES (BENGAL) CO.

C 37 & 38, College Street Market, CALCUTTA


Phone B B 524

Telegrams "BITISYND", Calcutta

Plants Protection & Anti-Malaria Campaign

DESTROY THE MOSQUITOES AND ALSO PROTECT YOUR PLANTS AGAINST ALL PREVAILING DISEASES BY THE USE OF OUR MOST DURABLE AND CHEAP APPARATUS

Manufacturers of
Paris Green
Mixers,
Screeners,
Hand
Bellows
Haversacks
etc
Approved
by D.D.M.S.
Northern
Command
and others



STOCKISTS
OF
PARIS GREEN
AND
ANTI-
MALARIA
STORES

Dust-Gun
For Blowing Paris Green Powder

Manufactured by:—

Messrs. DINA NATH & SONS

GOVERNMENT AND MILITARY CONTRACTORS AND ORDER
SUPPLIERS

GUMTI BAZAR, LAHORE

Pyefly

For Sprays that Kill!

PYEFLY is an Empire product a concentrated Liquid Extract of Pyrethrum manufactured in England from Empire grown flowers. It is biologically standardised to a specific strength so that when 1 part is diluted with 64 parts of kerosene a highly toxic insecticide is produced. Biological tests have proved that PYEFLY at this dilution gives a 100% efficient spray for use against flies etc. Much has been written and possibly will be written on the comparative values of chemical and biological tests. Both are necessary but it is the biological test which proves its actual toxicity.

The toxic principles Pyrethrins being organic bodies are prone to decomposition, but certain precautions introduced into our method of manufacture ensure their stability.

The Malaria scourge is sufficiently well known that comment is unnecessary so also is the fact that where adequate anti-mosquito measures have been applied there has been a marked decrease in Malaria fever. Until such time as all mosquito breeding places are exterminated Malaria fever will continue, unless the adult mosquitoes are destroyed. Proof that Pyrethrum sprays if properly used will do this is already known, thanks to the experiments carried out by various authorities in India.

The reason why PYEFLY is strongly recommended as the toxic principle of fly and mosquito sprays is that it has a guaranteed stabilised strength calculated by biological tests.

Besides mosquitoes there are other bacteria carrying insects and flies etc. which can be controlled with Pyrethrum sprays of the right sort if used thoroughly.

Formula —

Pyefly
Citronella oil
Kerosene

1 part
4 parts
59 parts

STAFFORD ALLEN & SONS LIMITED
Wharf Road, London, E.C. 2
England

Agents — **THE ANGLO-THAI CORPORATION, LTD.**
P. O. Box No 70, BOMBAY



Pest Fighters

OF THE NATION...

MYERS SPRAY PUMPS.

Highly efficient for all liquid spraying
and suitable for Anti-Malaria spraying.

Sole Agents

**THE WATER SUPPLY
SPECIALISTS LTD.**

MISSION ROW EXTENSION

(GUJRAT MANSION)

CALCUTTA

Phone: Cal. 4129. Grams: WATERFLOW.

Post Box 424.

ASK FOR CATALOGUE

PUMPS FOR **EVERY PURPOSE**

SAVE A LOT BY BUYING

"PYRENTOL-20"

Biologically and chemically standardised Extract of Pyrethrum Flowers
in its new form. It is a perfect and stable base for insecticidal sprays,
mosquitocidal and larvicidal mixtures.

(Approved and recommended by The Malaria Institute of India, Delhi,
as per Government of India Health Bulletin No 11, Fifth Edition-1940)

Chief features :—

It is cheapest in prices

It can be available in higher concentration up to 1 to 79.

It does not weaken in quality for a long time storage

Available from our own agents and stockists at Calcutta, Madras,
Karachi, Lahore, Delhi, Rangoon, Colombo, etc., etc

For further particulars and prices, please write to —

STANDARD CHEMICAL & PHARMACEUTICAL CO.

Atlas Mills, Reay Road, Bombay No. 10.

Telegraphic Address: "GERMOL."

Telephone: 46612.

Chemotherapy of Malaria

PRAEQUINE

THE SYNTHETIC GAMETOCIDE



QUINACRINE

THE SYN

MAY & BAKER have pleasure in announcing
they have undertaken the manufacture of this
essential anti malarial preparation

PRAEQUINE

is supplied in tablets of
0.01 gm and 0.05 gm and in
ampoules of 1 per cent solution

QUINACRINE

is supplied in
0.10 gm
containing

* T.M. & B.

MAY & BAKER (INDIA) LTD.
11 CLIVE STREET, CALCUTTA



PYROCIDE 20

STANDARDIZED EXTRACT OF PYRETHRUM FLOWERS

FOR

LARGE SCALE HOUSE SPRAYING SCHEMES
PYROCIDE 20 IS THE IDEAL BASE AND AT
THE SAME TIME THE CHEAPEST.

"Results go to show that an insecticidal mixture composed of one part of Pyrocide 20 in nineteen parts of Kerosene Oil will produce an insecticide which is equal in its mosquitocidal efficacy to the best proprietary sprays at present available on the market in India, and at about half the cost."
Vide September 1935 issue of the Records of the Malaria Survey of India.

SOLE AGENTS for :

INDIA, BURMAH and CEYLON

F. S. KERR & COMPANY


125, ESPLANADE ROAD
BOMBAY

Distributors for :

Bengal, Assam, Bihar and
Orissa
McGregor & Balfour Ltd.
11, Clive Street
Calcutta

Distributors for

Madras and Southern India
Vernon & Co.
36, Second Line Beach
Madras



MALARIA CONTROL BY ANTI-MOSQUITO MEASURES

DeVilbiss Equipment for Spraying
Insecticides has been approved by
the Malaria Institute of India and
initial orders have been placed
by the Government of India

Electric and Petrol driven Units
ranging from $\frac{1}{4}$ HP upwards are
in stock for immediate delivery

JOHN FLEMING & CO., LTD.

Bastion Road, BOMBAY

PYROCIDE 20

STANDARD INSECTICIDE OF THE FUTURE

FOR

LARGE SCALE HOUSE SPRAYING SCHEMES
PYROCIDE 20 IS THE IDEAL BASE AND AT
THE SAME TIME THE CHEAPEST

'Results go to show that an insecticidal mixture composed of one part of Pyrocide 20 in nineteen parts of Kerosene Oil will produce an insecticide which is equal in its mosquitocidal efficacy to the best proprietary sprays at present available on the market in India and at about half the cost
Vide September 1935 issue of the Records of the Malaria Survey of India

SOLE AGENTS for

INDIA, BURMAH and CEYLON

F. S. KERR & COMPANY


125 ESPLANADE ROAD
BOMBAY

Distributors for

Bengal Assam Bihar and
Orissa
McGregor & Balfour Ltd
11, Clive Street
Calcutta

Distributors for

Madras and Southern India
Vernon & Co.
36 Second Line Beach
Madras



MALARIA CONTROL BY ANTI-MOSQUITO MEASURES

DeVilbiss Equipment for Spraying Insecticides has been approved by the Malaria Institute of India and initial orders have been placed by the Government of India.

Electric and Petrol driven Units ranging from 1 H.P. upwards are in stock for immediate delivery.

JOHN FLEMING & CO., LTD.

Bastion Road, BOMBAY